

Parameter Approximations for Quantile Regressions with Measurement Error

ANDREW CHESHER

CENTRE FOR MICRODATA METHODS AND PRACTICE
INSTITUTE FOR FISCAL STUDIES AND UNIVERSITY COLLEGE LONDON

July 21st 2001

ABSTRACT. The impact of covariate measurement error on quantile regression functions is investigated using a small variance approximation. The approximation shows how the error contaminated and error free quantile regression functions are related, a key factor being the distribution of the error free covariate. Exact calculations probe the accuracy of the approximation. The order of the approximation error is unchanged if the error free covariate density is replaced by the error contaminated density. It is then possible to use the approximation to investigate the sensitivity of estimates to varying amounts of measurement error.

KEYWORDS: measurement error, parameter approximations, quantile regression.

1. INTRODUCTION

Quantile regression functions are of particular interest in problems where the dispersion or the shape of conditional distributions are expected to vary with covariate values. Since the seminal paper of Koenker and Bassett (1978), which introduced a method for estimation of quantile regression functions, there has been substantial development of estimation methods and algorithms, and gains in understanding of the properties of estimators. With increasing interest in econometrics in variation in response across individuals, and with the way in which the distribution of responses is affected by covariates, quantile regression estimation procedures are finding increasing use in applied econometric work.

Recent applications of quantile regression methods are: in finance, Bassett and Chen (2001), Chernozukov and Umantsev (2001), Taylor (2000); in analysis of the dynamics and distribution of wages and earnings, Bushinsky (2001, 1998a, 1998b, 1995), Eide and Showalter (1999), Fitzenberger, Hujer, McCurdy and Schnabel (2001), Reilly (1999); in analysis of the effect of, and achievement in, schooling, e.g., Arias, Hallock and Sosa-Escudero (2001), Hartog, Pereira and Vieira (2001), Levin (2001); in studies of nutrition, e.g., Variyam, Blaylock and Smallwood (1999); in economic demography, Abravaya (2001); in evaluation of treatment effects, Abadie, Angrist and Imbens (2001), Koenker and Biliias (2001); and in industrial economics, Machado and Mata (2000), Nahm (2001).

The conditional τ -quantile regression function (τ -QRF) for a scalar random variable Y given covariates X is defined implicitly by¹

$$F_{Y|X}(Q_{Y|X}(\tau, x)|x) = \tau$$

where

$$F_{Y|X}(y|x) = P[Y \leq y|X = x]$$

¹To keep discussion simple we assume throughout that conditional distribution functions are proper and strictly increasing in y .

is the conditional distribution function. The continuum of τ -quantile regression functions, $\tau \in (0, 1)$, captures all aspects of the dependence of conditional distribution of a response Y on covariates, X , unlike conditional moment regression functions. An attractive feature of QRFs is that they allow the nature of the dependence to be easily appreciated because they are expressed in the same metric as the response².

Despite the advances in QRF estimation and inference methods, and the many applications of quantile regression methods, some of the econometric issues given substantial attention in the study of mean regressions have received scant attention in the context of quantile regression estimation. One of these is *covariate measurement error*, a pervasive feature of econometric data, and likely a feature of many of the applications listed above.

Covariate measurement error causes many and subtle changes to conditional distributions, potentially attenuating mean regression functions, increasing dispersion, introducing heteroskedasticity in homoskedastic error free models and modifying the form of heteroskedasticity when it is present in an error free model.

There seems to have been little attention paid to measurement error in the context of quantile regression. Measurement error oriented texts such as Fuller (1987) and Carroll, Ruppert and Stefanski (1995) have no discussion of quantile regression. In the quantile regression literature Brown (1982) and He and Liang (2000) provide the few results available on estimation of parameters of QRFs when there is covariate measurement error. Brown (1982), studying robust estimation in errors-in-variables models, proposes a modified LAD estimator which can be regarded as an estimator of the slope of the median regression function with scalar error contaminated covariate, but rejects the estimator as unsatisfactory. He and Liang (2000) propose a consistent estimator of the slope of linear error free QRFs based on minimising the sum of the “check” functions

$$\rho_\tau(r) = r \times (\tau - 1_{[r < 0]})$$

applied to *orthogonal* residuals, r . They assume that the joint distribution of the response error and the covariate measurement error is spherically symmetric and independent of X .

This paper considers error free QRFs for a response Y , conditioned on X , and error contaminated QRFs for Y , conditioned on $Z = X + V$ where V is distributed independently of X , and of Y given X . Data on Y and the error contaminated Z provide information about the way in which the conditional quantiles of Y given Z vary with Z . Nonparametric quantile regression methods (see e.g., Yu and Jones (1990), Magee, Burbidge and Robb (1991), Nahm (2001)) can give detailed information about this variation. But this bears only indirectly on the way in which quantiles of Y conditional on error free X vary with X . In most applications this is what is of interest because economic theory will be informative about relationships between error free variates, and policy interventions, whose impact on the distribution of Y is of interest, will alter values of error free covariates.

The first part of this paper aims at improving understanding of the relationship between error contaminated and error free QRFs. This can help in interpreting the results of QRF estimation when measurement error is believed to be present. It can help explain differences in estimated QRFs using data sets with different amounts of measurement error. In cases where a functional form of error free QRFs is imposed upon error contaminated data it illuminates the misspecification that is thereby committed.

In the second part of the paper we consider problems in which error free QRFs are parametrically specified (e.g., as linear), and investigate the possibility of using information on the relationship between error free and error contaminated QRFs to retrieve

²Bushinsky (1998a) provides a good introduction to quantile regression methods and their implementation.

information about the values of parameters of error free QRFs. In some circumstances this is not possible because the error free QRF cannot be identified from knowledge of the form of error free QRFs because measurement error induces no change in that form. The model in which Y , X and V are jointly normally distributed is a leading example. Here measurement error changes the separation and slope of QRFs but they remain linear.

But in many other cases identification is possible, as pointed out in Riersøl (1950). However such identification, flowing from functional form assumptions, is fragile. For that reason the procedure we propose is presented mainly as a means of providing a form of sensitivity analysis. It provides a partial answer to the following question.

Were the error free QRF to be of a hypothesised form and covariate measurement error to be present, what are the likely values of the parameters of the error free QRFs, and how does our view of this change as the amount of measurement error allowed for increases?

An exact answer to this question requires a case by case analysis. The exact impact of measurement error on mean regressions can be derived in explicit form only in a few special cases³. Outside these cases, and in almost all interesting cases⁴ for QRFs, the exact impact of covariate measurement error can only be obtained by numerical calculation. Such calculations do not give insight into the generic effects of covariate measurement error and they do not provide a link between the effects of measurement error and easily grasped features of the error free QRF and the distributions of covariates and measurement error.

This paper provides a partial resolution of this problem by providing an *approximation* to an error contaminated QRF expressed in terms of functionals of the error free QRF and the density of either the error free or the error contaminated covariates. The approximation is developed using small parameter (variance) approximation methods and extends the results of Chesher (1991) to QRFs.

Section 2 gives the approximation to error contaminated QRFs. Details of the derivation are given in an appendix. The insights into the generic effects of measurement error on QRFs provided by the approximation are discussed in Section 3 where some leading special cases are examined.

Section 4 reports an investigation into the accuracy of the approximation in a rich class of models with a single error contaminated covariate. An error free covariate (X) and independently distributed measurement error (V) are given exponential power distributions⁵. The conditional distribution of the response given X (independent of V) is also specified as a member of the exponential power family with location parameter depending upon X and with scale and shape parameters independent of X . We use numerical methods to calculate the exact error free (conditional on X) and error contaminated (conditional on $Z = X + V$) QRFs and we calculate the approximation developed in Section 2. We find that for quite large amounts of measurement error the approximations are encouragingly accurate.

Section 5 considers one possible use of the approximation. It investigates the use of the approximate QRF to extract information about the error free QRF from error contaminated data. The results of Section 2 show that the approximate error contaminated QRF is determined by the error free QRF and derivatives of it, whose form is known once the

³A leading example is the model in which Y (with arbitrary distribution) has *polynomial* regression on error free X and additive independent measurement error is normally distributed, see Chesher (1998a).

⁴The fully Gaussian model in which Y has linear regression on X and Y , X and V are jointly normally distributed is the obvious exception.

⁵A random variable with an exponential power distribution has density function proportional to $\exp(-\lambda|w - \mu|^{\frac{2}{1+\gamma}})$. Normal ($\gamma = 0$), Laplace ($\gamma = 1$) and uniform ($\gamma = -1$) distributions are leading special cases.

error free QRF is specified, and by a functional of the density of the error contaminated covariate. This density can be estimated using realisations of the error contaminated covariate. Therefore, given a parametric form for the error free QRF, a parametric approximate error contaminated QRF can be specified and estimated. When identification permits, estimates of parameters of the error free QRF can be retrieved.

The performance of this procedure is investigated in Monte Carlo experiments. In practice, given the identification issues which arise here one would likely want to use this procedure to perform a sensitivity analysis. Thus one can ask: given a specified form for an error free QRF, how are my views about its parameters changed as I consider the possibility of there being varying amounts of measurement error. Section 6 concludes.

2. THE APPROXIMATE EFFECT OF MEASUREMENT ERROR

2.1. Error free and error contaminated QRFs. Consider a scalar response Y , continuously distributed given k element vector X , and let $F_{Y|X}(y|x)$ be the conditional distribution function of Y given $X = x$, as follows.

$$F_{Y|X}(y|x) = P[Y \leq y|X = x]$$

Let $Z = X + V$ where $V = \Psi U$, U and X are independently distributed and $E[U] = 0$, $Var[U] = I$. The matrix Ψ is lower triangular and $\Psi\Psi' = \Sigma$ so that $Var[V] = \Sigma$.

The conditional τ -QRFs, $Q_X(\tau, x)$, in which conditioning is on *error free* X , and $Q_Z(\tau, z)$, in which conditioning is on *error contaminated* Z , are defined by the following implicit equations.

$$\begin{aligned} F_{Y|X}(Q_X(\tau, x)|x) &= \tau \\ F_{Y|Z}(Q_Z(\tau, z)|z) &= \tau \end{aligned}$$

2.2. Approximate error contaminated QRFs. We seek an approximation to the error contaminated τ -QRF, $Q_Z(\tau, z)$. This is a functional of the conditional distribution function of Y given X and the marginal distribution functions of U and X and depends upon τ and Σ , a relationship we express as follows.

$$Q_Z(\tau, \cdot) = \mathcal{F}(F_{Y|X}, F_X, F_U; \tau, \Sigma)$$

Note that the error free QRF is got by setting $\Sigma = 0$.

$$Q_X(\tau, \cdot) = \mathcal{F}(F_{Y|X}, F_X, F_U; \tau, 0)$$

The approximation to the error contaminated QRF is given in equation (2) below, to which those not interested in the method of derivation can proceed directly.

The approximation is obtained by considering a Taylor series type approximation to $Q_Z(\tau, \cdot)$ as follows⁶.

$$Q_Z(\tau, \cdot) = Q_X(\tau, \cdot) + \sum_{i,j} \sigma_{ij} \frac{\partial}{\partial \sigma_{ij}} Q_Z(\tau, \cdot)|_{\Sigma=0} + o(\Sigma)$$

⁶Here and later unless indicated $\sum_{i,j}$ indicates double summation over i and j both from 1 to k . A term described as $o(\Sigma)$ has the property that

$$\lim_{\tau \rightarrow \omega} \frac{o(\Sigma)}{\omega} = 0$$

where $\omega = \text{trace}(\Sigma)$.

We use the following approximation (Chesher (1991)) to the conditional distribution function $F_{Y|Z}(y|z)$ in which conditioning is on error contaminated Z .

$$\begin{aligned} F_{Y|Z}(y|z) &= \tilde{F}_{Y|Z}(y|z) + o(\Sigma) \\ \tilde{F}_{Y|Z}(y|z) &= F_{Y|X}(y|z) + \sum_{i,j} \sigma_{ij} \left(F_{Y|X}^i(y|z) g_X^j(z) + \frac{1}{2} F_{Y|X}^{ij}(y|z) \right) \end{aligned} \quad (1)$$

Here superscripts i, j indicate differentiation with respect to the i th and j th conditioning arguments, for example

$$F_{Y|X}^{ij}(y|z) = \frac{\partial^2}{\partial x_i \partial x_j} F_{Y|X}(y|x) \Big|_{x=z}.$$

The function $g_X(\cdot)$, which plays a key role in what follows, is the log probability density function of X ,

$$g_X(z) = \log f_X(x)$$

with derivatives as follows.

$$g_X^j(z) = \frac{\partial}{\partial x_j} g_X(x) \Big|_{x=z}$$

For the approximation to have an error of the stated order we require the absolute third own and cross moments of U to be finite valued, the existence of bounded third own and cross derivatives of $F_{Y|X}(y|x)$ with respect to x , and that X has a continuous distribution with twice differentiable density function and support on \mathfrak{R}^k . This approximation to the distribution function does not require Y to be continuously distributed⁷ given X .

For the moment let Q_Z be shorthand for $Q_Z(\tau, z)$. Since $F_{Y|Z}(Q_Z|z) = \tau$ we have

$$\tilde{F}_{Y|Z}(Q_Z|z) = \tau + o(\Sigma),$$

that is:

$$F_{Y|X}(Q_Z|z) + \sum_{i,j} \sigma_{ij} \left(F_{Y|X}^i(Q_Z|z) g_X^j(z) + \frac{1}{2} F_{Y|X}^{ij}(Q_Z|z) \right) = \tau + o(\Sigma).$$

Considering variation in Q_Z and Σ and taking differentials gives

$$F_{Y|X}^Y(Q_Z|z) dQ_Z + \sum_{i,j} d\sigma_{ij} \left(F_{Y|X}^i(Q_Z|z) g_X^j(z) + \frac{1}{2} F_{Y|X}^{ij}(Q_Z|z) \right) + O(\Sigma) = \tau + o(\Sigma),$$

where the superscript “Y” denotes differentiation with respect to the response variable, that is:

$$F_{Y|X}^Y(Q_Z|z) = \frac{\partial}{\partial y} F_{Y|X}(y|z) \Big|_{y=Q_Z} = f_{Y|X}(Q_Z|z)$$

which is the conditional density of Y at the τ -quantile under consideration. Setting $\Sigma = 0$, yields the required derivatives,

$$\frac{\partial Q_Z}{\partial \sigma_{ij}} \Big|_{\Sigma=0} = - \frac{F_{Y|X}^i(Q_Z|z) g_X^j(z) + \frac{1}{2} F_{Y|X}^{ij}(Q_Z|z)}{F_{Y|X}^Y(Q_Z|z)}$$

and we therefore have the following approximation.

$$Q_Z(\tau, z) = Q_X(\tau, z) - \sum_{i,j} \sigma_{ij} \frac{F_{Y|X}^i(Q_Z|z) g_X^j(z) + \frac{1}{2} F_{Y|X}^{ij}(Q_Z|z)}{F_{Y|X}^Y(Q_Z|z)} + o(\Sigma)$$

⁷In its application to QRFs we do assume a continuous distribution for Y with strictly increasing distribution function mainly in order to avoid indeterminacy in QRFs.

The approximation is much more easily interpreted, and used, when expressed in terms of the error free QRF and its derivatives

$$\begin{aligned} Q_X^\tau(\tau, z) &= \frac{\partial}{\partial \tau} Q_X(\tau, z) \\ Q_X^i(\tau, z) &= \left. \frac{\partial}{\partial x_i} Q_X(\tau, x) \right|_{x=z} \end{aligned}$$

and so forth, as follows. Details of the derivation of this expression are given in Appendix 1.

$$\begin{aligned} Q_Z(\tau, z) &= Q_X(\tau, z) + \sum_{i,j} \sigma_{ij} \left(Q_X^i(\tau, z) g_X^j(z) + \frac{1}{2} Q_X^{ij}(\tau, z) \right) \\ &\quad - \frac{1}{2} \frac{1}{Q_X^\tau(\tau, z)} \sum_{i,j} \sigma_{ij} \left(Q_X^{\tau i}(\tau, z) Q_X^j(\tau, z) + Q_X^{\tau j}(\tau, z) Q_X^i(\tau, z) \right) \\ &\quad + \frac{1}{2} \frac{Q_X^{\tau\tau}(\tau, z)}{Q_X^\tau(\tau, z)^2} \sum_{i,j} \sigma_{ij} Q_X^i(\tau, z) Q_X^j(\tau, z) + o(\Sigma) \end{aligned} \quad (2)$$

2.3. Discussion. Section 3 provides interpretation of the terms in this approximation and considers some leading special cases. First it is worth noting that there are elements of generality that may not be obvious at first sight.

Non-additive measurement error. The approximation has been developed for the case of additive measurement error, but we have allowed the error free QRF to be nonlinear, so some other interesting cases can be easily obtained by considering transformations of the covariates. For example⁸ consider a scalar covariate X and let

$$Z = \lambda^{-1}(\lambda(X) + \lambda(V))$$

where $\lambda(\cdot)$ is a strictly monotonic function. Additive and multiplicative measurement error arise when $\lambda(\cdot)$ is respectively the identity function and the logarithmic function. The approximation (2) for additive measurement error applies when the error free QRF is expressed as a function of $\lambda(X)$. Then $g_X(\cdot)$ must be regarded as the log density of $\lambda(X)$. The result is easily “unbundled” to give an approximation in terms of an error free QRF written as a function of X and the log density of X . Of course the resulting approximation will involve the function $\lambda(\cdot)$ and its derivatives⁹.

Error free covariates. We have proceeded as if all elements of X are error contaminated, but in many leading cases of interest we may expect measurement error to be a serious issue for only one covariate. For example in considering household demand we may be confident in the accuracy of measures of household composition but suspect measurement error in household income. The approximation (2) is easily applied to such cases by setting elements of Σ to zero. Note that in this case, with X_F and X_C denoting respectively error free and error contaminated covariates, the log density derivative $g_X^j(z)$ that appears in (2) becomes the derivative of the log *conditional* density of X_C given X_F with respect to elements of X_C .

⁸I am grateful to Christian Schluter for suggesting this generalised additive formulation.

⁹This is essentially the approach taken in Chesher and Schluter (2001) and in Chesher, Dumangane and Smith (2001) in studying the impact of measurement error on respectively inequality measures (e.g., the Gini coefficient) and duration analysis. In both cases multiplicative measurement error is the leading case of interest.

Temporal variation. The approximation has been developed for classical measurement error but it can be applied (with care) to other problems. For example in demand analysis we might regard recorded income as the sum of permanent and transitory income and be interested in the dependence of demand on permanent income. Then, under suitable assumptions, the approximation gives information about the relationship between QRFs for permanent income and QRFs for income accruing over a short recording period.

Alternative forms of the approximation. The log density derivatives $g_X^j(z)$ that appear in (2) can be replaced by derivatives of the log density of Z , $g_Z^j(z)$, without increasing the order of the approximation error. This is proved in Appendix 2. This substitution has two benefits. First, in models with normal measurement error it can result in increased accuracy¹⁰. Second, unlike $g_X^j(z)$, $g_Z^j(z)$ can be estimated - using realisations of error contaminated Z . With an estimate of $g_Z^j(z)$ and knowledge of the form of the error free QRF one then has information about all aspects of the dependence on z of the (approximate) error contaminated QRF, a point that is crucial to our proposed sensitivity analysis procedure.

3. INTERPRETATION AND SPECIAL CASES

To start, it is interesting to compare the quantile regression approximation (2) with the approximate mean regression function given in Chesher (1991). For error free and error contaminated mean regression functions respectively $R_X(x) = E_{Y|X}[Y|X = x]$ and $R_Z(z) = E_{Y|Z}[Y|Z = z]$ this approximation is as follows.

$$R_Z(z) = R_X(z) + \sum_{i,j} \sigma_{ij} \left(R_X^i(z) g_X^j(z) + \frac{1}{2} R_X^{ij}(z) \right) + o(\Sigma) \quad (3)$$

This has the same form as the first line of (2)¹¹.

The second and third lines in (2) capture (approximately) the variance and distributional shape distortions produced by measurement error. Most of the message contained in these approximations can be uncovered by considering the case in which there is just one covariate.

3.1. Attenuation and curve damping. Let superscript “ x ” denote differentiation with respect to the single covariate and write the scalar measurement error variance as σ^2 . When there is one covariate (2) simplifies as follows.

$$\begin{aligned} Q_Z(\tau, z) &= Q_X(\tau, z) + \sigma^2 Q_X^x(\tau, z) g_X^x(z) + \frac{\sigma^2}{2} Q_X^{xx}(\tau, z) \\ &\quad - \sigma^2 \frac{Q_X^{\tau x}(\tau, z) Q_X^x(\tau, z)}{Q_X^\tau(\tau, z)} \\ &\quad + \frac{\sigma^2}{2} \frac{Q_X^{\tau\tau}(\tau, z) Q_X^x(\tau, z)^2}{Q_X^\tau(\tau, z)^2} + o(\sigma^2) \end{aligned} \quad (4)$$

The leading term is just the error free quantile function with argument z . The next two terms completing the first line of (4) do not involve derivatives with respect to τ . These are QRF analogues of the only $O(\Sigma)$ terms in the mean regression approximation (3).

¹⁰When error free mean regressions are linear a substitution of this sort renders the approximation exact, Chesher (1998a).

¹¹It also has the same form as (1) because the conditional distribution function $F_{Y|X}(y|x)$ is a regression function, namely for $1_{\{Y \leq y\}}$ given X .

Note that $g_X^x(z)$ is zero at every mode of the density of X . To the left (right) of each mode $g_X^x(z)$ is positive (negative). Consider x and τ where the error free QRF has a positive derivative. There the effect of the term $\sigma^2 Q_X^x(\tau, z) g_X^x(z)$ is to raise the error contaminated QRF relative to the error free QRF to the left of each mode of the density of X and to lower it to the right of each mode. This tends to “flatten” the QRF and is an expression of the *attenuating effect* of measurement error. There is the same attenuation effect where the error free QRF has a negative derivative. The effect is clear to see when the error free QRF is linear and is illustrated for mean regression in Chesher (1991). Then $Q_X^x(\tau, z)$ is constant and the term $\frac{\sigma^2}{2} Q_X^{xx}(\tau, z)$ vanishes. When $g_X^x(z)$ is also linear the approximate error contaminated QRF is then linear, but otherwise the term $g_X^x(z)$ introduces *nonlinearity*. Of course $g_X^x(z)$ is linear only if X has a normal distribution.

The opposite effect occurs at each antimode of the density of X . Near antimodes the error contaminated QRF is amplified. The result is that when the distribution of X is multimodal the error contaminated QRF tends to move sinuously relative to the error free QRF.

The final term in the first line of (4) is present only when the error free QRF is nonlinear. It is positive (negative) where that QRF is strictly concave (convex). The effect of this term is to dampen the curvature of the error contaminated QRF relative to the error free QRF.

The terms in the second and third lines of (4) are more complex and more easily understood in special cases. We first consider them in problems in which error free QRFs are parallel.

3.2. Parallel conditional quantiles.

Consider parallel error free QRFs

$$Q_X(\tau, x) = a(\tau) + b(x)$$

which arise when Y is a location shift of a random variable W , the latter distributed independently of X , that is

$$Y = b(X) + W.$$

With $Q_W(\tau) = a(\tau)$ denoting the τ -quantile of W ,

$$Q_X(\tau, x) = Q_W(\tau) + b(x).$$

In this case $Q_X^{\tau x}(\tau, z) = 0$ which *removes* the term in the second line of (4).

In this case, applying (4), the error contaminated quantile is approximately

$$Q_Z(\tau, z) = a(\tau) + b(z) + \sigma^2 b^x(z) g_X^x(z) + \frac{\sigma^2}{2} b^{xx}(z) + \frac{\sigma^2}{2} \frac{a^{\tau\tau}(\tau) b^x(z)^2}{a^\tau(\tau)^2} + o(\sigma^2) \quad (5)$$

where superscripts “ x ” and “ τ ” denote differentiation with respect to x and τ respectively. The following points are of interest¹².

1. Even though the error free quantiles are parallel, the error contaminated quantiles are *not* in general parallel, because in the final term of (5) there are functions of z and τ which interact.
2. However if the error free quantile functions are *linear* the final term in (5) is a function of τ alone and measurement error does *not* destroy the parallel quantile property, though it may render quantile functions non-linear through the impact of the term $\sigma^2 b^x(z) g_X^x(z)$ in (5).

¹²Where statements are made about some manifestation of measurement error being present or absent it should be taken to mean to the order of approximation considered in this analysis.

3. If $b(z)$ does not depend upon z then measurement error has no impact - this is, of course, an exact result!
4. Regarding $a(\tau)$ as the quantile function of the random variable W , we have

$$\frac{a^{\tau\tau}(\tau)}{a\tau(\tau)^2} = \frac{Q_W^{\tau\tau}(\tau)}{Q_W^\tau(\tau)^2} = - \left. \frac{\partial}{\partial w} \log f_W(w) \right|_{w=Q_W(\tau)} = -g_W^w(Q_W(\tau))$$

where (A1.3) and (A1.5) of Appendix 1 have been used to obtain the final expression and $f_W(w)$ is the density function of W .

- (a) This term, and so the final term in (5), is zero at each mode (and antimode) of the density of W .
- (b) When the density of W is unimodal, this final term in (5) is negative for small τ and positive for large τ , and captures the impact of measurement error in increasing the dispersion of the conditional distribution of Y .
- (c) This dispersion increasing effect is larger for values of z at which $b^x(z)$ is large in magnitude and zero when $b^x(z)$ is zero. In the nonlinear quantile function case the variations with z in the sensitivity of $b(z)$ to z induce heteroskedasticity.

In summary, parallel nonlinear quantile regressions contaminated by measurement error become non-parallel, the effect being greater at covariate values at which error free QRFs are more nonlinear. The discussion of Section 2.3 implies that this effect will also be present in linear error free QRF problems when measurement error is not additive.

Error contaminated QRFs tend to be more widely separated than error free QRFs. This expansion effect is larger when the slope of the error free QRF is large in magnitude. It is larger for τ -QRFs for which τ corresponds to a quantile on an sharply increasing or decreasing part of the conditional density, in many cases away from the mode of this distribution but in the main body of the distribution.

Non-parallel conditional quantiles. With non-parallel quantiles there is heteroskedasticity and/or conditional shape variation in the error free model and these are altered by the introduction of measurement error. This effect is captured in the term in (4) involving $Q_X^{\tau x}(\tau, z)$ which is nonzero only at points where quantile functions are non-parallel. Consider the simple case in which

$$Q_X(\tau, z) = a(\tau)c(x) + b(x)$$

which arises when

$$Y = b(X) + c(X)W$$

and W is independent of X with τ -quantile $Q_W(\tau) = a(\tau)$. The error free τ -quantile is ($c(x) \geq 0$ is assumed)

$$Q_X(\tau, x) = c(x)Q_W(\tau) + b(x).$$

The relevant term in (4) is

$$\frac{Q_X^{\tau x}(\tau, z)Q_X^x(\tau, z)}{Q_X^\tau(\tau, z)} = \frac{c^x(z)}{c(z)} (Q_W(\tau)c^x(z) + b^x(z)).$$

This term further modifies the τ -free part of the QRF adding the term $c^x(z)b^x(z)/c(z)$ and modifies the form of the covariate dependence of shape and dispersion.

4. ACCURACY OF THE APPROXIMATION

This section examines the accuracy of the approximation to error contaminated QRFs. Most of the results are obtained using numerical methods to calculate the exact error contaminated QRF.

First consider the fully Gaussian model in which the error contaminated QRF can be obtained in closed form. Here we find that the approximation is in fact *exact* so far as capturing the dependence of the QRF on covariates is concerned. Approximation error causes the approximate QRFs to be more widely separated than the exact QRFs an effect not exactly captured in the approximation.

4.1. Analytic results for the Gaussian model. Let Y given $X = x$ and $V = v$ be $N(x'\beta, \eta^2)$ with X and V jointly normally distributed as follows.

$$\begin{bmatrix} X \\ V \end{bmatrix} \sim N \left(\begin{bmatrix} \mu_X \\ 0 \end{bmatrix}, \begin{bmatrix} \Sigma_{XX} & 0 \\ 0 & \Sigma \end{bmatrix} \right)$$

Let $Q_N(\tau)$ be the τ -quantile of a $N(0,1)$ variate. Exact error free and contaminated QRFs are

$$Q_{Y|X}(\tau, x) = x'\beta + \eta Q_N(\tau)$$

$$Q_{Y|Z}(\tau, z) = \mu'_X (\Sigma_{XX} + \Sigma)^{-1} \Sigma \beta + z' \left(I - \Sigma (\Sigma_{XX} + \Sigma)^{-1} \Sigma \right) \beta + a(\beta, \Sigma_{XX}, \Sigma) Q_N(\tau)$$

where

$$a(\beta, \Sigma_{XX}, \Sigma) = \left(\eta^2 + \beta' \left(\Sigma - \Sigma (\Sigma_{XX} + \Sigma)^{-1} \Sigma \right) \beta \right)^{1/2}.$$

Using $g_Z^j(z)$ in (2), as suggested in Section 2.3 gives the following approximation to the error contaminated QRF.

$$\begin{aligned} \tilde{Q}_{Y|Z}(\tau, z) &= \mu'_X (\Sigma_{XX} + \Sigma)^{-1} \Sigma \beta + z' \left(I - \Sigma (\Sigma_{XX} + \Sigma)^{-1} \Sigma \right) \beta + \tilde{a}(\beta, \Sigma) Q_N(\tau) \\ \tilde{a}(\beta, \Sigma) &= \eta + \frac{1}{2\eta} \beta' \Sigma \beta \end{aligned}$$

The regression coefficients of the approximate QRF are the *same* as those of the exact QRF. Approximation error arises only in the intercept, because of error in \tilde{a} as an approximation¹³ to a .

The approximate QRFs are more widely separated than the exact QRFs, that is $|\tilde{Q}_{Y|Z}(\tau, z) - h(z)| \geq |Q_{Y|Z}(\tau, z) - h(z)|$ where

$$h(z) = \mu'_X (\Sigma_{XX} + \Sigma)^{-1} \Sigma \beta + z' \left(I - \Sigma (\Sigma_{XX} + \Sigma)^{-1} \Sigma \right) \beta.$$

To see this, write

$$\begin{aligned} |\tilde{Q}_{Y|Z}(\tau, z) - h(z)|^2 &= \tilde{a}^2 Q_N(\tau)^2 \\ |Q_{Y|Z}(\tau, z) - h(z)|^2 &= a^2 Q_N(\tau)^2, \end{aligned}$$

suppressing parameter dependence of a and \tilde{a} in the notation, then

$$\tilde{a}^2 - a^2 = \frac{1}{4\eta^2} (\beta' \Sigma \beta)^2 + \beta' \Sigma (\Sigma_{XX} + \Sigma)^{-1} \Sigma \beta \geq 0$$

which delivers the required result. The approximation error is small when $\beta' \Sigma \beta$ is close to zero and large when η is small relative to $\beta' \Sigma \beta$. It tends to be small when error free covariates have a dispersed distribution.

¹³Note that a Taylor series expansion around $\Sigma = 0$ gives $\tilde{a}(\beta, \Sigma) = a(\beta, \Sigma_{XX}, \Sigma) + o(\Sigma)$ showing the the approximation is indeed correct apart from terms of order $o(\Sigma)$.

4.2. Numerical calculations for exponential power distributions. It is difficult to find other cases in which exact error contaminated QRFs can be obtained in closed form so the accuracy of the approximation is now examined using numerical methods. Attention is confined to models with a single covariate.

Let Y be determined by a location shift model in which

$$Y = \beta_0 + \beta_1 X + \sigma_W W \quad (6)$$

$$Z = X + \sigma U \quad (7)$$

and W and U (both mean 0, variance 1), and X (mean 0, variance 3) are mutually independently distributed with exponential power (EP) distributions¹⁴ with shape parameters γ_W , γ_X and γ_U .

A random variable S with mean μ and variance λ^2 and an exponential power distribution with shape parameter $\gamma \in (-1, 1)$ has the following probability density function.

$$f_S(s) = A \exp \left(-B \left| \frac{s - \mu}{\lambda} \right|^{\frac{2}{1+\gamma}} \right)$$

The constants A and B are defined in Appendix 3. Setting γ equal to 0 and 1 gives respectively normal and Laplace distributions. As $\gamma \rightarrow -1$ the density approaches the uniform density function on $(\mu - \sqrt{3}\lambda, \mu + \sqrt{3}\lambda)$.

Let $Q_\gamma(\tau)$ denote the τ -quantile¹⁵ of a zero mean unit variance EP variate with shape parameter γ . Then the error free QRF is

$$Q_{Y|X}(\tau, x) = \beta_0 + \beta_1 X + \sigma_W Q_{\gamma_W}(\tau). \quad (8)$$

To obtain the exact error contaminated QRF the conditional distribution function of Y given Z is calculated by numerical integration¹⁶ and the value of the QRF at values of z is obtained using a Newton type method¹⁷.

Figures 1 - 3 - show error free (dotted), exact error contaminated (solid) and approximate error contaminated (dashed) τ -QRFs when $\beta_0 = 0$, $\beta_1 = 1$, $\eta = 1$, $\sigma_{XX} = 3$, and $\sigma^2 = 1$. At these settings R^2 in the error free mean regression is 0.75, the signal to noise ratio for the error contaminated covariate is 0.75 and for mean regression the attenuation of the error contaminated regression is 25%, that is $E[Y|Z = z] = 0.75z$ compared with $E[Y|X = x] = x$.

The graphs show τ -QRFs for $\tau \in \{0.5, 0.75, 0.9\}$. Figures 1, 2 and 3 are distinguished by the choice of shape parameter in the EP distribution for W , with γ_W equal to 0.5, 0 and -0.5 respectively. The variance of the error contaminated covariate is 4 and the graphs show QRFs for $z \in [-4, 4]$, that is ± 2 standard deviations around the mean.

In each 3×3 array of graphs the shape parameter of the EP distribution of X varies across rows with γ_X equal to -0.5 in the top row, then 0 and 0.5. The shape parameter of the EP distribution of measurement error, V , varies across columns with γ_X equal to -0.5 in the left column, then 0 and 0.5. Thus the centre pane on each page shows QRFs when both X and V are normally distributed.

First consider the *exact* error contaminated QRFs (solid lines). Attenuation (around 25%) is evident in every case. The exact error contaminated QRFs are nonlinear except

¹⁴Box and Tiao (1973) give a discussion of the properties of EP distributions.

¹⁵An easily computed expression for the EP τ -quantiles is given in Appendix 3.

¹⁶The Splus 2000 (1999) procedure *integrate* is used. This employs an adaptive 15-point Gauss-Kronrod quadrature based on the Fortran function *dqage* and *dqgie* from QUADPACK (Piessens et al. (1983)) in NETLIB (Dongarra and Grosse (1987)).

¹⁷The Splus 2000 procedure *uniroot* is used. This implements Brent's (1973) safeguarded polynomial interpolation procedure for solving a univariate nonlinear equation, based on the Fortran function *zeroin* from NETLIB (Dongarra and Grosse 1987).

Figure 1: Exact and approximate τ -QRFs: $\tau \in \{0.5, 0.75, 0.9\}$, $\gamma_Y = +0.5$




Figure 2: Exact and approximate τ -QRFs: $\tau \in \{0.5, 0.75, 0.9\}$, $\gamma_Y = 0.0$





Figure 3: Exact and approximate QRFs: $\tau \in \{0.5, 0.75, 0.9\}$, $\gamma_Y = -0.5$



when X and V are both normally distributed although the nonlinearity is very weak when the error free covariate is normal (centre rows).

Varying the shape of the distribution of W (compare graph arrays) and V (compare columns) has *little* effect on the error contaminated QRFs. Varying the shape of the distribution of the error free covariate X (compare rows) has a *substantial* effect. When this distribution is peaked (bottom rows) attenuation is most marked at values of Z near the centre of the distribution of X . When it is platykurtic (top rows) attenuation is most marked for values of z in the tail area of the distribution of X .

The shapes of the error contaminated QRFs vary little as τ is altered. The additional noise introduced by measurement error moves the QRFs away from the median QRF.

Now consider the *approximate* error contaminated QRFs (dashed lines). These are calculated using (5) with $g_Z^j(z)$ in place of $g_X^j(z)$ because it is in this form that the approximation is used in the sensitivity analysis described in Section 2.3. In every case the approximation accurately captures the attenuation and nonlinearity in the error contaminated QRF. The location of the error contaminated QRF is very accurately captured by the approximate median regressions ($\tau = 0.5$) but the approximate QRFs tend to be located a little above (below¹⁸) the exact QRFs for $\tau > 0.5$ (< 0.5). The quality of the approximations varies only a little as the three EP shape parameters are altered.

In summary, with linear error free QRFs, in the cases studied, error contaminated QRFs are significantly nonlinear unless the error free covariate is normally distributed. The main QRF deforming impact of measurement error is driven by the shape of the distribution of the error free covariate. When the variance of measurement error is not too large, this shape is reflected in the shape of the distribution of the error contaminated covariate which is the driving force in the approximation (5). As a result the approximation captures the nonlinearity in the error contaminated QRFs well, although there is some error in locating the vertical location of the extreme QRFs. In the ‘‘bias correction’’ procedure and sensitivity analysis developed in Section 5 this location error has little impact because data on Y is used to ‘‘calibrate’’ the locations of the QRFs.

5. BIAS CORRECTION AND SENSITIVITY ANALYSIS

Small variance approximations like that developed here can be used to gauge the sensitivity of estimators to varying amounts of measurement error and to obtain estimators with reduced inconsistency. Examples are provided in Chesher and Schluter (1999) and Chesher and Santos Silva (1992). In this Section we examine the potential of small variance approximations in this regard in the context of QRF estimation.

Suppose a parametric form of a QRF is specified - here we just consider the simplest case in which error free QRFs are linear and parallel so that Y is generated by the location shift model (6). The τ -QRF of Y given X is

$$Q_X(\tau, x) = \beta_0 + \beta_1 x + \sigma_W Q_W(\tau)$$

where $Q_W(\tau)$ is the τ -quantile of W .

The results in Section 3.2 give the following expression for the approximate error contaminated τ -QRF.

$$\begin{aligned} \tilde{Q}_Z(\tau, z) &= \beta_0^*(\tau) + \beta_1 (z + \sigma^2 g_Z^x(z)) \\ \beta_0^*(\tau) &= \beta_0 + \sigma_W Q_W(\tau) - \frac{\sigma^2}{2\sigma_W} \beta_1^2 g_W^w(Q_W(\tau)) \end{aligned} \tag{9}$$

The function $g_Z^x(z)$ is the derivative of the log density of the error contaminated covariate, a function that can be estimated ($\hat{g}_X^x(z)$) with the data available. It is used here following

¹⁸Not shown.

the suggestion in Section 2.3 where it is noted that substituting this function for $g_X^x(z)$ (a) does not alter the order of the approximation error and (b) allows realisations of Z to be used to estimate the approximate error contaminated QRF.

If we knew the variance of measurement error then we could estimate the error contaminated QRF using $z + \sigma^2 \hat{g}_Z^x(z)$ as the right hand side variable and regard its estimated coefficient as an estimator of the slope of the error free QRF, β_1 . If the approximation is accurate then we expect the inconsistency of this estimator to be small. The argument in Chesher and Santos Silva (2001) suggests that the difference between the pseudo-true value of this estimate and the error free QRF coefficient, β_1 , will be of order $o(\sigma^2)$. We cannot identify other parameters of the error free QRF. This presents an alternative to the method of He and Liang (2000) who also impose a restriction on the variances in this problem and study linear QRFs.

A similar procedure could be implemented if we had an estimate of the measurement error variance, perhaps from an ancillary sample.

In the absence of knowledge of σ^2 a sensitivity analysis could be conducted, fixing σ^2 at a sequence of values in some plausible range, estimating the parameters of (9) at each chosen value of σ^2 .

Finally, if $g_Z^x(z)$ is sufficiently nonlinear we could estimate using z and an estimate of $g_Z^x(z)$ as separate explanatory variables producing approximately consistent estimators of β_1 and σ^2 .

Since the approximate error contaminated QRF is to some extent misspecified, inference should be conducted in the context of inference for misspecified M-estimators set out in White (1994).

The method proposed involves two step estimation with a nonparametric plug-in estimator used at the first stage but that plug-in estimate is determined entirely by realisations of the error contaminated covariate and, note, it *is* the error contaminated QRF that we are attempting to estimate. The principle of conditionality suggests that we should make inference conditional on covariate's realised values. If we follow that principle, the impact of variation in the plug in estimate on the sampling variance of the QRF estimator will be carried into the conditional (on the realised values of Z) standard errors through the realised values of Z and $\hat{g}_Z^x(z)$.

In order to examine the performance of a procedure of this sort the results of a series of Monte Carlo experiments are now reported. The error free QRF is linear with $\beta_0 = 0$, $\beta_1 = 1$, $\sigma_W = 1$ and the distributions of W , X and V are exponential power distributions with mean zero and shape parameters $\gamma_W, \gamma_X, \gamma_V \in \{-0.5, 0, +0.5\}$, a total of 27 cases in all. The variances of W and V were set to one and the variance of X was set to 3. At these settings the R^2 in an error free mean regression is 0.75 and the attenuation of the error contaminated mean regression is 25%, that is the OLS estimator of β_1 using error contaminated Z has probability limit equal to 0.75. In each experiment a sample size of 400 was used and there were 2000 replications.

We show results of two types. In the first we use the exact function $g_Z^x(z)$ in constructing the new right hand side variables. In the second we estimate the function using an exponential series estimator.

5.1. Log density derivative $g_Z^x(z)$ known. There are three tables of results, each showing means and standard deviations of estimates across the 2000 Monte Carlo replications¹⁹. The first, second and third sets of 9 rows show results for τ equal to respectively 0.5 (median regression), 0.75 and 0.90.

Table 1 shows results for the QRF estimator ignoring measurement error. The attenuation effect of measurement error is plain to see. In all cases the mean of the estimates of

¹⁹The sampling distributions seem close to symmetric, with means very close to medians, the latter thus not reported.

β_1 is very close to 0.75. The standard deviation of the estimates increases as τ increases as one would expect from the sampling theory of QRF estimators. There is little variation in the average value of the QRF estimator across values of the EP distribution shape parameters and across τ -QRFs.

Table 2 shows results for the QRF estimator with σ^2 “known”. The improvement is substantial. The mean of the estimates of β_1 is very close to 1 (the error free QRF value), deviating at most by 3.6%. The accuracy of estimation is slightly impaired - the standard deviations of the measurement error “corrected” estimates are around 25% higher than the standard deviations of the naive estimator which ignores measurement error. There is a small amount of variation as the EP distribution shape parameters are altered. In the case in which the measurement error distribution is platykurtic the slope estimates are slightly downward biased at $\tau = 0.5$ and slightly upward biased at $\tau = 0.9$. There is the opposite effect when the measurement error distribution is leptokurtic with slight upward bias at $\tau = 0.5$ and slight downward bias at $\tau = 0.9$. These biases are, in all cases, very small.

Table 3 shows results when σ^2 is “estimated”. When X is normally distributed there is extreme multicollinearity between z and $g_Z^x(z)$ and results are not shown for this case. Of course when measurement error is also normally distributed $g_Z^x(z) \propto z$ and σ^2 cannot be identified from the approximate QRF, or indeed at all, without further information.

Estimating σ^2 brings significant degradation in performance and now we find that one of the EP distribution shape parameters has a substantial influence, the shape parameter for the distribution of measurement error. The results vary only a little as the other shape parameters and τ are altered. With normal measurement error ($\gamma_V = 0$) the average of the “corrected” slope estimates is still very close to 1, deviating at most by 2.9%. With $\gamma_V = -0.5$, in which case the measurement error distribution is distinctly platykurtic, the “corrected estimates” are around 15% downward biased (compared with 25% for the naive estimator). With $\gamma_V = +0.5$ (leptokurtic) there is around 8% upward bias.

When σ^2 is estimated there is degradation in accuracy, standard deviations of the slope estimates increasing roughly fourfold. This is an effect that can be driven down by using larger samples. Of course in situations when $g_Z^x(z)$ is highly nonlinear this problem will be eased, but note that for real benefit to arise, this should be a nonlinearity arising from the distribution of error free X - if it arises from the distribution of V then the residual bias is likely to be large.

5.2. Log density derivative $g_Z^x(z)$ unknown. There are two sets of tables, laid out as in the previous section. Table 4 gives results with σ^2 known and Table 5 gives results with σ^2 unknown.

The estimated log density derivative $g_Z^x(z)$ is derived from the exponential series density estimator of Barron and Sheu (1993). The data are mapped by affine transformation onto the unit interval²⁰ and the unknown density of z is specified as

$$f_Z(z) \propto f_Z^0(z) \exp \left(\sum_{j=1}^m \theta_j h_j(z) \right) \quad (10)$$

where $f_Z^0(z) = 1$ is the uniform kernel density on $[0, 1]$ and the $h_j(\cdot)$ is the j th order Legendre polynomial. The required log density derivative is simply

$$g_Z^x(z) = \sum_{j=1}^m \theta_j h_j'(z) \quad (11)$$

²⁰The minimum and maximum of the realised values of Z are associated with respectively 0.1 and 0.9 to avoid end effects.

where $h'_j(\cdot)$ is the first derivative of the j th order Legendre polynomial.

The parameters θ are estimated by maximum likelihood regarding (10) as specifying the form of the likelihood contributions, the constant of integration being found by numerical methods²¹. We choose $m = 8$ to produce the results given here. In a truly nonparametric estimation one would regard m as a smoothing parameter and determine a data driven appropriate value, for example by cross validation. In these Monte Carlo experiments m was fixed at a value which allowed the essential features of the density of Z to be captured while avoiding excessive roughness in the estimate.

First consider the case in which σ^2 is known and compare Tables 2 and 4. It is clear that estimating $g_Z^x(\cdot)$ has little effect on the bias of the measurement error corrected slope estimator, but it does slightly reduce the accuracy of the estimator, standard deviations across Monte Carlo replications rising by around 20%.

When σ^2 is estimated (compare Tables 3 and 5) the standard deviations of the slope estimates rise by two to four fold compared with the case when σ^2 is known and $g_Z^x(\cdot)$ is estimated, and by around 15% compared with the case in which σ^2 is estimated and $g_Z^x(\cdot)$ is known.

There is a significant increase in bias which is downward in all the cases considered. Since $\hat{g}_Z^x(z)$ is $g_Z^x(z)$ contaminated with measurement error, this could itself be a measurement error effect. Much smaller bias is found using smaller values of the smoothing parameter²², m , but then the variance of the measurement error corrected estimator is much larger. If an attempt at estimating the measurement error variance is to be made, then, to avoid attenuation it seems to be important not to undersmooth when estimating $g_Z^x(z)$, and to have a large sample to hand.

5.3. Discussion. In the simple cases considered, estimation of approximate measurement error contaminated QRFs brings about a substantial reduction in bias but with an increase in variance that is small if the variance of measurement error is known, but sizeable otherwise. The proposed procedures are likely to work well in real problems only in large samples. But in many cases in microeconomic work in which QRF estimation would be contemplated large samples will be available²³, so perhaps this is not a great drawback.

Of more concern are the difficulties that would likely be encountered were more flexible forms of the error free QRF to be entertained. Once the error free QRF is specified as flexible and nonlinear there is the likelihood of collinearity between the derivatives of the error free QRF that appear in (2) and $g_Z^x(z)$. Another difficulty in nonlinear models is that if there are values of X at which the QRF is highly nonlinear then we can expect the approximation to have a large remainder term because it depends on the magnitude of the third derivatives of the error free QRF.

There is a further issue to consider. In practice QRFs are sometimes estimated in order to investigate heteroskedasticity. Dependence on X in the error free QRF that depends upon τ is manifested in the error contaminated QRF differently from dependence that is τ independent - see Section 3.2. To use the procedure developed here one must be specific about the interaction between X and τ in determining the error free QRF. In practice arriving at such a specification might be difficult and the resulting additional functions of z that arise may be highly collinear.

²¹Further details of the implementation of this procedure can be found in Chesher (1998b). The Monte Carlo experiments were conducted using R (Ihaka and Gentleman (1996), Hornik (2001)). In the density estimation, maximum likelihood estimation was done using the `nlm` procedure in R, a Newton type procedure described in Dennis and Schnabel (1983) and Schnabel, Koontz and Weiss (1985). QRFs were estimated using the procedure `rq` in the R contributed package `quantreg` which is an implementation of the modified Barrodale Roberts algorithm described in Koenker and d'Orey (1987,1994).

²²For example the bias is reduced by around 50% on choosing $m = 4$.

²³See the applications cited in Section 1.

We have only examined performance when there is a single covariate. Results in Chesher (1998b) for mean regression suggest that we can expect similarly good performance in multiple covariate problems as long as *only one* covariate is measured with error and the conditional density of the error contaminated covariates given the error free covariates depends on the latter through a single index.

6. CONCLUDING REMARKS

Covariate measurement error causes fundamental changes in conditional quantile regression functions, altering their shape, orientation and location. This paper has provided information about the generic effects of measurement error by developing a small measurement error variance approximation to measurement error contaminated τ -QRFs. The approximation depends upon the error free QRF and its derivatives up to order two, the variance of measurement error, and the density of the error contaminated covariates. It does not depend upon, and to use it one needs no knowledge of, the specific form of the density of measurement error.

Exact calculations suggest that the approximation can be accurate when the amount of measurement error is small to moderate, as long as the error free QRF is not too nonlinear and the measurement error distribution is not too far from normal.

A number of uses of the approximation have been proposed.

1. It allows one to gauge the likely effects of measurement error on a particular form for an error free QRF that is proposed for use in analysis of data. With realisations of the error contaminated covariate one can estimate the terms in the approximation that depend on the density of this variate and, with a particular form for the error free QRF to hand, one can derive the remaining terms.
2. With knowledge of, or an estimate of, the variance of measurement error, it can be used to produce a measurement error corrected estimate of the parameters of the error free QRF. This works well for linear QRFs when the error contaminated covariate is distinctly non-normal. But away from this class of cases there are likely to be difficulties because of high collinearity between the variables that appear in the error free QRF and the additional terms that appear in the approximation.
3. It can be used to examine the sensitivity of QRF estimates to alternative assumed amounts of measurement error by estimating the approximate error contaminated QRF for a range of values of the measurement error variance.

Obtaining consistent estimates of error free QRFs when only error contaminated covariate data are available is a challenging problem. This paper has made progress in one direction, namely (a) improving understanding of the impact of covariate measurement error on QRFs and (b) providing a tool for sensitivity analysis, but it does not offer a widely applicable solution to the consistent estimation problem. It does not seem possible to develop an instrumental variable based solution to this problem. An approach exploiting replicate measurements or validation samples offers more prospect of success, an approach under investigation for mean regression when the form of the error free regression is unknown²⁴.

²⁴See Li and Vuong (1998) and Schennach (2000).

APPENDIX 1: EXPRESSING APPROXIMATE QRFs AS FUNCTIONALS OF ERROR FREE QRFs

We use an abbreviated notation and consider conditional quantiles defined by the following equation.

$$F(Q|x) = \tau \quad (\text{A1.1})$$

Of course $Q = Q(\tau, x)$ a dependence we make explicit where otherwise there might be confusion.

Considering variations in x , τ and Q subject to (A1.1) there is

$$F^Y(Q|x)dQ + \sum_i F^i(Q|x)dx_i = d\tau \quad (\text{A1.2})$$

where

$$\begin{aligned} F^Y(Q|x) &= \left. \frac{\partial}{\partial y} F^Y(y|x) \right|_{y=Q} \\ F^i(Q|x) &= \left. \frac{\partial}{\partial x_i} F^Y(y|x) \right|_{y=Q}. \end{aligned}$$

Shortly second partial derivatives appear, F^{YY} , F^{Yi} and F^{ij} , defined similarly. Equation (A1.2) leads directly to the following expressions for the first partial derivatives of the conditional quantile function.

$$Q^\tau(\tau, x) = \frac{1}{F^Y(Q|x)} \quad (\text{A1.3})$$

$$Q^i(\tau, x) = -\frac{F^i(Q|x)}{F^Y(Q|x)} \quad (\text{A1.4})$$

The second order partial derivatives of the quantile function follow on differentiating (A1.3) and (A1.4).

$$Q^{\tau\tau}(\tau, x) = -\frac{F^{YY}(Q|x)}{F^Y(Q|x)^2} Q^\tau(\tau, x) = -\frac{F^{YY}(Q|x)}{F^Y(Q|x)^3} \quad (\text{A1.5})$$

$$\begin{aligned} Q^{\tau i}(\tau, x) &= -\frac{1}{F^Y(Q|x)^2} (F^{Yi}(Q|x) + F^{YY}(Q|x)Q^i(\tau, x)) \\ &= -\frac{F^{Yi}(Q|x)}{F^Y(Q|x)^2} + \frac{F^{YY}(Q|x)F^i(Q|x)}{F^Y(Q|x)^3} \end{aligned} \quad (\text{A1.6})$$

$$\begin{aligned} Q^{ij}(\tau, x) &= -\frac{1}{F^Y(Q|x)} (F^{Yi}(Q|x)Q^j(\tau, x) + F^{ij}(Q|x)) \\ &\quad + \frac{F^i(Q|x)}{F^Y(Q|x)^2} (F^{YY}(Q|x)Q^j(\tau, x) + F^{Yj}(Q|x)) \\ &= -\frac{F^{ij}(Q|x)}{F^Y(Q|x)} + \frac{F^{Yi}(Q|x)F^j(Q|x)}{F^Y(Q|x)^2} + \frac{F^{Yj}(Q|x)F^i(Q|x)}{F^Y(Q|x)^2} \\ &\quad - \frac{F^{YY}(Q|x)F^i(Q|x)F^j(Q|x)}{F^Y(Q|x)^3} \end{aligned} \quad (\text{A1.7})$$

In the main text we noted that

$$\left. \frac{\partial Q_Z}{\partial \sigma_{ij}} \right|_{\Sigma=0} = -\frac{F_{Y|X}^i(Q_Z|z)g_X^j(z)}{F_{Y|X}^Y(Q_Z|z)} - \frac{1}{2} \frac{F_{Y|X}^{ij}(Q_Z|z)}{F_{Y|X}^Y(Q_Z|z)} \quad (\text{A1.8})$$

which we now wish to express in terms of the conditional QRF and its derivatives. The leading term is given directly by (A1.4) with suitable expansion of notation. Now note that, from (A1.6),

$$\frac{F^{Yi}(Q|x)F^j(Q|x)}{F^Y(Q|x)^2} = \frac{Q^{\tau i}(\tau, x)Q^j(\tau, x)}{Q^\tau(\tau, x)} - \frac{Q^{\tau\tau}(\tau, x)Q^i(\tau, x)Q^j(\tau, x)}{Q^\tau(\tau, x)^2}.$$

and from (A1.7), exploiting (A1.3) and (A1.4)

$$\frac{F^{ij}(Q|x)}{F^Y(Q|x)} = -Q^{ij}(\tau, x) + \frac{Q^{\tau i}(\tau, x)Q^j(\tau, x)}{Q^\tau(\tau, x)} + \frac{Q^{\tau j}(\tau, x)Q^i(\tau, x)}{Q^\tau(\tau, x)} - \frac{Q^{\tau\tau}(\tau, x)Q^i(\tau, x)Q^j(\tau, x)}{Q^\tau(\tau, x)^2}$$

Substituting this final expression in (A1.8) gives equation (2) in the main text.

APPENDIX 2: THE EFFECT ON THE APPROXIMATION OF USING THE LOG DENSITY OF Z RATHER THAN X .

Chesher (1991) shows that the densities of Z and X satisfy

$$f_Z(z) = f_X(z) + \sum_{s,t} \sigma_{st} f_X^{st}(z) + o(\Sigma).$$

The log densities therefore satisfy

$$g_Z(z) = g_X(z) + \sum_{s,t} \sigma_{st} \frac{f_X^{st}(z)}{f_X(z)} + o(\Sigma)$$

and their derivatives satisfy

$$g_Z^j(z) = g_X^j(z) + \sum_{s,t} \sigma_{st} \left(\frac{f_X^{stj}(z)}{f_X(z)} - \frac{f_X^{st}(z)f_X^j(z)}{f_X(z)^2} \right) + o(\Sigma).$$

It follows immediately that

$$\sum_{i,j} \sigma_{ij} Q_X^i(\tau, z) g_X^j(z) - \sum_{i,j} \sigma_{ij} Q_X^i(\tau, z) g_Z^j(z) = o(\Sigma)$$

and then directly that the order of the approximation error in (2) is not increased on substituting $g_Z^j(z)$ for $g_X^j(z)$.

APPENDIX 3: EXPONENTIAL POWER DISTRIBUTIONS: QUANTILES AND RANDOM NUMBER GENERATION

Let S have an exponential power distribution with mean μ and variance λ^2 and shape parameter $\gamma \in (-1, 1)$. The probability density function of S is

$$f_S(s) = A \exp\left(-B \left|\frac{s - \mu}{\lambda}\right|^{\frac{2}{1+\gamma}}\right)$$

where A and B are defined as follows.

$$A = \frac{1}{\lambda} \left(\frac{\Gamma(\frac{3}{2}(1+\gamma))}{(1+\gamma)\Gamma(\frac{1}{2}(1+\gamma))^{3/2}} \right)^{1/2} \quad B = \left(\frac{\Gamma(\frac{3}{2}(1+\gamma))}{\Gamma(\frac{1}{2}(1+\gamma))} \right)^{\frac{1}{1+\gamma}}$$

Let G have a Gamma distribution with mean and variance δ . The density function of $G \in [0, \infty]$ is

$$f_G(g) = \Gamma(\delta)^{-1} g^{\delta-1} \exp(-g).$$

Quantiles. Many statistical packages have fast routines for calculating Gamma quantiles. These can be used to calculate EP quantiles, as follows.

Let $Q_G(\tau; \delta)$ be the τ -quantile of G . Let $Q_S(\tau; \mu, \lambda, \gamma)$ be the τ -quantile of S . Quantiles of S are related to quantiles of G as follows.

$$Q_S(\tau; \mu, \lambda, \gamma) = \mu + \lambda \operatorname{sign}(\tau - 0.5) \left(B^{-1} Q_G\left(1 - \frac{2 \min(\tau, 1 - \tau) \lambda^{1/2}}{(1 + \gamma)^{1/2} \Gamma(\frac{1}{2}(1 + \gamma))^{\frac{3}{4}}}, \frac{1}{2}(1 + \gamma)\right) \right)^{\frac{1+\gamma}{2}}$$

Pseudo-random number generation. The EP quantile formula leads directly to fast pseudo-random number generation because, if K has a uniform distribution on $[0, 1]$, then $Q_S(K; \mu, \lambda, \gamma)$ has an EP distribution with mean μ , variance λ^2 and shape parameter γ .

ACKNOWLEDGMENTS

I am grateful to the ESRC for support of part of this research under grant number R0027836, and to the Leverhulme Trust for support at the Centre for Microdata Methods and Practice.

REFERENCES

- ABADIE, A., ANGRIST, J., AND G. IMBENS (2001): "Instrumental variables estimates of the effect of subsidized training on the quantiles of trainee earnings", *Econometrica*, forthcoming.
- ABRAVAYA, J., (2001): "The effects of demographics and maternal behavior on the distribution of birth outcomes", *Empirical Economics*, 19, 247-257.
- ARIAS, O., HALLOCK, K.F., AND W. SOSA-ESCUADERO (2001): "Individual heterogeneity in the returns to schooling: instrumental variables quantile regression using twins data", *Empirical Economics*, 26, 7-40.

BARRON, A.R., AND C-H. SHEU (1991): "Approximation of density functions by sequences of exponential families", *Annals of Statistics*, 19, 1347-1369.

BASSETT, G.W. JR., AND H-L. CHEN (2001): "Portfolio style: Return-based attribution using quantile regression", *Empirical Economics*, 19, 293-305.

BASSETT, G. JR., AND R KOENKER (1982): "An empirical quantile function for linear models with iid errors," *Journal of the American Statistical Association*, 77, 407-415.

BOX, G.E.P., AND TIAO, G.C. (1973) *Bayesian Inference in Statistical Analysis*, New York: Wiley Classics.

BRENT, R. (1973): *Algorithms for Minimization without Derivatives*. Prentice-Hall, Englewood Cliffs, NJ, USA.

BROWN, M.L. (1982): "Robust line estimation with errors in both variables," *Journal of the American Statistical Association*, 77, 71-79. Correction in 78, 1008.

BUSHINSKY, M., (2001): "Quantile regression with sample selection: Estimating women's return to education in the U.S.," *Empirical Economics*, 19, 87-113.

BUSHINSKY, M., (1998a): "Recent advances in quantile regression models - A practical guideline for empirical research", *Journal of Human Resources*, 33, 88-126.

BUCHINSKY, M., (1998b): "The dynamics of changes in the female wage distribution in the USA: A quantile regression approach", *Journal of Applied Econometrics*, 13, 1-30.

BUCHINSKY, M., (1995): "Quantile regression, Box-Cox transformation model, and the U.S. wage structure, 1963-1987", *Journal of Econometrics*, 65, 109-154.

CARROL, R.J., RUPPERT, D., AND L.A. STEFANSKI (1995): *Measurement error in nonlinear models*, London: Chapman and Hall.

CHERNOZHUKOV, V., AND L. UMANTSEV (2001): "Conditional value-at-risk: Aspects of modeling and estimation", *Empirical Economics*, 19, 271-292.

CHESHER, A.D., (1991): "The effect of measurement error," *Biometrika*, 78, 451 - 462.

CHESHER, A.D. (1998a): "Polynomial regression with covariate measurement error", University of Bristol Economics Department Discussion Paper No. 98/448.

CHESHER, A.D., (1998b): "Measurement error bias reduction", University of Bristol Economics Department Discussion Paper No. 98/449.

CHESHER, A.D., DUMANGANE, M.B.G., AND R.J. SMITH (2001): "Duration response measurement error", forthcoming *Journal of Econometrics*.

CHESHER, A.D., AND J.M.C. SANTOS SILVA, (2001): "Taste variation in discrete choice models", forthcoming, *Review of Economic Studies*.

CHESHER, A.D., AND C. SCHLUTER (1999) "Welfare measurement and measurement error," University of Bristol Department of Economics Discussion Paper No. 99/463.

DENNIS, J. E. AND SCHNABEL, R. B., (1983): *Numerical Methods for Unconstrained Optimization and Nonlinear Equations*. Englewood Cliffs, NJ: Prentice-Hall.

DONGARRA, J. J., AND E. GROSSE (1987): "Distribution of mathematical software via electronic mail", *Communications of the ACM*, 30, 403-407.

EIDE, E.R., AND M.H. SHOWALTER (1999): "Factors affecting the transmission of earnings across generations: A quantile regression approach", *Journal of Human Resources*, 34, 253-267.

FITZENBERGER, B., HUJER, R., MCCURDY, T.E., AND R. SCHNABEL (2001): "Testing for uniform wage trends in West-Germany: A cohort analysis using quantile regressions for censored data", *Empirical Economics*, 26, 41-86.

FULLER, W.A., (1987): *Measurement Error Models*, New York: Wiley.

GINTHER, D.K., (2000) "A nonparametric analysis of the US male earnings distribution", in *Advance in Econometrics, Volume 14*, Cambridge: Cambridge University Press

HARTOG, J., PEREIRA, P.T., AND J.A.C. VIEIRA (2001): "Changing returns to education in Portugal during the 1980s and early 1990s: OLS and quantile regression estimators", *Applied Economics*, 33, 1021-1037.

- HE, X., AND H LIANG (2000): "Quantile regression estimates for a class of linear and partially linear errors-in-variables models," *Statistica Sinica*, 10, 129-140.
- HORNİK, K., (2001): "The R FAQ", <http://www.ci.tuwien.ac.at/~hornik/R/>.
- IHAKA, R., AND R. GENTLEMAN. (1996): "R: A language for data analysis and graphics", *Journal of Computational and Graphical Statistics*, 5, 299-314.
- KOENKER, R., AND G. BASSETT JR. (1978): "Regression Quantiles," *Econometrica*, 46, 33-50.
- KOENKER, R., AND Y. BILIAS (2001): "Quantile regression for duration data: A reappraisal of the Pennsylvania Reemployment Bonus Experiments", *Empirical Economics*, 19, 199-220.
- KOENKER, R.W. AND V. D'OREY (1987): "Computing regression quantiles", *Journal of the Royal Statistical Society, Series C*, 36, 383-393.
- KOENKER, R.W. AND V. D'OREY (1994): "Remark on Algorithm AS229; Computing dual regression quantiles and regression rank scores", *Applied Statistics*, 43, 410-414.
- LEVIN, J. (2001): "For whom the reductions count: A quantile regression analysis of class size and peer effects on scholastic achievement", *Empirical Economics*, 19, 221-246.
- LI, T., AND Q. VUONG (1998): "Nonparametric estimation of the measurement error model using multiple indicators," *Journal of Multivariate Analysis*, 65, 139-165.
- MACHADO, J.A.F., AND J. MATA (2000): "Box-Cox quantile regression and the distribution of firm sizes", *Journal of Applied Econometrics*, 15, 253-274.
- NAHM, J-W., (2001): "Nonparametric quantile regression analysis of R&D-sales relationship for Korean firms", *Empirical Economics*, 19, 259-270.
- PIESSENS, R., DEDONCKER-KAPENGA, E., UBERHUBER, C., AND D. KAHANER (1983): *QUADPACK: A Subroutine Package for Automatic Integration*. Springer, Berlin.
- REIERSOL, O., (1950) "Identifiability of a linear relation between variables which are subject to error," *Econometrica*, 18, 357-389.
- REILLY, B., (1999): "The gender pay gap in Russia during the transition, 1992-96", *Economics of Transition*, 7, 245-264.
- SCHENNACH, S., (2000): "Estimation of nonlinear models with measurement error, mimeo.
- SCHNABEL, R. B., KOONTZ, J. E. AND WEISS, B. E., (1985): "A modular system of algorithms for unconstrained minimization", *ACM Trans. Math. Software*, 11, 419-440.
- S-PLUS 2000 USER'S GUIDE. (1999): Data Analysis Products Division, Mathsoft, Seattle, WA.
- TAYLOR, J.W., (2000): "A quantile regression neural network approach to estimating the conditional density of multiperiod returns", *Journal of Forecasting*, 19, 299-311.
- VARIYAM, J., BLAYLOCK, J., AND D. SMALLWOOD (1999): "Characterizing the distribution of macronutrient intake among US adults: A quantile regression approach", *American Journal of Agricultural Economics*, 81, 1293-1293.
- WHITE, H. (1994): *Estimation, Inference, and Specification Analysis*. New York: Cambridge University Press.

Table 1: Means and standard deviations of QRF slope estimates ignoring measurement error

τ	γ_Y	γ_X	$\gamma_V = -0.5$		$\gamma_V = 0.0$		$\gamma_V = +0.5$	
			mean	s.d.	mean	s.d.	mean	s.d.
0.50	-0.5	-0.5	.738	.029	.755	.031	.772	.033
		0.0	.734	.031	.750	.033	.769	.034
		+0.5	.728	.034	.744	.035	.761	.038
	0.0	-0.5	.736	.030	.755	.031	.774	.032
		0.0	.732	.031	0.750	.033	.771	.034
		+0.5	.725	.034	.743	.035	.763	.035
	+0.5	-0.5	.736	.028	.756	.029	.778	.032
		0.0	.730	.030	.750	.032	.772	.033
		+0.5	.723	.033	.743	.034	.764	.037
0.75	-0.5	-0.5	.746	.034	.753	.034	.764	.036
		0.0	.742	.034	.750	.036	.761	.037
		+0.5	.739	.038	.747	.038	.757	.040
	0.0	-0.5	.746	.033	.752	.034	.763	.036
		0.0	.743	.034	.750	.036	.761	.037
		+0.5	.740	.036	.745	.038	.756	.039
	+0.5	-0.5	.747	.032	.753	.034	.763	.036
		0.0	.743	.034	.750	.035	.760	.037
		+0.5	.739	.036	.746	.038	.756	.039
0.90	-0.5	-0.5	.765	.042	.748	.044	.736	.044
		0.0	.766	.043	.750	.044	.740	.046
		+0.5	.769	.045	.754	.047	.743	.048
	0.0	-0.5	.766	.043	.747	.043	.735	.047
		0.0	.768	.044	.750	.044	.738	.047
		+0.5	.770	.045	.752	.045	.744	.048
	+0.5	-0.5	.770	.045	.746	.044	.733	.046
		0.0	.771	.044	.750	.045	.737	.047
		+0.5	.773	.046	.754	.046	.742	.048

Table 2: Means and standard deviations of measurement error corrected QRF slope estimates with σ^2 known and $g_Z^x(\cdot)$ known

τ	γ_Y	γ_X	$\gamma_V = -0.5$		$\gamma_V = 0.0$		$\gamma_V = +0.5$	
			mean	s.d.	mean	s.d.	mean	s.d.
0.50	-0.5	-0.5	0.989	.040	1.011	.040	1.028	.042
		0.0	0.978	.042	1.000	.044	1.026	.046
		+0.5	0.972	.043	0.996	.046	1.021	.050
	0.0	-0.5	0.986	.041	1.010	.040	1.031	.041
		0.0	0.976	.041	1.000	.043	1.028	.045
		+0.5	0.970	.044	0.995	.046	1.024	.047
	+0.5	-0.5	0.987	.039	1.013	.038	1.036	.041
		0.0	0.974	.040	1.000	.043	1.030	.044
		+0.5	0.966	.042	0.995	.044	1.025	.048
0.75	-0.5	-0.5	0.994	.045	1.007	.044	1.018	.046
		0.0	0.989	.046	1.000	.047	1.015	.050
		+0.5	0.988	.049	0.998	.050	1.011	.053
	0.0	-0.5	0.992	.044	1.005	.044	1.018	.046
		0.0	0.990	.046	1.000	.048	1.014	.049
		+0.5	0.988	.047	0.996	.049	1.013	.052
	+0.5	-0.5	0.993	.044	1.005	.044	1.018	.046
		0.0	0.991	.046	1.000	.047	1.014	.049
		+0.5	0.989	.047	0.997	.049	1.012	.052
0.90	-0.5	-0.5	1.004	.056	0.994	.058	0.984	.058
		0.0	1.020	.058	1.000	.059	0.986	.062
		+0.5	1.029	.058	1.005	.062	0.984	.064
	0.0	-0.5	1.005	.056	0.990	.057	0.982	.059
		0.0	1.023	.059	1.000	.059	0.984	.062
		+0.5	1.032	.059	1.003	.059	0.986	.063
	+0.5	-0.5	1.007	.059	0.988	.059	0.978	.059
		0.0	1.026	.059	1.001	.059	0.981	.062
		+0.5	1.036	.059	1.004	.059	0.984	.065

Table 3: Means and standard deviations of measurement error corrected QRF slope estimates with σ^2 unknown and $g_Z^x(\cdot)$ known

τ	γ_Y	γ_X	$\gamma_V = -0.5$		$\gamma_V = 0.0$		$\gamma_V = +0.5$	
			mean	s.d.	mean	s.d.	mean	s.d.
0.50	-0.5	-0.5	0.870	0.107	1.024	.127	1.087	.130
		0.0	-	-	-	-	-	-
		+0.5	1.117	.168	1.017	.161	0.910	.149
	0.0	-0.5	0.867	.106	1.023	.122	1.095	.129
		0.0	-	-	-	-	-	-
		+0.5	1.123	.160	1.018	.161	0.909	.152
	+0.5	-0.5	0.874	.105	1.029	.120	1.101	.128
		0.0	-	-	-	-	-	-
		+0.5	1.122	.164	1.020	.158	0.908	.152
0.75	-0.5	-0.5	0.892	.121	1.013	.137	1.074	.142
		0.0	-	-	-	-	-	-
		+0.5	1.106	.180	1.008	.180	0.899	.161
	0.0	-0.5	0.888	.119	1.017	.133	1.078	.146
		0.0	-	-	-	-	-	-
		+0.5	1.098	.170	1.004	.175	0.903	.161
	+0.5	-0.5	0.890	.116	1.013	.136	1.073	.144
		0.0	-	-	-	-	-	-
		+0.5	1.102	.178	1.011	.170	0.903	.162
0.90	-0.5	-0.5	0.933	.152	0.988	.181	1.015	.188
		0.0	-	-	-	-	-	-
		+0.5	1.077	.218	0.988	.216	0.880	.194
	0.0	-0.5	0.931	.158	0.993	.169	1.020	.192
		0.0	-	-	-	-	-	-
		+0.5	1.066	.227	0.980	.221	0.886	.194
	+0.5	-0.5	0.934	.158	0.981	.182	1.013	.196
		0.0	-	-	-	-	-	-
		+0.5	1.064	.227	0.987	.217	0.887	.201

Table 4: Means and standard deviations of measurement error corrected QRF slope estimates with σ^2 known and $g_Z^x(\cdot)$ estimated

τ	γ_Y	γ_X	$\gamma_V = -0.5$		$\gamma_V = 0.0$		$\gamma_V = +0.5$	
			mean	s.d.	mean	s.d.	mean	s.d.
0.50	-0.5	-0.5	0.979	.048	1.002	.049	1.024	.052
		0.0	0.972	.047	0.994	.050	1.021	.052
		+0.5	0.968	.047	0.991	.051	1.017	.056
	0.0	-0.5	0.977	.049	1.003	.049	1.027	.051
		0.0	0.969	.046	0.994	.049	1.024	.052
		+0.5	0.965	.048	0.991	.051	1.020	.052
	+0.5	-0.5	0.978	.048	1.005	.047	1.032	.051
		0.0	0.968	.047	0.993	.049	1.024	.052
		+0.5	0.963	.046	0.992	.049	1.021	.055
0.75	-0.5	-0.5	0.986	.053	0.999	.053	1.015	.055
		0.0	0.984	.051	0.993	.053	1.012	.056
		+0.5	0.986	.052	0.994	.054	1.008	.060
	0.0	-0.5	0.984	.052	0.999	.052	1.016	.055
		0.0	0.985	.051	0.993	.053	1.011	.057
		+0.5	0.986	.051	0.994	.053	1.009	.057
	+0.5	-0.5	0.986	.052	0.997	.052	1.015	.054
		0.0	0.986	.051	0.994	.054	1.010	.057
		+0.5	0.985	.051	0.994	.053	1.008	.057
0.90	-0.5	-0.5	0.999	.063	0.987	.064	0.979	.067
		0.0	1.015	.064	0.994	.064	0.983	.067
		+0.5	1.027	.063	1.003	.065	0.983	.068
	0.0	-0.5	0.999	.064	0.985	.064	0.977	.068
		0.0	1.019	.063	0.992	.064	0.980	.068
		+0.5	1.029	.061	1.002	.064	0.984	.067
	+0.5	-0.5	1.003	.064	0.983	.063	0.975	.067
		0.0	1.021	.064	0.997	.066	0.977	.069
		+0.5	1.032	.063	1.002	.063	0.982	.069

Table 5: Means and standard deviations of measurement error corrected QRF slope estimates with σ^2 unknown and $g_Z^x(\cdot)$ estimated

τ	γ_Y	γ_X	$\gamma_V = -0.5$		$\gamma_V = 0.0$		$\gamma_V = +0.5$	
			mean	s.d.	mean	s.d.	mean	s.d.
0.50	-0.5	-0.5	0.820	.102	0.903	.136	0.972	.169
		0.0	-	-	-	-	-	-
		+0.5	0.944	.182	0.907	.170	0.863	.148
	0.0	-0.5	0.818	.101	0.906	.137	0.974	.173
		0.0	-	-	-	-	-	-
		+0.5	0.947	.181	0.904	.153	0.865	.156
	+0.5	-0.5	0.817	.097	0.908	.128	0.976	.188
		0.0	-	-	-	-	-	-
		+0.5	0.950	.172	0.906	.150	0.862	.147
0.75	-0.5	-0.5	0.835	.118	0.900	.152	0.958	.183
		0.0	-	-	-	-	-	-
		+0.5	0.940	.187	0.902	.180	0.845	.162
	0.0	-0.5	0.830	.116	0.903	.151	0.955	.187
		0.0	-	-	-	-	-	-
		+0.5	0.939	.187	0.888	.175	0.853	.180
	+0.5	-0.5	0.830	.117	0.896	.136	0.949	.196
		0.0	-	-	-	-	-	-
		+0.5	0.941	.178	0.896	.165	0.845	.168
0.90	-0.5	-0.5	0.856	.163	0.884	.173	0.906	.220
		0.0	-	-	-	-	-	-
		+0.5	0.939	.214	0.888	.212	0.824	.199
	0.0	-0.5	0.859	.158	0.883	.193	0.902	.222
		0.0	-	-	-	-	-	-
		+0.5	0.933	.218	0.878	.219	0.829	.214
	+0.5	-0.5	0.857	.155	0.878	.173	0.898	.235
		0.0	-	-	-	-	-	-
		+0.5	0.933	.214	0.883	.203	0.823	.206