

Econ 204 2010

Lecture 4

Outline

1. Open and Closed Sets
2. Continuity in Metric Spaces

Announcements:


- PS1 due tomorrow 7/30
in lecture
- revised notes for lecture 5 (tomorrow) posted

Open and Closed Sets

Definition 1. Let (X, d) be a metric space. A set $A \subseteq X$ is open if

$$\forall x \in A \exists \varepsilon > 0 \text{ s.t. } B_\varepsilon(x) \subseteq A$$

A set $C \subseteq X$ is closed if $X \setminus C$ is open.



$$\{y \in X : d(x, y) < \epsilon\}$$

$B_\epsilon(x)$

A open

B not open
 $\nexists \epsilon > 0$ s.t.
 $B_\epsilon(x) \subseteq B$

Open and Closed Sets


Example: (a, b) is open in the metric space \mathbf{E}^1 (\mathbf{R} with the usual Euclidean metric). Given $x \in (a, b)$, $a < x < b$. Let

$$\varepsilon = \min\{x - a, b - x\} > 0$$

Then

$$\begin{aligned} -\varepsilon &\geq -(x-a) \\ \varepsilon &\leq b-x \end{aligned}$$

$$\begin{aligned} y \in B_\varepsilon(x) &\Rightarrow y \in (x - \varepsilon, x + \varepsilon) \\ &\subseteq (x - (x - a), x + (b - x)) \\ &= (a, b) \end{aligned}$$



so $B_\varepsilon(x) \subseteq (a, b)$, so (a, b) is open.

Notice that ε depends on x ; in particular, ε gets smaller as x nears the boundary of the set.

Open and Closed Sets

Example: In \mathbf{E}^1 , $[a, b]$ is closed. $\mathbf{R} \setminus [a, b] = (-\infty, a) \cup (b, \infty)$ is a union of two open sets, which must be open.

Example: In the metric space $[0, 1]$, $[0, 1]$ is open. With $[0, 1]$ as the underlying metric space,

$$B_\varepsilon(0) = \{x \in [0, 1] : |x - 0| < \varepsilon\} = [0, \varepsilon) \subset [0, 1]$$

$x \in X$
 $\text{if } \varepsilon < 1$

Thus, openness and closedness depend on the underlying metric space as well as on the set.

Open and Closed Sets

Example: Most sets are neither open nor closed. For example, in E^1 , $[0, 1] \cup (2, 3)$ is neither open nor closed.

Example: An open set may consist of a single point. For example, if $X = \mathbf{N}$ and $d(m, n) = |m - n|$, then

$$B_{1/2}(1) = \{m \in \mathbf{N} : |m - 1| < 1/2\} = \{1\}$$

Claim: $\{1\}$ is open.
Proof: take any $x \in \{1\}$. Well, then
 $x = 1$. Then $B_{1/2}(1) = \{1\} \subseteq \{1\}$
 $\Rightarrow \{1\}$ is open.

Open and Closed Sets

Example: In any metric space (X, d) both \emptyset and X are open, and both \emptyset and X are closed.

To see that \emptyset is open, note that the statement

$$\forall x \in \emptyset \exists \varepsilon > 0 B_\varepsilon(x) \subseteq \emptyset$$

is vacuously true since there aren't any $x \in \emptyset$. To see that X is open, note that since $B_\varepsilon(x)$ is by definition $\{z \in X : d(z, x) < \varepsilon\}$, $\subseteq X$ it is trivially contained in X .

Since \emptyset is open, X is closed; since X is open, \emptyset is closed.

$$= X \setminus \emptyset$$

$$\subseteq X \setminus X$$

Open and Closed Sets


Example: Open balls are open sets. $\forall x \in X, \varepsilon > 0$

Suppose $y \in B_\varepsilon(x)$. Then $d(x, y) < \varepsilon$. Let $\delta = \varepsilon - d(x, y) > 0$. If $d(z, y) < \delta$, then

$$\begin{aligned} d(z, x) &\leq d(z, y) + d(y, x) \\ &< \delta + d(x, y) \\ &= \underbrace{\varepsilon - d(x, y)} + d(x, y) \\ &= \varepsilon \end{aligned}$$

$$\Rightarrow z \in B_\varepsilon(x).$$

so $B_\delta(y) \subseteq B_\varepsilon(x)$, so $B_\varepsilon(x)$ is open.



Open and Closed Sets

Theorem 1 (Thm. 4.2). *Let (X, d) be a metric space. Then*

- 1. \emptyset and X are both open, and both closed.*
- 2. The union of an arbitrary (finite, countable, or uncountable) collection of open sets is open.*
- 3. The intersection of a finite collection of open sets is open.*

Proof. 1. We have already shown this.

2. Suppose $\{A_\lambda\}_{\lambda \in \Lambda}$ is a collection of open sets.

$$x \in \bigcup_{\lambda \in \Lambda} A_\lambda \Rightarrow \exists \lambda_0 \in \Lambda \text{ s.t. } x \in A_{\lambda_0}$$

$$\Rightarrow \exists \varepsilon > 0 \text{ s.t. } B_\varepsilon(x) \subseteq A_{\lambda_0} \subseteq \bigcup_{\lambda \in \Lambda} A_\lambda$$

so $\bigcup_{\lambda \in \Lambda} A_\lambda$ is open.

3. Suppose $A_1, \dots, A_n \subseteq X$ are open sets. If $x \in \bigcap_{i=1}^n A_i$, then

$$x \in A_1, x \in A_2, \dots, x \in A_n$$

so

\uparrow \uparrow \dots \uparrow
 open open ... open

$$\exists \varepsilon_1 > 0, \dots, \varepsilon_n > 0 \text{ s.t. } \underline{B_{\varepsilon_1}(x) \subseteq A_1}, \dots, \underline{B_{\varepsilon_n}(x) \subseteq A_n}$$



Let*

$$\varepsilon = \min\{\varepsilon_1, \dots, \varepsilon_n\} > 0$$

Then

$$\underbrace{B_\varepsilon(x) \subseteq B_{\varepsilon_1}(x) \subseteq A_1, \dots, B_\varepsilon(x) \subseteq B_{\varepsilon_n}(x) \subseteq A_n}$$

so

$$B_\varepsilon(x) \subseteq \bigcap_{i=1}^n A_i$$

which proves that $\bigcap_{i=1}^n A_i$ is open.



*Note this is where we need the fact that we are taking a finite intersection. The infimum of an infinite set of positive numbers could be zero. And the intersection of an infinite collection of open sets need not be open.

Interior, Closure, Exterior and Boundary

Definition 2. • The interior of A , denoted $\text{int } A$, is the largest open set contained in A (the union of all open sets contained in A).

$$A \text{ not open} \Leftrightarrow \text{int } A \subsetneq A$$

• The closure of A , denoted \bar{A} , is the smallest closed set containing A (the intersection of all closed sets containing A)

$$A \text{ not closed} \Leftrightarrow A \subsetneq \bar{A}$$

• The exterior of A , denoted $\text{ext } A$, is the largest open set contained in $X \setminus A$. ($= \text{int } (X \setminus A)$)

• The boundary of A , denoted $\partial A = \overline{(X \setminus A)} \cap \bar{A}$

Interior, Closure, Exterior and Boundary


Example: Let $A = [0, 1] \cup (2, 3)$. Then

$$\text{int } A = (0, 1) \cup (2, 3)$$

$$\bar{A} = [0, 1] \cup [2, 3]$$

$$\begin{aligned} \text{ext } A &= \text{int}(X \setminus A) = \text{int}((-\infty, 0) \cup (1, 2) \cup (3, +\infty)) \\ &= (-\infty, 0) \cup (1, 2) \cup (3, +\infty) \end{aligned}$$

$$\begin{aligned} \partial A &= \overline{(X \setminus A)} \cap \bar{A} \\ &= ((-\infty, 0] \cup [1, 2] \cup [3, +\infty)) \cap ([0, 1] \cup [2, 3]) \\ &= \{0, 1, 2, 3\} \end{aligned}$$



Sequences and Closed Sets

Theorem 2 (Thm. 4.13). *A set A in a metric space (X, d) is closed if and only if*


$$\{x_n\} \subset A, x_n \rightarrow x \in X \Rightarrow x \in A$$

Proof. Suppose A is closed. Then $X \setminus A$ is open. Consider a convergent sequence $x_n \rightarrow x \in X$, with $x_n \in A$ for all n . If $x \notin A$, $x \in X \setminus A$, so there is some $\varepsilon > 0$ such that $B_\varepsilon(x) \subseteq X \setminus A$ (why?). Since $x_n \rightarrow x$, there exists $N(\varepsilon)$ such that

$$\begin{aligned} n > N(\varepsilon) &\Rightarrow x_n \in B_\varepsilon(x) \subseteq X \setminus A \\ &\Rightarrow x_n \in X \setminus A \\ &\Rightarrow x_n \notin A \end{aligned}$$

contradiction. Therefore,

$$\{x_n\} \subset A, x_n \rightarrow x \in X \Rightarrow x \in A$$



Conversely, suppose


$$\{x_n\} \subset A, x_n \rightarrow x \in X \Rightarrow x \in A$$

We need to show that A is closed, i.e. $X \setminus A$ is open. Suppose not, so $X \setminus A$ is not open. Then there exists $x \in X \setminus A$ such that for every $\varepsilon > 0$,

$$B_\varepsilon(x) \not\subset X \setminus A$$

so there exists $y \in B_\varepsilon(x)$ such that $y \notin X \setminus A$. Then $y \in A$, hence

$$B_\varepsilon(x) \cap A \neq \emptyset$$



Construct a sequence $\{x_n\}$ as follows: for each n , choose

$$x_n \in B_{\frac{1}{n}}(x) \cap A$$

Given $\varepsilon > 0$, we can find $N(\varepsilon)$ such that $N(\varepsilon) > \frac{1}{\varepsilon}$ by the Archimedean Property, so $n > N(\varepsilon) \Rightarrow \frac{1}{n} < \frac{1}{N(\varepsilon)} < \varepsilon$, so $x_n \rightarrow x$. Then $\{x_n\} \subseteq A$, $x_n \rightarrow x$, so $x \in A$, contradiction. Therefore, $X \setminus A$ is open, so A is closed. \square

Continuity in Metric Spaces

Definition 3. Let (X, d) and (Y, ρ) be metric spaces. A function $f : X \rightarrow Y$ is continuous at a point $x_0 \in X$ if

$$\forall \varepsilon > 0 \exists \delta(x_0, \varepsilon) > 0 \text{ s.t. } d(x, x_0) < \delta(x_0, \varepsilon) \Rightarrow \rho(f(x), f(x_0)) < \varepsilon$$

$$\forall \varepsilon > 0 \quad \exists \delta > 0$$

f is continuous if it is continuous at every element of its domain.

Note that δ can depend on x_0 and ε .

Continuity in Metric Spaces

Continuity at x_0 requires:

- $f(x_0)$ is defined; and
- either
 - x_0 is an isolated point of X , i.e. $\exists \varepsilon > 0$ s.t. $B_\varepsilon(x_0) = \{x_0\}$;
or
 - $\lim_{x \rightarrow x_0} f(x)$ exists and equals $f(x_0)$

Continuity in Metric Spaces

Suppose $f : X \rightarrow Y$ and $A \subseteq Y$. Define

$$f^{-1}(A) = \{x \in X : f(x) \in A\}$$

Theorem 3 (Theorem 6.14). *Let (X, d) and (Y, ρ) be metric spaces, and $f : X \rightarrow Y$. Then f is continuous if and only if*


$$\left\{ f^{-1}(A) \text{ is open in } X \ \forall A \subseteq Y \text{ s.t. } A \text{ is open in } Y \right\}$$

Alternatively, f is continuous $\iff f^{-1}(C)$ is closed in X for every closed $C \subseteq Y$.

Proof. Suppose f is continuous. Given $A \subseteq Y$, A open, we must show that $f^{-1}(A)$ is open in X . Suppose $x_0 \in f^{-1}(A)$. Let $y_0 = f(x_0) \in A$. Since A is open, we can find $\varepsilon > 0$ such that $B_\varepsilon(y_0) \subseteq A$. Since f is continuous, there exists $\delta > 0$ such that

$$\begin{aligned}
 \underline{d(x, x_0) < \delta} &\Rightarrow \rho(f(x), f(x_0)) < \varepsilon \\
 &\Rightarrow f(x) \in B_\varepsilon(y_0) = \mathcal{B}_\varepsilon(f(x_0)) \\
 &\Rightarrow f(x) \in A \\
 &\Rightarrow \underline{x \in f^{-1}(A)}
 \end{aligned}$$

so $\underline{B_\delta(x_0) \subseteq f^{-1}(A)}$, so $f^{-1}(A)$ is open.



f continuous \Rightarrow

$\exists \delta > 0$

$B_\delta(x_0)$

s.t.

$x \in B_\delta(x_0)$

$\Rightarrow f(x) \in B_\epsilon(f(x_0))$


$\exists \epsilon > 0$
s.t.

Conversely, suppose

$f^{-1}(A)$ is open in $X \forall A \subseteq Y$ s.t. A is open in Y

We need to show that f is continuous. Let $x_0 \in X$, $\varepsilon > 0$. Let $A = B_\varepsilon(f(x_0))$. A is an open ball, hence an open set, so $f^{-1}(A)$ is open in X . $x_0 \in f^{-1}(A)$, so there exists $\delta > 0$ such that $B_\delta(x_0) \subseteq f^{-1}(A)$.

$$\begin{aligned}
 \underline{d(x, x_0) < \delta} &\Rightarrow x \in B_\delta(x_0) \\
 &\Rightarrow x \in f^{-1}(A) \\
 &\Rightarrow f(x) \in A (= B_\varepsilon(f(x_0))) \\
 &\Rightarrow \underline{\rho(f(x), f(x_0)) < \varepsilon}
 \end{aligned}$$



Thus, we have shown that f is continuous at x_0 ; since x_0 is an arbitrary point in X , f is continuous. □

Continuity in Metric Spaces

The composition of continuous functions is continuous:

Theorem 4 (Slightly weaker version of Thm. 6.10). *Let (X, d_X) , (Y, d_Y) and (Z, d_Z) be metric spaces. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then $g \circ f : X \rightarrow Z$ is continuous.*

Proof. Suppose $A \subseteq Z$ is open. Since g is continuous, $g^{-1}(A)$ is open in Y ; since f is continuous, $f^{-1}(g^{-1}(A))$ is open in X .

We claim that

$$f^{-1}(g^{-1}(A)) = (g \circ f)^{-1}(A)$$

Observe

$$\begin{aligned}x \in f^{-1}(g^{-1}(A)) &\Leftrightarrow f(x) \in g^{-1}(A) \\&\Leftrightarrow g(f(x)) \in A \\&\Leftrightarrow (g \circ f)(x) \in A \\&\Leftrightarrow x \in (g \circ f)^{-1}(A)\end{aligned}$$

which establishes the claim. This shows that $(g \circ f)^{-1}(A)$ is open in X , so $g \circ f$ is continuous. \square

Uniform Continuity

Definition 4 (Uniform Continuity). Suppose $f : (X, d) \rightarrow (Y, \rho)$.
 f is uniformly continuous if

$$\forall \varepsilon > 0 \exists \delta(\varepsilon) > 0 \text{ s.t. } \forall x_0 \in X, d(x, x_0) < \delta(\varepsilon) \Rightarrow \rho(f(x), f(x_0)) < \varepsilon$$

Notice the important contrast with continuity: f is continuous means

$$\forall x_0 \in X, \varepsilon > 0 \exists \delta(x_0, \varepsilon) > 0 \text{ s.t. } d(x, x_0) < \delta(x_0, \varepsilon) \Rightarrow \rho(f(x), f(x_0)) < \varepsilon$$


Uniform Continuity

Example: Consider

$$f(x) = \frac{1}{x}, \quad x \in (0, 1]$$

f is continuous (why?). We will show that f is **not** uniformly continuous.

Fix $\epsilon > 0$, look at x^* .
Then go to x^b



Fix $\varepsilon > 0$ and $x_0 \in (0, 1]$. If $x = \frac{x_0}{1 + \varepsilon x_0}$, then

$$\begin{aligned} 1 + \varepsilon x_0 &> 1 \\ x = \frac{x_0}{1 + \varepsilon x_0} &< x_0 \\ \frac{1}{x} - \frac{1}{x_0} &> 0 \end{aligned}$$

$$\begin{aligned} |f(x) - f(x_0)| &= \left| \frac{1}{x} - \frac{1}{x_0} \right| \\ &= \frac{1}{x} - \frac{1}{x_0} \\ &= \frac{1 + \varepsilon x_0}{x_0} - \frac{1}{x_0} \\ &= \frac{\varepsilon x_0}{x_0} \\ &= \varepsilon \end{aligned}$$

Thus, $\delta(x_0, \varepsilon)$ must be chosen small enough so that

$$\left| \frac{x_0}{1 + \varepsilon x_0} - x_0 \right| \geq \delta(x_0, \varepsilon)$$

$$\begin{aligned} \delta(x_0, \varepsilon) &\leq x_0 - \frac{x_0}{1 + \varepsilon x_0} \\ &= \frac{\varepsilon(x_0)^2}{1 + \varepsilon x_0} \\ &< \varepsilon(x_0)^2 \end{aligned}$$

which converges to zero as $x_0 \rightarrow 0$. So there is no $\delta(\varepsilon)$ that will work for all $x_0 \in (0, 1]$.

Uniform Continuity

$$\exists C > 0 \text{ s.t.}$$

$$|f'(x)| \leq C$$

$$\forall x \in [a, b]$$

Example: If $f : \mathbf{R} \rightarrow \mathbf{R}$ and $f'(x)$ is defined and uniformly bounded on an interval $[a, b]$, then f is uniformly continuous on $[a, b]$. However, even a function with an unbounded derivative may be uniformly continuous. Consider

$$f(x) = \sqrt{x}, \quad x \in [0, 1]$$

f is continuous (why?). We will show that f is uniformly continuous. Given $\varepsilon > 0$, let $\delta = \varepsilon^2$. Then given any $x_0 \in [0, 1]$,

$|x - x_0| < \delta$ implies by the Fundamental Theorem of Calculus

$$\begin{aligned} |f(x) - f(x_0)| &= \left| \int_{x_0}^x \frac{1}{2\sqrt{t}} dt \right| \\ &\leq \int_0^{|x-x_0|} \frac{1}{2\sqrt{t}} dt \\ &= \sqrt{|x - x_0|} \\ &< \sqrt{\delta} \\ &= \sqrt{\varepsilon^2} \\ &= \varepsilon \end{aligned}$$

Thus, f is uniformly continuous on $[0, 1]$, even though $f'(x) \rightarrow \infty$ as $x \rightarrow 0$.

Lipschitz Continuity

Definition 5. Let X, Y be normed vector spaces, $E \subseteq X$. A function $f : X \rightarrow Y$ is Lipschitz on E if

$$\exists K > 0 \text{ s.t. } \|f(x) - f(z)\|_Y \leq K\|x - z\|_X \quad \forall x, z \in E$$

f is locally Lipschitz on E if

$$\forall x_0 \in E \exists \varepsilon > 0 \text{ s.t. } f \text{ is Lipschitz on } B_\varepsilon(x_0) \cap E$$

$\exists K > 0 \forall x \neq z :$

$$\frac{\|f(x) - f(z)\|_Y}{\|x - z\|_X} \leq K$$

Notions of Continuity

Lipschitz continuity is stronger than either continuity or uniform continuity:

Lipschitz \Rightarrow locally Lipschitz \Rightarrow continuous

Lipschitz \Rightarrow uniformly continuous

Every C^1 function is locally Lipschitz. (Recall that a function $f : \mathbf{R}^m \rightarrow \mathbf{R}^n$ is said to be C^1 if all its first partial derivatives exist and are continuous.)

Homeomorphisms

Definition 6. *Let (X, d) and (Y, ρ) be metric spaces. A function $f : X \rightarrow Y$ is called a homeomorphism if it is one-to-one, onto, continuous, and its inverse function is continuous.*

Topological properties are invariant under homeomorphism:

$$\begin{array}{l} \text{identity on } Y \\ \text{"} \\ f \circ g : Y \rightarrow Y \end{array}$$

$$\begin{array}{l} \text{identity on } X \\ \text{"} \\ g \circ f : X \rightarrow X \end{array}$$

Homeomorphisms

$$f: X \rightarrow Y \quad , \quad g = f^{-1} : Y \rightarrow X$$

Suppose that f is a homeomorphism and $U \subset X$. Let $g = f^{-1} : Y \rightarrow X$.

$$y \in (g)^{-1}(U) \Leftrightarrow g(y) \in U$$

$$\Leftrightarrow y \in f(U)$$

$$\underline{U \text{ open in } X} \Rightarrow (g)^{-1}(U) \text{ is open in } \underline{(f(X), \rho)}$$

$$\Rightarrow f(U) \text{ is open in } (f(X), \rho)$$

This says that (X, d) and $(f(X), \rho|_{f(X)})$ are identical in terms of properties that can be characterized solely in terms of open sets; such properties are called "topological properties."