

Econ 204 2016

Lecture 11

Outline

1. Derivatives
2. Chain Rule
3. Mean Value Theorem
4. Taylor's Theorem

Derivatives

Definition 1. Let $f : I \rightarrow \mathbf{R}$, where $I \subseteq \mathbf{R}$ is an open interval. f is differentiable at $x \in I$ if

$$\lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} = a$$

for some $a \in \mathbf{R}$.

This is equivalent to $\exists a \in \mathbf{R}$ such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - (f(x) + ah)}{h} = 0$$

$$\Leftrightarrow \forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } 0 < |h| < \delta \Rightarrow \left| \frac{f(x+h) - (f(x) + ah)}{h} \right| < \varepsilon$$

$$\Leftrightarrow \forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } 0 < |h| < \delta \Rightarrow \frac{|f(x+h) - (f(x) + ah)|}{|h|} < \varepsilon$$

$$\Leftrightarrow \lim_{h \rightarrow 0} \frac{|f(x+h) - (f(x) + ah)|}{|h|} = 0$$

Derivatives

Definition 2. If $X \subseteq \mathbf{R}^n$ is open, $f : X \rightarrow \mathbf{R}^m$ is differentiable at $x \in X$ if $\exists T_x \in L(\mathbf{R}^n, \mathbf{R}^m)$ such that

$$\lim_{h \rightarrow 0, h \in \mathbf{R}^n} \frac{|f(x+h) - (f(x) + T_x(h))|}{|h|} = 0 \quad (1)$$

f is differentiable if it is differentiable at all $x \in X$.

Note that T_x is uniquely determined by Equation (1).

The definition requires that **one** linear operator T_x works no matter how h approaches zero.

In this case, $f(x) + T_x(h)$ is the best linear approximation to $f(x+h)$ for sufficiently small h .

Big-Oh and little-oh

Notation:

- $y = O(|h|^n)$ as $h \rightarrow 0$ – read “ y is big-Oh of $|h|^n$ ” – means

$$\exists K, \delta > 0 \text{ s.t. } |h| < \delta \Rightarrow |y| \leq K|h|^n$$

- $y = o(|h|^n)$ as $h \rightarrow 0$ – read “ y is little-oh of $|h|^n$ ” – means

$$\lim_{h \rightarrow 0} \frac{|y|}{|h|^n} = 0$$

Note that $y = O(|h|^{n+1})$ as $h \rightarrow 0$ implies $y = o(|h|^n)$ as $h \rightarrow 0$.

Using this notation: f is differentiable at $x \Leftrightarrow \exists T_x \in L(\mathbf{R}^n, \mathbf{R}^m)$
such that

$$f(x + h) = f(x) + T_x(h) + o(h) \text{ as } h \rightarrow 0$$

More Notation

Notation:

- df_x is the linear transformation T_x
- $Df(x)$ is the matrix of df_x with respect to the standard basis.
This is called the *Jacobian* or *Jacobian matrix* of f at x
- $E_f(h) = f(x + h) - (f(x) + df_x(h))$ is the *error term*

Using this notation,

f is differentiable at $x \Leftrightarrow E_f(h) = o(h)$ as $h \rightarrow 0$

What's $Df(x)$?

Now compute $Df(x) = (a_{ij})$. Let $\{e_1, \dots, e_n\}$ be the standard basis of \mathbf{R}^n . Look in direction e_j (note that $|\gamma e_j| = |\gamma|$).

$$\begin{aligned} o(\gamma) &= f(x + \gamma e_j) - (f(x) + T_x(\gamma e_j)) \\ &= f(x + \gamma e_j) - \left(f(x) + \begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & \cdots & \vdots & \cdots & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \gamma \\ 0 \\ \vdots \\ 0 \end{pmatrix} \right) \\ &= f(x + \gamma e_j) - \left(f(x) + \begin{pmatrix} \gamma a_{1j} \\ \vdots \\ \gamma a_{mj} \end{pmatrix} \right) \end{aligned}$$

For $i = 1, \dots, m$, let f^i denote the i^{th} component of the function f :

$$f^i(x + \gamma e_j) - (f^i(x) + \gamma a_{ij}) = o(\gamma)$$
$$\text{so } a_{ij} = \frac{\partial f^i}{\partial x_j}(x)$$

Derivatives and Partial Derivatives

Theorem 1 (Thm. 3.3). *Suppose $X \subseteq \mathbf{R}^n$ is open and $f : X \rightarrow \mathbf{R}^m$ is differentiable at $x \in X$. Then $\frac{\partial f^i}{\partial x_j}(x)$ exists for $1 \leq i \leq m$, $1 \leq j \leq n$, and*

$$Df(x) = \begin{pmatrix} \frac{\partial f^1}{\partial x_1}(x) & \cdots & \frac{\partial f^1}{\partial x_n}(x) \\ \vdots & \cdots & \vdots \\ \frac{\partial f^m}{\partial x_1}(x) & \cdots & \frac{\partial f^m}{\partial x_n}(x) \end{pmatrix}$$

i.e. the Jacobian at x is the matrix of partial derivatives at x .

Derivatives and Partial Derivatives

Remark: If f is differentiable at x , then all first-order partial derivatives $\frac{\partial f^i}{\partial x_j}$ exist at x . However, the converse is false: existence of all the first-order partial derivatives does not imply that f is differentiable.

The missing piece is continuity of the partial derivatives:

Theorem 2 (Thm. 3.4). *If all the first-order partial derivatives $\frac{\partial f^i}{\partial x_j}$ ($1 \leq i \leq m$, $1 \leq j \leq n$) exist and are continuous at x , then f is differentiable at x .*

Directional Derivatives

Suppose $X \subseteq \mathbf{R}^n$ open, $f : X \rightarrow \mathbf{R}^m$ is differentiable at x , and $|u| = 1$.

$$\begin{aligned} f(x + \gamma u) - (f(x) + T_x(\gamma u)) &= o(\gamma) \text{ as } \gamma \rightarrow 0 \\ \Rightarrow f(x + \gamma u) - (f(x) + \gamma T_x(u)) &= o(\gamma) \text{ as } \gamma \rightarrow 0 \\ \Rightarrow \lim_{\gamma \rightarrow 0} \frac{f(x + \gamma u) - f(x)}{\gamma} &= T_x(u) = Df(x)u \end{aligned}$$

i.e. the directional derivative in the direction u (with $|u| = 1$) is

$$Df(x)u \in \mathbf{R}^m$$

Chain Rule

Theorem 3 (Thm. 3.5, Chain Rule). *Let $X \subseteq \mathbf{R}^n$, $Y \subseteq \mathbf{R}^m$ be open, $f : X \rightarrow Y$, $g : Y \rightarrow \mathbf{R}^p$. Let $x_0 \in X$ and $F = g \circ f$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$, then $F = g \circ f$ is differentiable at x_0 and*

$$dF_{x_0} = dg_{f(x_0)} \circ df_{x_0}$$

(composition of linear transformations)

$$DF(x_0) = Dg(f(x_0))Df(x_0)$$

(matrix multiplication)

Remark: The statement is exactly the same as in the univariate case, except we replace the univariate derivative by a linear transformation. The proof is more or less the same, with a bit of linear algebra added.

Mean Value Theorem

Theorem 4 (Thm. 1.7, Mean Value Theorem, Univariate Case).
Let $a, b \in \mathbf{R}$. Suppose $f : [a, b] \rightarrow \mathbf{R}$ is continuous on $[a, b]$ and differentiable on (a, b) . Then there exists $c \in (a, b)$ such that

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

that is, such that

$$f(b) - f(a) = f'(c)(b - a)$$

Proof. Consider the function

$$g(x) = f(x) - f(a) - \frac{f(b) - f(a)}{b - a}(x - a)$$

Then $g(a) = 0 = g(b)$. Note that for $x \in (a, b)$,


$$g'(x) = f'(x) - \frac{f(b) - f(a)}{b - a}$$

so it suffices to find $c \in (a, b)$ such that $g'(c) = 0$.

Case I: If $g(x) = 0$ for all $x \in [a, b]$, choose an arbitrary $c \in (a, b)$, and note that $g'(c) = 0$, so we are done.

Case II: Suppose $g(x) > 0$ for some $x \in [a, b]$. Since g is continuous on $[a, b]$, it attains its maximum at some point $c \in (a, b)$. Since g is differentiable at c and c is an interior point of the domain of g , we have $g'(c) = 0$, and we are done.

Case III: If $g(x) < 0$ for some $x \in [a, b]$, the argument is similar to that in Case II. □



Mean Value Theorem

Notation:

$$\ell(x, y) = \{\alpha x + (1 - \alpha)y : \alpha \in [0, 1]\}$$

is the line segment from x to y .

Theorem 5 (Mean Value Theorem). *Suppose $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is differentiable on an open set $X \subseteq \mathbf{R}^n$, $x, y \in X$ and $\ell(x, y) \subseteq X$. Then there exists $z \in \ell(x, y)$ such that*

$$f(y) - f(x) = Df(z)(y - x)$$

Notice that the statement is exactly the same as in the univariate case. For $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$, we can apply the Mean Value Theorem to each component, to obtain $z_1, \dots, z_m \in \ell(x, y)$ such that

$$f^i(y) - f^i(x) = Df^i(z_i)(y - x)$$

However, we cannot find a single z which works for every component.

Note that each $z_i \in \ell(x, y) \subset \mathbf{R}^n$; there are m of them, one for each component in the range.

Mean Value Theorem

Theorem 6. *Suppose $X \subset \mathbf{R}^n$ is open and $f : X \rightarrow \mathbf{R}^m$ is differentiable. If $x, y \in X$ and $\ell(x, y) \subseteq X$, then there exists $z \in \ell(x, y)$ such that*

$$\begin{aligned} |f(y) - f(x)| &\leq |df_z(y - x)| \\ &\leq \|df_z\| |y - x| \end{aligned}$$

Mean Value Theorem

Remark: To understand why we don't get equality, consider $f : [0, 1] \rightarrow \mathbf{R}^2$ defined by

$$f(t) = (\cos 2\pi t, \sin 2\pi t)$$

f maps $[0, 1]$ to the unit circle in \mathbf{R}^2 . Note that $f(0) = f(1) = (1, 0)$, so $|f(1) - f(0)| = 0$. However, for any $z \in [0, 1]$,

$$\begin{aligned} |df_z(1 - 0)| &= |2\pi(-\sin 2\pi z, \cos 2\pi z)| \\ &= 2\pi\sqrt{\sin^2 2\pi z + \cos^2 2\pi z} \\ &= 2\pi \end{aligned}$$

Taylor's Theorem – \mathbf{R}

Theorem 7 (Thm. 1.9, Taylor's Theorem in \mathbf{R}). *Let $f : I \rightarrow \mathbf{R}$ be n -times differentiable, where $I \subseteq \mathbf{R}$ is an open interval. If $x, x + h \in I$, then*

$$f(x + h) = f(x) + \sum_{k=1}^{n-1} \frac{f^{(k)}(x)h^k}{k!} + E_n$$

where $f^{(k)}$ is the k^{th} derivative of f and

$$E_n = \frac{f^{(n)}(x + \lambda h)h^n}{n!} \text{ for some } \lambda \in (0, 1)$$

Motivation: Let

$$\begin{aligned}T_n(h) &= f(x) + \sum_{k=1}^n \frac{f^{(k)}(x)h^k}{k!} \\&= f(x) + f'(x)h + \frac{f''(x)h^2}{2} + \dots + \frac{f^{(n)}(x)h^n}{n!} \\T_n(0) &= f(x) \\T'_n(h) &= f'(x) + f''(x)h + \dots + \frac{f^{(n)}(x)h^{n-1}}{(n-1)!} \\T'_n(0) &= f'(x) \\T''_n(h) &= f''(x) + \dots + \frac{f^{(n)}(x)h^{n-2}}{(n-2)!} \\T''_n(0) &= f''(x) \\&\vdots \\T_n^{(n)}(0) &= f^{(n)}(x)\end{aligned}$$

so $T_n(h)$ is the unique n^{th} degree polynomial such that

$$\begin{aligned}T_n(0) &= f(x) \\T'_n(0) &= f'(x) \\&\vdots \\T_n^{(n)}(0) &= f^{(n)}(x)\end{aligned}$$

Taylor's Theorem – \mathbf{R}

Theorem 8 (Alternate Taylor's Theorem in \mathbf{R}). *Let $f : I \rightarrow \mathbf{R}$ be n times differentiable, where $I \subseteq \mathbf{R}$ is an open interval and $x \in I$. Then*

$$f(x + h) = f(x) + \sum_{k=1}^n \frac{f^{(k)}(x)h^k}{k!} + o(h^n) \text{ as } h \rightarrow 0$$

If f is $(n + 1)$ times continuously differentiable, then

$$f(x + h) = f(x) + \sum_{k=1}^n \frac{f^{(k)}(x)h^k}{k!} + O(h^{n+1}) \text{ as } h \rightarrow 0$$

Remark: The first equation in the statement of the theorem is essentially a restatement of the definition of the n^{th} derivative. The second statement is proven from Theorem 1.9, and the continuity of the derivative.

C^k Functions

Definition 3. Let $X \subseteq \mathbf{R}^n$ be open. A function $f : X \rightarrow \mathbf{R}^m$ is continuously differentiable on X if

- f is differentiable on X and
- df_x is a continuous function of x from X to $L(\mathbf{R}^n, \mathbf{R}^m)$, with respect to the operator norm $\|df_x\|$

f is C^k if all partial derivatives of order $\leq k$ exist and are continuous in X .

C^k Functions

Theorem 9 (Thm. 4.3). *Suppose $X \subseteq \mathbf{R}^n$ is open and $f : X \rightarrow \mathbf{R}^m$. Then f is continuously differentiable on X if and only if f is C^1 .*

Taylor's Theorem – Linear Terms

Theorem 10. *Suppose $X \subseteq \mathbf{R}^n$ is open and $x \in X$. If $f : X \rightarrow \mathbf{R}^m$ is differentiable, then*

$$f(x + h) = f(x) + Df(x)h + o(h) \text{ as } h \rightarrow 0$$

This is essentially a restatement of the definition of differentiability.

Taylor's Theorem – Linear Terms

Theorem 11 (Corollary of 4.4). *Suppose $X \subseteq \mathbf{R}^n$ is open and $x \in X$. If $f : X \rightarrow \mathbf{R}^m$ is C^2 , then*

$$f(x + h) = f(x) + Df(x)h + O(|h|^2) \text{ as } h \rightarrow 0$$

Taylor's Theorem – Quadratic Terms

We treat each component of the function separately, so consider $f : X \rightarrow \mathbf{R}$, $X \subseteq \mathbf{R}^n$ an open set. Let

$$D^2 f(x) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2}(x) & \frac{\partial^2 f}{\partial x_2 \partial x_1}(x) & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_1}(x) \\ \frac{\partial^2 f}{\partial x_1 \partial x_2}(x) & \frac{\partial^2 f}{\partial x_2^2}(x) & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_2}(x) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_1 \partial x_n}(x) & \cdots & \cdots & \frac{\partial^2 f}{\partial x_n^2}(x) \end{pmatrix}$$

$$f \in C^2 \Rightarrow \frac{\partial^2 f}{\partial x_i \partial x_j}(x) = \frac{\partial^2 f}{\partial x_j \partial x_i}(x)$$

$\Rightarrow D^2 f(x)$ is symmetric

$\Rightarrow D^2 f(x)$ has eigenvectors that are an orthonormal basis and thus can be diagonalized

Taylor's Theorem – Quadratic Terms

Theorem 12 (Stronger Version of Thm. 4.4). *Let $X \subseteq \mathbf{R}^n$ be open, $f : X \rightarrow \mathbf{R}$, $f \in C^2(X)$, and $x \in X$. Then*

$$f(x + h) = f(x) + Df(x)h + \frac{1}{2}h^\top (D^2f(x))h + o(|h|^2) \text{ as } h \rightarrow 0$$

If $f \in C^3$,

$$f(x + h) = f(x) + Df(x)h + \frac{1}{2}h^\top (D^2f(x))h + O(|h|^3) \text{ as } h \rightarrow 0$$

Characterizing Critical Points

Definition 4. *We say f has a saddle at x if $Df(x) = 0$ but f has neither a local maximum nor a local minimum at x .*

Characterizing Critical Points

Corollary 1. *Suppose $X \subseteq \mathbf{R}^n$ is open and $x \in X$. If $f : X \rightarrow \mathbf{R}$ is C^2 , there is an orthonormal basis $\{v_1, \dots, v_n\}$ and corresponding eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbf{R}$ of $D^2f(x)$ such that*

$$\begin{aligned} f(x + h) &= f(x + \gamma_1 v_1 + \dots + \gamma_n v_n) \\ &= f(x) + \sum_{i=1}^n (Df(x)v_i) \gamma_i + \frac{1}{2} \sum_{i=1}^n \lambda_i \gamma_i^2 + o(|\gamma|^2) \end{aligned}$$

where $\gamma_i = h \cdot v_i$.

1. If $f \in C^3$, we may strengthen $o(|\gamma|^2)$ to $O(|\gamma|^3)$.

2. If f has a local maximum or local minimum at x , then

$$Df(x) = 0$$

3. If $Df(x) = 0$, then

- $\lambda_1, \dots, \lambda_n > 0 \Rightarrow f$ has a local minimum at x
- $\lambda_1, \dots, \lambda_n < 0 \Rightarrow f$ has a local maximum at x
- $\lambda_i < 0$ for some i , $\lambda_j > 0$ for some $j \Rightarrow f$ has a saddle at x
- $\lambda_1, \dots, \lambda_n \geq 0$, $\lambda_i > 0$ for some $i \Rightarrow f$ has a local minimum or a saddle at x
- $\lambda_1, \dots, \lambda_n \leq 0$, $\lambda_i < 0$ for some $i \Rightarrow f$ has a local maximum or a saddle at x
- $\lambda_1 = \dots = \lambda_n = 0$ gives no information.

Proof. (Sketch) From our study of quadratic forms, we know the behavior of the quadratic terms is determined by the signs of the eigenvalues. If $\lambda_i = 0$ for some i , then we know that the quadratic form arising from the second partial derivatives is identically zero in the direction v_i , and the higher derivatives will determine the behavior of the function f in the direction v_i . For example, if $f(x) = x^3$, then $f'(0) = 0$, $f''(0) = 0$, but we know that f has a saddle at $x = 0$; however, if $f(x) = x^4$, then again $f'(0) = 0$ and $f''(0) = 0$ but f has a local (and global) minimum at $x = 0$. □