

Econ 204 2020

Lecture 10

Outline

1. Diagonalization of Real Symmetric Matrices
2. Application to Quadratic Forms
3. Linear Maps Between Normed Spaces

Announcements

- PS 3 due today 1pm
→ solns posted
≈ 2 pm today
- PS 4 due
Tuesday at 1pm
- last year's exam
posted ≈ Sunday
- Zoom details
for next
week later
today

How Might This Matter

$$\begin{aligned}c_{t+1} &= b_{11}c_t + b_{12}k_t \\k_{t+1} &= b_{21}c_t + b_{22}k_t\end{aligned}\quad \forall t$$

- Why does diagonalizability matter?

Consider a two-dimensional linear difference equation:

$$\begin{pmatrix} c_{t+1} \\ k_{t+1} \end{pmatrix} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \begin{pmatrix} c_t \\ k_t \end{pmatrix} \quad \forall t = 0, 1, 2, 3, \dots$$

given an initial condition c_0, k_0 , or, setting

$$y_t = \begin{pmatrix} c_t \\ k_t \end{pmatrix} \quad \forall t \quad \text{and} \quad B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$

we can rewrite this more compactly as

$$y_{t+1} = By_t \quad \forall t$$

where $b_{ij} \in \mathbf{R}$ each i, j .

We want to find a solution y_t , $t = 1, 2, 3, \dots$ given initial condition y_0 . (Why?)

Such a dynamical system will arise for example as a characterization of the solution to a standard infinite-horizon optimal growth problem (202a, lecture 2).

If B is diagonalizable, this can be easily solved after a change of basis. If B is diagonalizable, choose an invertible 2×2 real matrix P such that

$$P^{-1}BP = D = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}$$

Then

$$\begin{aligned}
 y_{t+1} = By_t \quad \forall t &\iff P^{-1}y_{t+1} = P^{-1}By_t \quad \forall t && \text{(mult. by } P^{-1}\text{)} \\
 &\iff (P^{-1}y_{t+1}) = (P^{-1}BP)(P^{-1}y_t) \quad \forall t && PP^{-1} = I \\
 &\iff \bar{y}_{t+1} = D\bar{y}_t \quad \forall t \\
 & && = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix} \bar{y}_t \\
 \text{where } \bar{y}_t &= P^{-1}y_t \quad \forall t
 \end{aligned}$$

$$\Leftrightarrow \bar{y}_{i,t+1} = d_i \bar{y}_{i,t} \quad \forall t, \quad i=1,2$$

where $\bar{y}_t = P^{-1}y_t \quad \forall t$.

Since D is diagonal, after a change of basis to \bar{y}_t , we need to solve two **independent** linear univariate difference equations, which is easy:

$$\bar{y}_{i,t} = d_i^t \bar{y}_{i,0} \quad \forall t \quad i=1,2$$

- Not all real $n \times n$ matrices are diagonalizable (not even all invertible $n \times n$ matrices are)...so can we identify some classes that are? *yesterday:*
 - basis of eigenvectors (\Leftrightarrow)
 - n distinct eigenvalues (\Rightarrow)
- Some types of matrices appear more frequently than others – especially real symmetric $n \times n$ matrices (matrix representation of second derivatives of C^2 functions, quadratic forms...). *e.g. second order conditions in optimization, checking concavity and convexity, Taylor series approximation of function*

- Recall that an $n \times n$ real matrix A is *symmetric* if $a_{ij} = a_{ji}$ for all i, j , where a_{ij} is the $(i, j)^{\text{th}}$ entry of A .

Rest of this section: work in \mathbb{R}^n

- vector space

- norm

- inner product

$$(x \cdot y) = \sum_{i=1}^n x_i y_i$$

Orthonormal Bases

Definition 1. Let

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

A basis $V = \{v_1, \dots, v_n\}$ of \mathbb{R}^n is orthonormal if $v_i \cdot v_j = \delta_{ij} = \begin{cases} 1 & i=j \\ 0 & i \neq j \end{cases}$

In other words, a basis is orthonormal if each basis element has unit length ($\|v_i\|^2 = v_i \cdot v_i = 1 \ \forall i$), and distinct basis elements are perpendicular ($v_i \cdot v_j = 0$ for $i \neq j$).

$$\|x\| = \left(\sum_{i=1}^n x_i^2 \right)^{1/2} = (x \cdot x)^{1/2}$$

Orthonormal Bases

Remark: Suppose that $x = \sum_{j=1}^n \alpha_j v_j$ where $\{v_1, \dots, v_n\}$ is an orthonormal basis of \mathbf{R}^n . Then

$$\begin{aligned} x \cdot v_k &= \left(\sum_{j=1}^n \alpha_j v_j \right) \cdot v_k \\ &= \sum_{j=1}^n \alpha_j (v_j \cdot v_k) \\ &= \sum_{j=1}^n \alpha_j \delta_{jk} = \begin{cases} \alpha_k & j=k \\ 0 & j \neq k \end{cases} \\ &= \alpha_k \end{aligned}$$

so

$$x = \sum_{j=1}^n (x \cdot v_j) v_j$$

Orthonormal Bases

Example: The standard basis of \mathbf{R}^n is orthonormal.


$$e_i = (0, \dots, 1, \dots, 0) \quad i = 1, \dots, n$$

(Why?)

e.g. \mathbb{R}^2 : $e_1 = (1, 0)$, $e_2 = (0, 1)$

others? e.g. $v_1 = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$, $v_2 = \left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$

also many bases that are not orthonormal



Unitary Matrices

Recall that for a real $n \times m$ matrix A , A^\top denotes the transpose of A : the $(i, j)^{th}$ entry of A^\top is the $(j, i)^{th}$ entry of A .

So the i^{th} row of A^\top is the i^{th} column of A .

Definition 2. A real $n \times n$ matrix A is unitary if $A^\top = A^{-1}$.

Notice that by definition every unitary matrix is invertible.

Unitary Matrices

Theorem 1. *A real $n \times n$ matrix A is unitary if and only if the columns of A are orthonormal.*

Proof. Let v_j denote the j^{th} column of A .

$$\begin{aligned} A^T &= A^{-1} && \iff A^T A = I && = (\delta_{ij}) = \begin{cases} 1 & i=j \\ 0 & i \neq j \end{cases} \\ &&& \iff v_i \cdot v_j = \delta_{ij} \quad \forall i, j \\ &&& \iff \{v_1, \dots, v_n\} \text{ is orthonormal} \end{aligned}$$

□

$$A = \begin{pmatrix} v_1 & \dots & v_n \end{pmatrix} = \text{Mtx}_{W,V}(\text{id})$$

$\begin{matrix} \text{ord}_W(v_1) & \dots & \text{ord}_W(v_n) \end{matrix}$

$$= \text{ord}_W \left(\begin{pmatrix} 1 & 2 \end{pmatrix} \right)$$

$$= 1 \cdot \begin{pmatrix} 1 & 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 & 1 \end{pmatrix}$$

Unitary Matrices

If A is unitary, let V be the set of columns of A and W be the standard basis of \mathbf{R}^n . Since A is unitary, it is invertible, so V is a basis of \mathbf{R}^n . ($\{v_1, \dots, v_n\}$ linearly independent)

$$A^T = A^{-1} = \text{Mtx}_{V,W}(\text{id}) = \text{change of basis from } W \text{ to } V$$

\uparrow
 standard basis

Since V is orthonormal, the transformation between bases W and V preserves all geometry, including lengths and angles.

Thus : Let C be an $n \times n$ real symmetric matrix.
Then C is diagonalizable. In addition,

$$C = P^{-1} D P$$

where D is a diagonal matrix and P is unitary.

Note : The diagonal elements $\{\lambda_1, \dots, \lambda_n\}$ of D
are the eigenvalues of C

- C has orthonormal eigenvectors $\{v_1, \dots, v_n\}$
that are a basis for \mathbb{R}^n .

Diagonalization of Real Symmetric Matrices

Theorem 2. *Let $T \in L(\mathbf{R}^n, \mathbf{R}^n)$ and W be the standard basis of \mathbf{R}^n . Suppose that $Mtx_W(T)$ is symmetric. Then the eigenvectors of T are all real, and there is an orthonormal basis $V = \{v_1, \dots, v_n\}$ of \mathbf{R}^n consisting of eigenvectors of T , so that $Mtx_W(T)$ is diagonalizable:*

$$Mtx_W(T) = Mtx_{W,V}(id) \cdot Mtx_V(T) \cdot Mtx_{V,W}(id)$$

where $Mtx_V T$ is diagonal and the change of basis matrices $Mtx_{V,W}(id)$ and $Mtx_{W,V}(id)$ are unitary.

The proof of the theorem requires a lengthy digression into the linear algebra of *complex* vector spaces. A brief outline is in the notes.

quadratic form: polynomial with all terms of degree 2

Quadratic Forms

Example: Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$

$$f(x) = \alpha x_1^2 + \beta x_1 x_2 + \gamma x_2^2$$

Let write as $f(x) = x^T A x$, A symmetric

$$A = \begin{pmatrix} \alpha & \frac{\beta}{2} \\ \frac{\beta}{2} & \gamma \end{pmatrix}$$

$$x^T A x = (x_1 \ x_2) \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

so A is symmetric and

$$\begin{aligned}x^{\top}Ax &= (x_1, x_2) \begin{pmatrix} \alpha & \frac{\beta}{2} \\ \frac{\beta}{2} & \gamma \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= (x_1, x_2) \begin{pmatrix} \alpha x_1 + \frac{\beta}{2}x_2 \\ \frac{\beta}{2}x_1 + \gamma x_2 \end{pmatrix} \\ &= \alpha x_1^2 + \beta x_1 x_2 + \gamma x_2^2 \\ &= f(x)\end{aligned}$$

Notice $f(0) = 0$.

Can we determine anything about $f(x)$ for $x \neq 0$?

e.g. $f(x) \geq 0 \quad \forall x$?

easy if $\beta = 0 \dots$

Quadratic Forms

general form:

Consider a quadratic form

$$f(x_1, \dots, x_n) = \sum_{i=1}^n \alpha_{ii} x_i^2 + \sum_{i < j} \beta_{ij} x_i x_j \quad (1)$$

Let

$$\alpha_{ij} = \begin{cases} \frac{\beta_{ij}}{2} & \text{if } i < j \quad \leftarrow \text{above diagonal} \\ \frac{\beta_{ji}}{2} & \text{if } i > j \quad \leftarrow \text{below diagonal} \end{cases}$$

Let

$$A = \begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & \ddots & \vdots \\ \alpha_{n1} & \cdots & \alpha_{nn} \end{pmatrix} \quad \text{so } f(x) = x^T A x$$

\uparrow
real symmetric

Quadratic Forms

A is symmetric, so let $V = \{v_1, \dots, v_n\}$ be an orthonormal basis of eigenvectors of A with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$.

$$\text{Then } A = U^T D U \quad = \omega^{-1} \Omega \omega$$

$$\text{where } D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$$

ω and $U = \text{Mtx}_{V,W}(\text{id})$ is unitary

The columns of U^T (the rows of U) are the coordinates of v_1, \dots, v_n , expressed in terms of the standard basis W . Given $x \in \mathbf{R}^n$, recall

$$x = \sum_{i=1}^n \gamma_i v_i \quad \text{where } \gamma_i = x \cdot v_i$$

Quadratic Forms

So

$$\begin{aligned} f(\gamma) = f(x) &= f\left(\sum \gamma_i v_i\right) \\ &= \left(\sum \gamma_i v_i\right)^\top A \left(\sum \gamma_i v_i\right) = x^\top A x \\ &= \left(\sum \gamma_i v_i\right)^\top U^\top D U \left(\sum \gamma_i v_i\right) \\ &= \left(U \sum \gamma_i v_i\right)^\top D \left(U \sum \gamma_i v_i\right) && ((EF)^\top = F^\top E^\top) \\ &= \left(\sum \gamma_i U v_i\right)^\top D \left(\sum \gamma_i U v_i\right) && (U \text{ linear}) \\ &= (\gamma_1, \dots, \gamma_n) D \begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix} \leftarrow U \text{ is change of} \\ &= \sum \lambda_i \gamma_i^2 && \text{basis matrix} \\ & && \text{from } W \text{ to } V \\ & && \Rightarrow U v_i = e_i \quad \forall i \end{aligned}$$

↑
eigenvalues
of A

Quadratic Forms

The equation for a level set of f is

$$\{\gamma \in \mathbb{R}^n : f(\gamma) = C\} = \left\{ \gamma \in \mathbb{R}^n : \sum_{i=1}^n \lambda_i \gamma_i^2 = C \right\} \quad C \in \mathbb{R}$$

- If $\lambda_i \geq 0$ for all i , the level set is an ellipsoid, with principal axes in the directions v_1, \dots, v_n . The length of the principal axis along v_i is $\sqrt{C/\lambda_i}$ if $C \geq 0$ (if $\lambda_i = 0$, the level set is a degenerate ellipsoid with principal axis of infinite length in that direction). The level set is empty if $C < 0$.

$\Rightarrow f$ has global min at 0, $f(x) \geq 0 \quad \forall x$

- If $\lambda_i \leq 0$ for all i , the level set is an ellipsoid, with principal axes in the directions v_1, \dots, v_n . The length of the principal

$\Rightarrow f$ has global max at 0, $f(x) \leq 0 \quad \forall x$

axis along v_i is $\sqrt{|C/\lambda_i|}$ if $C \leq 0$ (if $\lambda_i = 0$, the level set is a degenerate ellipsoid with principal axis of infinite length in that direction). The level set is empty if $C > 0$.

- If $\lambda_i > 0$ for some i and $\lambda_j < 0$ for some j , the level set is a hyperboloid. For example, suppose $n = 2$, $\lambda_1 > 0$, $\lambda_2 < 0$. The equation is

$$\begin{aligned} C &= \lambda_1 \gamma_1^2 + \lambda_2 \gamma_2^2 \\ &= \left(\sqrt{\lambda_1 \gamma_1} + \sqrt{|\lambda_2| \gamma_2} \right) \left(\sqrt{\lambda_1 \gamma_1} - \sqrt{|\lambda_2| \gamma_2} \right) \end{aligned}$$

$\Rightarrow f$ has a saddle point at 0
min with respect to v_i
max with respect to v_j

This is a hyperbola with asymptotes


$$0 = \sqrt{\lambda_1 \gamma_1} + \sqrt{|\lambda_2| \gamma_2}$$

$$\Rightarrow \gamma_1 = -\sqrt{\frac{|\lambda_2|}{\lambda_1}} \gamma_2$$

$$0 = \left(\sqrt{\lambda_1 \gamma_1} - \sqrt{|\lambda_2| \gamma_2} \right)$$

$$\Rightarrow \gamma_1 = \sqrt{\frac{|\lambda_2|}{\lambda_1}} \gamma_2$$


$$\lambda_1 > 0, \lambda_2 > 0$$



f has a global min at 0

$$\lambda_1 > 0, \lambda_2 < 0$$

$$v_1 = \sqrt{|\lambda_2|/\lambda_1} \gamma_2$$



$$\gamma_1 = -\sqrt{\frac{|\lambda_1|}{\lambda_2}} \gamma_2$$

$$\{x \in \mathbb{R}^n : f(x) = c\}$$

f has a saddle point at 0

Quadratic Forms

$$f(x) = x^T A x$$

This proves the following corollary of Theorem 2.

Corollary 1. Consider the quadratic form (1). Let $\{v_1, \dots, v_n\}$ be an orthonormal basis of eigenvectors of A with corresponding eigenvalues $\{\lambda_1, \dots, \lambda_n\}$

1. f has a global minimum at 0 if and only if $\lambda_i \geq 0$ for all i ; the level sets of f are ellipsoids with principal axes aligned with the orthonormal eigenvectors v_1, \dots, v_n .
2. f has a global maximum at 0 if and only if $\lambda_i \leq 0$ for all i ; the level sets of f are ellipsoids with principal axes aligned with the orthonormal eigenvectors v_1, \dots, v_n .

3. *If $\lambda_i < 0$ for some i and $\lambda_j > 0$ for some j , then f has a saddle point at 0 ; the level sets of f are hyperboloids with principal axes aligned with the orthonormal eigenvectors v_1, \dots, v_n .*

Bounded Linear Maps

(over \mathbb{R})

Definition 3. Suppose X, Y are normed vector spaces and $T \in L(X, Y)$. We say T is bounded if

$$\exists \beta \in \mathbf{R} \text{ s.t. } \|T(x)\|_Y \leq \beta \|x\|_X \quad \forall x \in X$$

Note this implies that T is Lipschitz with constant β .

why not previous notion of bounded:
 $\exists B \in \mathbb{R}$ s.t. $\|T(x)\| \leq B \quad \forall x$??

$$T(\alpha x) = \alpha T(x) \quad \forall \alpha \in \mathbb{R}$$
$$\|T(\alpha x)\| = \|\alpha T(x)\| = |\alpha| \|T(x)\|$$

Bounded Linear Maps

Much more is true:

Theorem 3 (Thms. 4.1, 4.3). *Let X and Y be normed vector spaces and $T \in L(X, Y)$. Then*

- T is continuous at some point $x_0 \in X$*
- $\iff T$ is continuous at every $x \in X$*
- $\iff T$ is uniformly continuous on X*
- $\iff T$ is Lipschitz*
- $\iff T$ is bounded*

Proof. Suppose T is continuous at x_0 . Fix $\varepsilon > 0$. Then there exists $\delta > 0$ such that

$$\|z - x_0\| < \delta \Rightarrow \|T(z) - T(x_0)\| < \varepsilon$$

Now suppose x is any element of X . If $\|y - x\| < \delta$, let $z =$
 $z = y - x + x_0$, so $\|z - x_0\| = \|y - x\| < \delta$.

$$\underbrace{z = y - x + x_0}_{z - x_0 = y - x}$$

$$\begin{aligned} & \|T(y) - T(x)\| \\ &= \|T(y - x)\| && (T \text{ linear}) \\ &= \|T(y - x + x_0 - x_0)\| = \|T(z - x_0)\| \\ &= \|T(z) - T(x_0)\| && (T \text{ linear}) \\ &< \varepsilon \end{aligned}$$

which proves that T is continuous at every x , and uniformly continuous.

We claim that T is bounded if and only if T is continuous at 0. Suppose T is not bounded. Then

$$\exists \{x_n\} \text{ s.t. } \|T(x_n)\| > n\|x_n\| \quad \forall n$$

Note that $x_n \neq 0$. Let $\varepsilon = 1$. Fix $\delta > 0$ and choose n such that $\frac{1}{n} < \delta$. Let

$$\begin{aligned}
 x'_n &= \frac{x_n}{n\|x_n\|} = \frac{1}{n} \frac{x_n}{\|x_n\|} \\
 \|x'_n - 0\| &= \|x'_n\| = \frac{\|x_n\|}{n\|x_n\|} \\
 &= \frac{1}{n} \\
 &< \delta \\
 \|T(x'_n) - T(0)\| &= \|T(x'_n)\| \\
 &= \frac{1}{n\|x_n\|} \|T(x_n)\| \quad \left(\begin{array}{l} \text{defn of } x'_n \\ + T \text{ linear} \end{array} \right) \\
 &> \frac{n\|x_n\|}{n\|x_n\|} \quad \left(\text{defn of } x_n \right) \\
 &= 1 \\
 &= \varepsilon
 \end{aligned}$$

Since this is true for every δ , T is not continuous at 0. Therefore, T continuous at 0 implies T is bounded. Now, suppose T is bounded, so find $M > 0$ such that $\|T(x)\| \leq M\|x\|$ for every $x \in X$. Given $\varepsilon > 0$, let $\delta = \varepsilon/M$. Then

$$\begin{aligned}\|x - 0\| < \delta &\Rightarrow \|x\| < \delta \\ &\Rightarrow \|T(x) - T(0)\| = \|T(x)\| < M\delta \quad (\text{def of } M) \\ &\Rightarrow \|T(x) - T(0)\| < \varepsilon = M\delta\end{aligned}$$

so T is continuous at 0.

Thus, we have shown that continuity at some point x_0 implies uniform continuity, which implies continuity at every point, which implies T is continuous at 0, which implies that T is bounded, which implies that T is continuous at 0, which implies that T is

continuous at some x_0 , so all of the statements except possibly the Lipschitz statement are equivalent.

Suppose T is bounded, with constant M .^{so} Then

$$\begin{aligned}\|T(x) - T(y)\| &= \|T(x - y)\| && (T \text{ linear}) \\ &\leq M\|x - y\|\end{aligned}$$

so T is Lipschitz with constant M ; conversely, if T is Lipschitz with constant M , then T is bounded with constant M . So all the statements are equivalent. \square

$$\forall x \in X: \|T(x) - T(0)\| = \|T(x)\| \leq M\|x - 0\| = M\|x\|$$

C_{00} - sequences of real numbers with finitely many entries $\neq 0$

Bounded Linear Maps

Every linear map on a finite-dimensional normed vector space is bounded (and thus continuous, uniformly continuous, and Lipschitz continuous).

Theorem 4 (Thm. 4.5). Let X and Y be normed vector spaces, with $\dim X = n$. Every $T \in L(X, Y)$ is bounded.

($n \in \mathbb{N}$)

Proof. See de la Fuente. □

$$T: C_{00} \rightarrow C_{00}$$

$$\|x\| = \sup_n |x_n|$$

$$T(x_1, x_2, \dots) = (x_1, 2x_2, 3x_3, \dots)$$

$$y^n = (1, 1, 1, \dots, \underset{\uparrow}{1}, 0, 0, \dots)$$

$$\|y^n\| = 1 \quad \text{for } n \geq 1$$

$$T(y^n) = (1, 2, 3, \dots, n, 0, \dots, 0)$$

$$\|T(y^n)\| = n \rightarrow \infty \text{ as } n \rightarrow \infty$$

Topological Isomorphism

Definition 4. A topological isomorphism *between normed vector spaces* X and Y is a linear transformation $T \in L(X, Y)$ that is invertible (one-to-one, onto), continuous, and has a continuous inverse.

Two normed vector spaces X and Y are topologically isomorphic if there is a topological isomorphism $T : X \rightarrow Y$.

$$\|f(x) - f(y)\| \leq \|Df(x)\| \|x - y\|$$

The Space $B(X, Y)$

$$\Rightarrow \exists \beta > 0 \text{ s.t.} \\ \|T(x)\| \leq \beta \|x\| \quad \forall x$$

Suppose X and Y are normed vector spaces. We define

$$B(X, Y) = \{T \in L(X, Y) : T \text{ is bounded}\}$$

Define :

$$\|T\|_{B(X, Y)} = \sup \left\{ \frac{\|T(x)\|_Y}{\|x\|_X} : x \in X, x \neq 0 \right\}$$

$$= \sup \{ \|T(x)\|_Y : \|x\|_X = 1 \}$$

$$\Rightarrow \frac{\|T(x)\|}{\|x\|} \leq \beta \quad \forall x \neq 0$$

$$\Rightarrow \|T(x)\| \leq \|T\| \|x\| \quad \forall x \in X \text{ by definition}$$

We skip the proofs of the rest of these results – read dIF.

$$\frac{\|T(x)\|}{\|x\|} = \frac{1}{\|x\|} \|T(x)\|$$

$$= \left\| \frac{1}{\|x\|} T(x) \right\| = \|T\left(\frac{x}{\|x\|}\right)\|$$

$$= \sup \left\{ \left\| T \left(\frac{x}{\|x\|} \right) \right\| : x \in X, x \neq 0 \right\}$$

$$x \neq 0, \quad y = \frac{x}{\|x\|} = \frac{1}{\|x\|} x \Leftrightarrow \|y\| = 1, y \in X$$
$$= \sup \left\{ \|T(y)\| : y = \frac{1}{\|x\|} x, x \in X, x \neq 0 \right\}$$

$$\|y\| = \left\| \frac{1}{\|x\|} x \right\| = \frac{1}{\|x\|} \|x\| = 1$$

The Space $B(X, Y)$

Theorem 5 (Thm. 4.8). *Let X, Y be normed vector spaces. Then*

$$\left(B(X, Y), \|\cdot\|_{B(X, Y)} \right)$$

is a normed vector space.

The Space $B(\mathbf{R}^n, \mathbf{R}^m)$

Theorem 6 (Thm. 4.9). *Let $T \in L(\mathbf{R}^n, \mathbf{R}^m)$ ($= B(\mathbf{R}^n, \mathbf{R}^m)$) with matrix $A = (a_{ij})$ with respect to the standard bases. Let*

$$M = \max\{|a_{ij}| : 1 \leq i \leq m, 1 \leq j \leq n\}$$

Then

$$M \leq \|T\| \leq M\sqrt{mn}$$

.

Compositions

Theorem 7 (Thm. 4.10). *Let $R \in L(\mathbf{R}^m, \mathbf{R}^n)$ and $S \in L(\mathbf{R}^n, \mathbf{R}^p)$.
Then*

$$\|S \circ R\| \leq \|S\| \|R\|$$

$$\ker T = \{0\} \Leftrightarrow T \text{ is invertible}$$

Invertibility

Define $\Omega(\mathbb{R}^n) = \{T \in L(\mathbb{R}^n, \mathbb{R}^n) : T \text{ is invertible}\}$

Theorem 8 (Thm. 4.11'). Suppose $T \in L(\mathbb{R}^n, \mathbb{R}^n)$ and E is the standard basis of \mathbb{R}^n . Then

T is invertible

$$\Leftrightarrow \ker T = \{0\}$$

$$\Leftrightarrow \det(Mtx_E(T)) \neq 0$$

$$\Leftrightarrow \det(Mtx_{V,V}(T)) \neq 0 \text{ for every basis } V$$

$$\Leftrightarrow \det(Mtx_{V,W}(T)) \neq 0 \text{ for every pair of bases } V, W$$

} nice exercise

Invertibility

Theorem 9 (Thm. 4.12). *If $S, T \in \Omega(\mathbf{R}^n)$, then $S \circ T \in \Omega(\mathbf{R}^n)$ and*

$$(S \circ T)^{-1} = T^{-1} \circ S^{-1}$$

Invertibility

Theorem 10 (Thm. 4.14). *Let $S, T \in L(\mathbf{R}^n, \mathbf{R}^n)$. If T is invertible and*

$$\|T - S\| < \frac{1}{\|T^{-1}\|}$$

then S is invertible. In particular, $\Omega(\mathbf{R}^n)$ is open in $L(\mathbf{R}^n, \mathbf{R}^n) = B(\mathbf{R}^n, \mathbf{R}^n)$.

Theorem 11 (Thm. 4.15). *The function $(\cdot)^{-1} : \Omega(\mathbf{R}^n) \rightarrow \Omega(\mathbf{R}^n)$ that assigns T^{-1} to each $T \in \Omega(\mathbf{R}^n)$ is continuous.*

$\{x_n\}$ Cauchy

$$\sup \{d(x_k, x_n) : k \geq n\}$$

$$d(x_m, x_n) = \sum_t \beta^{-t} \rho(x_{m+t}, x_{n+t}) < \varepsilon$$

$$\Rightarrow \beta^{-t} \rho(x_{m+t}, x_{n+t}) < \varepsilon \quad \forall t$$

$\{x_{n+t}\} \subseteq \mathbb{R}$ is Cauchy $\forall t$

$$x \in [0, 1]^n$$

$$d(x, y) = \sum_t \beta^{-t} p(x_t, y_t)$$

$$x_t \in [0, 1]$$

$$\forall \epsilon > 0$$

$$\exists T$$

s.t.

$$\sum_{t=T}^{\infty} \beta^{-t}$$

$$< \epsilon$$