

Announcements  
• PS 3 due tomorrow

# Econ 204 2020

## Lecture 9

### Outline

1. Quotient Vector Spaces
2. Matrix Representations of Linear Transformations
3. Change of Basis and Similarity
4. Eigenvalues and Eigenvectors
5. Diagonalization

$$x \sim y \Leftrightarrow x - y = \bar{w} \in W$$

$$y = x - \bar{w} \quad w = -\bar{w} \in W$$

## Quotient Vector Spaces

Given a vector space  $X$  and a vector subspace  $W$  of  $X$ , define an equivalence relation by

$$x \sim y \iff x - y \in W$$

Form a new vector space  $X/W$ : the set of vectors is

$$\{[x] : x \in X\}$$

where  $[x]$  denotes the equivalence class of  $x$  with respect to  $\sim$ .

$X/W$  is read “ $X$  mod  $W$ ”.

Note that the vectors in  $X/W$  are **sets** of vectors in  $X$ : for  $x \in X$ ,

$$[x] = \{x + w : w \in W\}$$

# Quotient Vector Spaces

We claim that  $X/W$  can be viewed as a vector space over  $F$ . Define the vector space operations  $+$ ,  $\cdot$  in  $X/W$  as follows:

Define

$$\begin{aligned}[x] + [y] &= [x + y] \\ \alpha[x] &= [\alpha x]\end{aligned}$$

**Exercise:** Verify that  $\sim$  is an equivalence relation and that vector addition and scalar multiplication are well-defined.\*

Then  $X/W$  is a vector space over  $F$  with these definitions for  $+$  and  $\cdot$ . (more exercise)

\* Show:  $\forall x, x', y, y' \in X, \forall \alpha \in F:$

$$[x] = [x'], [y] = [y'] \Rightarrow [x + y] = [x' + y'], [\alpha x] = [\alpha x']$$

$$X/W \neq X \setminus W$$

## Quotient Vector Spaces

(0, 0, 1)


**Example:** Let  $X = \mathbf{R}^3$  and let  $W = \{x \in \mathbf{R}^3 : x_1 = x_2 = 0\}$ .  
Then for  $x, y \in \mathbf{R}^3$ ,

$$\begin{aligned}x \sim y &\iff x - y \in W \\ &\iff x_1 - y_1 = 0, x_2 - y_2 = 0 \\ &\iff x_1 = y_1, x_2 = y_2\end{aligned}$$

and

$$[x] = \{x + w : w \in W\} = \{(x_1, x_2, z) : z \in \mathbf{R}\}$$

So the equivalence class corresponding to  $x$  is the line in  $\mathbf{R}^3$  through  $x$  parallel to the axis of the third coordinate.



## Example, cont.

What is  $X/W$ ? Intuitively this equivalence relation ignores the third coordinate, and we can identify the equivalence class  $[x]$  with the vector  $(x_1, x_2) \in \mathbf{R}^2$ .

The next two results show how to formalize this connection.

# Quotient Vector Spaces

**Theorem 1.** *If  $X$  is a vector space with  $\dim X = n$  for some  $n \in \mathbf{N}$  and  $W$  is a vector subspace of  $X$ , then*

$$\dim(X/W) = \dim X - \dim W$$

*Proof.* (Sketch) Begin with a basis  $\{w_1, \dots, w_c\}$  for  $W$ , and a basis  $\{[x_1], \dots, [x_k]\}$  for  $X/W$ . Show that

$$\{w_1, \dots, w_c\} \cup \{x_1, \dots, x_k\}$$

is a basis for  $X$ .



## Quotient Vector Spaces

**Theorem 2.** *Let  $X$  and  $Y$  be vector spaces over the same field  $F$  and  $T \in L(X, Y)$ . Then  $\text{Im } T$  is isomorphic to  $X/\ker T$ .*

*Proof.* Notice that if  $X$  is finite-dimensional, then

$$\begin{aligned}\dim(X/\ker T) &= \dim X - \dim \ker T && \text{(by the previous theorem)} \\ &= \text{Rank } T && \text{(by the Rank-Nullity Theorem)} \\ &= \dim \text{Im } T\end{aligned}$$

so  $X/\ker T$  is isomorphic to  $\text{Im } T$ . (why??)

We prove that this is true in general, and that the isomorphism is natural.



Define  $\tilde{T} : X / \ker T \rightarrow \text{Im } T$  by

$$\tilde{T}([x]) = T(x) \quad (\text{natural})$$

We first need to check that this is well-defined, that is, that if  $[x] = [x']$  then  $\tilde{T}([x]) = \tilde{T}([x'])$ .

$$\begin{aligned} [x] = [x'] &\Rightarrow x \sim x' \\ &\Rightarrow x - x' \in \ker T \\ &\Rightarrow T(x - x') = 0 = T(x) - T(x') \quad (T \text{ linear}) \\ &\Rightarrow T(x) = T(x') \end{aligned}$$

so  $\tilde{T}$  is well-defined.

Clearly,  $\tilde{T} : X / \ker T \rightarrow \text{Im } T$ . It is easy to check that  $\tilde{T}$  is linear,

so  $\tilde{T} \in L(X/\ker T, \text{Im } T)$ . Next we show that  $\tilde{T}$  is an isomorphism.

$$\begin{aligned} \underline{1-1:} \quad \tilde{T}([x]) = \tilde{T}([y]) &\Rightarrow T(x) = T(y) \\ &\Rightarrow T(x - y) = 0 && (T \text{ linear}) \\ &\Rightarrow x - y \in \ker T \\ &\Rightarrow x \sim y \\ &\Rightarrow [x] = [y] \end{aligned}$$

so  $\tilde{T}$  is one-to-one.

$$\begin{aligned} \underline{\text{onto:}} \quad y \in \text{Im } T &\Rightarrow \exists x \in X \text{ s.t. } T(x) = y \\ &\Rightarrow \tilde{T}([x]) = y = T(x) \end{aligned}$$

so  $\tilde{T}$  is onto, hence  $\tilde{T}$  is an isomorphism. □

$$T(x) = (x_1 + x_3, x_2 + x_3)$$

$$T(x) = (0, 0) \Leftrightarrow \begin{cases} x_1 + x_3 = 0 \\ x_2 + x_3 = 0 \end{cases}$$

**Example:** Consider  $T \in L(\mathbf{R}^3, \mathbf{R}^2)$  defined by

$$T(x_1, x_2, x_3) = (x_1, x_2) \Leftrightarrow \begin{cases} x_1 = x_2 = -x_3 \end{cases}$$

Then

$$\ker T = \{x \in \mathbf{R}^3 : x_1 = x_2 = 0\} \quad \left( = \omega \text{ from previous example} \right)$$

is the  $x_3$ -axis.

Given  $x$ , the equivalence class  $[x]$  is just the line through  $x$  parallel to the  $x_3$ -axis.

$$\tilde{T}([x]) = T(x_1, x_2, x_3) = (x_1, x_2)$$

and

$$\text{Im } T = \mathbf{R}^2, \quad X / \ker T \cong \mathbf{R}^2 = \text{Im } T$$

as we suggested intuitively above (here the symbol  $\cong$  denotes isomorphism, that is, we write  $Y \cong Z$  if  $Y$  and  $Z$  are isomorphic.)

# Coordinate Representations

(over  $\mathbb{R}$ )

Every real vector space  $X_n$  with dimension  $n$  is isomorphic to  $\mathbf{R}^n$ .  
What's the isomorphism?

Let  $X$  be a finite-dimensional vector space over  $\mathbf{R}$  with  $\dim X = n$ . Fix any Hamel basis  $V = \{v_1, \dots, v_n\}$  of  $X$ . Any  $x \in X$  has a unique representation

$$x = \sum_{j=1}^n \beta_j v_j$$

$$\begin{aligned} \text{crd}_V &: X \rightarrow \mathbb{R}^n \\ \text{crd}_W &: Y \rightarrow \mathbb{R}^m \end{aligned}$$

(here, we allow  $\beta_j = 0$ ).

Define:

$$\text{crd}_V(x) = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_n \end{pmatrix} \in \mathbf{R}^n$$

"coordinate representation of  $x$  with respect to  $V$ " 10

$V = \{v_1, \dots, v_n\}$  basis for  $X$

$crd_V(x)$  is the vector of coordinates of  $x$  with respect to the basis  $V$ .

Notice:

$$crd_V(v_1) = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad crd_V(v_2) = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad \dots \quad crd_V(v_n) = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

$crd_V$  is an isomorphism from  $X$  to  $\mathbf{R}^n$

# Matrix Representations of Linear Transformations

Suppose  $T \in L(X, Y)$ ,  $\dim X = n$ ,  $\dim Y = m$ . Fix bases

$$V = \{v_1, \dots, v_n\} \text{ of } X$$

$$W = \{w_1, \dots, w_m\} \text{ of } Y$$

$T(v_j) \in Y$ , so

$$T(v_j) = \sum_{i=1}^m \alpha_{ij} w_i$$

for some  $\alpha_{ij}$ ,  $i=1, \dots, m$   
(again allow  $\alpha_{ij}=0$ )

Define

$$Mtx_{W,V}(T) = \begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & \cdots & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix} \quad m \times n \text{ matrix}$$

"matrix representation  
of  $T$  with respect  
to  $V$  and  $W$ "

↑ coordinates of  $T(v_1)$  w.r.t.  $W$       ↑ coordinates of  $T(v_n)$  w.r.t.  $W$

# Matrix Representations of Linear Transformations

Notice that the columns are the coordinates (expressed with respect to  $W$ ) of  $T(v_1), \dots, T(v_n)$ .

Observe

$$\begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & \cdots & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \begin{pmatrix} \alpha_{11} \\ \vdots \\ \alpha_{m1} \end{pmatrix}$$

so

$$\begin{aligned} Mtx_{W,V}(T) \cdot crd_V(v_j) &= crd_W(T(v_j)) \quad \forall v_j \in V \\ \Rightarrow Mtx_{W,V}(T) \cdot crd_V(x) &= crd_W(T(x)) \quad \forall x \in X \end{aligned}$$

# Matrix Representations

Multiplying a vector by a matrix does two things:

- Computes the action of  $T$
- Accounts for the change in basis



basis for  $X$   
↑

basis for  $Y$   
↑

**Example:**  $X = Y = \mathbf{R}^2$ ,  $V = \{(1, 0), (0, 1)\}$ ,  $W = \{(1, 1), (-1, 1)\}$ ,  
 $T = id$ , that is,  $T(x) = x$  for each  $x$ .

$$Mtx_{W,V}(T) \neq \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$Mtx_{W,V}(T)$  is the matrix that *changes basis* from  $V$  to  $W$ .

$$V = \{ (1,0), (0,1) \}$$

$$W = \{ w_1, w_2 \} = \{ (1,1), (-1,1) \}$$

$$\text{ord}_V(w_1) = (1,1)$$

$$\text{ord}_W(w_1) = (1,0)$$

How do we compute it?

Look for coord. representations of  $T(v_1) = v_1$  and  $T(v_2) = v_2$ :

$$T(v_1) = v_1 = (1,0) = \alpha_{11}(1,1) + \alpha_{21}(-1,1)$$

$$\alpha_{11} - \alpha_{21} = 1$$

$$\alpha_{11} + \alpha_{21} = 0$$

$$2\alpha_{11} = 1, \alpha_{11} = \frac{1}{2}$$

$$\alpha_{21} = -\frac{1}{2}$$

$$\Rightarrow \text{ord}_W(v_1) = \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}$$

$$T(v_2) = v_2 = (0,1) = \alpha_{12}(1,1) + \alpha_{22}(-1,1)$$

$$\alpha_{12} - \alpha_{22} = 0$$

$$\alpha_{12} + \alpha_{22} = 1$$

$$2\alpha_{12} = 1, \alpha_{12} = \frac{1}{2}$$

$$\alpha_{22} = \frac{1}{2}$$

$$\Rightarrow \text{ord}_W(v_2) = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$$

$$W = \{w_1, w_2\}$$

$$\text{ord}_W(w_i) ?$$

$$T: X \rightarrow Y, T = \text{id}$$

$$y = \alpha_1 w_1 + \alpha_2 w_2$$

$$\text{ord}_W(y) = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}$$

$$w_1 = 1 \cdot w_1 + 0 \cdot w_2$$

So

$$Mtx_{W,V}(\text{id}) = \begin{pmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{pmatrix}$$

$$\begin{array}{cc} \uparrow & \uparrow \\ \text{ord}_W(T(v_1)) & \text{ord}_W(T(v_2)) \\ \text{"} & \text{"} \\ \text{ord}_W(v_1) & \text{ord}_W(v_2) \end{array}$$

## Matrix Representations

**Theorem 3** (Thm. 3.5'). *Let  $X$  and  $Y$  be vector spaces over the same field  $F$ , with  $\dim X = n$ ,  $\dim Y = m$ . Then  $L(X, Y)$ , the space of linear transformations from  $X$  to  $Y$ , is isomorphic to  $F_{m \times n}$ , the vector space of  $m \times n$  matrices over  $F$ . If  $V = \{v_1, \dots, v_n\}$  is a basis for  $X$  and  $W = \{w_1, \dots, w_m\}$  is a basis for  $Y$ , then*

$$Mtx_{W,V} \in L(L(X, Y), F_{m \times n})$$

*and  $Mtx_{W,V}$  is an isomorphism from  $L(X, Y)$  to  $F_{m \times n}$ .*

# Matrix Representations

**Theorem 4** (From Handout). *Let  $X, Y, Z$  be finite-dimensional vector spaces over the same field  $F$  with bases  $U, V, W$  respectively. Let  $S \in L(X, Y)$  and  $T \in L(Y, Z)$ . Then*

$$\text{Mtx}_{W,V}(T) \cdot \text{Mtx}_{V,U}(S) = \text{Mtx}_{W,U}(T \circ S)$$

*i.e. matrix multiplication corresponds via the matrix representation isomorphism to composition of linear transformations.*

*Proof.* See handout. □

Note that  $\text{Mtx}_{W,V}$  is a function from  $L(X, Y)$  to the space  $F_{m \times n}$  of  $m \times n$  matrices, while  $\text{Mtx}_{W,V}(T)$  is an  $m \times n$  matrix.

# Matrix Representations

The theorem can be summarized by the following “Commutative Diagram:”


$$\begin{array}{ccccc}
 & & S & & T \\
 & X & \rightarrow & Y & \rightarrow & Z \\
 \text{crd}_U & \updownarrow & & \updownarrow \text{crd}_V & & \updownarrow \text{crd}_W \\
 & \mathbf{R}^n & \rightarrow & \mathbf{R}^m & \rightarrow & \mathbf{R}^r \\
 & & \text{Mtx}_{V,U}(S) & & \text{Mtx}_{W,V}(T) & & 
 \end{array}$$

We say the diagram commutes because you get the same answer any way you go around the diagram (in directions allowed by the arrows). The *crd* arrows go in both directions because *crd* is an isomorphism.

# Change of Basis

Let  $X$  be a finite-dimensional vector space with basis  $V$ . If  $T \in L(X, X)$  it is customary to use the same basis in the domain and range. In this case,  $Mtx_V(T)$  denotes  $Mtx_{V,V}(T)$ .

**Question:** If  $W$  is another basis for  $X$ , how are  $Mtx_V(T)$  and  $Mtx_W(T)$  related?



$$\begin{aligned}
Mtx_{V,W}(id) \cdot Mtx_W(T) \cdot Mtx_{W,V}(id) &= Mtx_{V,W}(id) \cdot Mtx_{W,V}(T \circ id) \\
&= Mtx_{V,V}(id \circ T \circ id) \\
&= Mtx_V(T)
\end{aligned}$$

and

$$\begin{aligned}
Mtx_{V,W}(id) \cdot Mtx_{W,V}(id) &= Mtx_{V,V}(id) \\
&= \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}
\end{aligned}$$



$$V = \{v_1, \dots, v_n\} \quad U = \{u_1, \dots, u_m\}$$

$$T^{ij}(v_i) = u_j$$

$$T^{ij}(v_k) = 0 \quad \forall k \neq i$$

So this says that

$$\text{Mtx}_V(T) = P^{-1} \text{Mtx}_W(T) P$$

for the invertible matrix

$$P = \text{Mtx}_{W,V}(id)$$

that is the change of basis matrix.

On the other hand, if  $P$  is any invertible matrix, then  $P$  is also a change of basis matrix for appropriate corresponding bases (see handout).

# Similarity

**Definition 1.** <sup>( $n \times n$ )</sup> Square matrices  $A$  and  $B$  are similar if

$$A = P^{-1}BP$$

for some invertible matrix  $P$ .

# Similarity

**Theorem 5.** *Suppose that  $X$  is a finite-dimensional vector space.*

- 1. If  $T \in L(X, X)$  then any two matrix representations of  $T$  are similar. That is, if  $U, W$  are any two bases of  $X$ , then  $Mtx_W(T)$  and  $Mtx_U(T)$  are similar.*
- 2. Conversely, two similar matrices represent the same linear transformation  $T$ , relative to suitable bases. That is, given similar matrices  $A, B$  with  $A = P^{-1}BP$  and any basis  $U$ , there is a basis  $W$  and  $T \in L(X, X)$  such that*

$$\begin{aligned} B &= Mtx_U(T) \\ A &= Mtx_W(T) \\ P &= Mtx_{U,W}(id) \\ P^{-1} &= Mtx_{W,U}(id) \end{aligned}$$

*Proof.* See Handout on Diagonalization and Quadratic Forms.



# Eigenvalues and Eigenvectors

Here, we define eigenvalues and eigenvectors of a linear transformation and show that  $\lambda$  is an eigenvalue of  $T$  if and only if  $\lambda$  is an eigenvalue for some matrix representation of  $T$  if and only if  $\lambda$  is an eigenvalue for every matrix representation of  $T$ .

**Definition 2.** *Let  $X$  be a vector space and  $T \in L(X, X)$ . We say that  $\lambda$  is an eigenvalue of  $T$  and  $v \neq 0$  is an eigenvector corresponding to  $\lambda$  if  $T(v) = \lambda v$ .*

# Eigenvalues and Eigenvectors

**Theorem 6** (Theorem 4 in Handout). *Let  $X$  be a finite-dimensional vector space, and  $U$  a basis. Then  $\lambda$  is an eigenvalue of  $T$  if and only if  $\lambda$  is an eigenvalue of  $Mtx_U(T)$ .  $v$  is an eigenvector of  $T$  corresponding to  $\lambda$  if and only if  $crd_U(v)$  is an eigenvector of  $Mtx_U(T)$  corresponding to  $\lambda$ .*

*Proof.* By the Commutative Diagram Theorem,

$$\begin{aligned} T(v) = \lambda v &\Leftrightarrow crd_U(T(v)) = crd_U(\lambda v) \\ &\Leftrightarrow Mtx_U(T)(crd_U(v)) = \lambda(crd_U(v)) \end{aligned}$$

$$A = Mtx_U(T), \quad x = crd_U(v),$$

$$\Leftrightarrow Ax = \lambda x$$

□

# Computing Eigenvalues and Eigenvectors

Suppose  $\dim X = n$ ; let  $I$  be the  $n \times n$  identity matrix. Given  $T \in L(X, X)$ , fix a basis  $U$  and let

$$A = Mtx_U(T)$$

Find the eigenvalues of  $T$  by computing the eigenvalues of  $A$ :

$$\begin{aligned} \exists v \neq 0 \text{ s.t. } Av = \lambda v &\iff (A - \lambda I)v = 0 && \text{for some } v \neq 0 \\ &\iff (A - \lambda I) \text{ is not invertible} \\ &\iff \det(A - \lambda I) = 0 \end{aligned}$$

We have the following facts:

- If  $A \in \mathbf{R}_{n \times n}$ ,

$$f(\lambda) = \det(A - \lambda I)$$

is an  $n^{\text{th}}$  degree polynomial in  $\lambda$  with real coefficients; it is called the *characteristic polynomial* of  $A$ .

- $f$  has  $n$  roots in  $\mathbf{C}$ , counting multiplicity:

$$f(\lambda) = (\lambda - c_1)(\lambda - c_2) \cdots (\lambda - c_n)$$

( may have  
 $c_i = c_j, i \neq j$  )

where  $c_1, \dots, c_n \in \mathbf{C}$  are the eigenvalues; the  $c_j$ 's are not necessarily distinct. Notice that  $f(\lambda) = 0$  if and only if  $\lambda \in \{c_1, \dots, c_n\}$ , so the roots are the solutions of the equation  $f(\lambda) = 0$ .



- the roots that are not real come in conjugate pairs:

$$f(a + bi) = 0 \Leftrightarrow f(a - bi) = 0$$

- if  $\lambda = c_j \in \mathbf{R}$ , there is a corresponding eigenvector in  $\mathbf{R}^n$ .
- if  $\lambda = c_j \notin \mathbf{R}$ , the corresponding eigenvectors are in  $\mathbf{C}^n \setminus \mathbf{R}^n$ .

# Diagonalization

**Definition 3.** *Suppose  $X$  is a finite-dimensional vector space with basis  $U$ . Given a linear transformation  $T \in L(X, X)$ , let*

$$A = Mtx_U(T)$$

*We say that  $A$  can be diagonalized if there is a basis  $W$  for  $X$  such that  $Mtx_W(T)$  is a diagonal matrix, that is,*

$$Mtx_W(T) = \begin{pmatrix} \lambda_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & \lambda_n \end{pmatrix}$$

So

$A$  can be diagonalized  $\iff A$  is similar to a diagonal matrix  
 $\iff A = P^{-1}BP$  where  $B$  is diagonal

$$\begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$$

Suppose there is a basis  $W$  such that

$$Mtx_W(T) = \begin{pmatrix} \lambda_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & \lambda_n \end{pmatrix}$$

$\Rightarrow \lambda_1, \dots, \lambda_n$  are eigenvalues of  $Mtx_W(T)$  and  $T$

Then the standard basis vectors of  $\mathbf{R}^n$  are eigenvectors of  $Mtx_W(T)$ .

In general:

$z_j$  is an eigenvector of  $T$  corresponding to  $\lambda_j \iff crd_W(z_j)$  is an eigenvector of  $Mtx_W(T)$  corresponding to  $\lambda_j$ .

So an eigenvector <sup>of  $T$</sup>  corresponding to  $\lambda_j$  is  $w_j$ , since  $crd_W(w_j) = e_j$ , the  $j^{th}$  standard basis vector in  $\mathbf{R}^n$ .

$$\begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

Thus  $Mtx_W(T)$  is diagonal if and only if  $W = \{w_1, \dots, w_n\}$  where  $w_j$  is an eigenvector of  $T$  corresponding to  $\lambda_j$  for each  $j$ .

Then the action of  $T$  is clear: it stretches each basis element  $w_i$  by the factor  $\lambda_i$ .

$$\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$$

## Diagonalization

**Theorem 7** (Thm. 6.7'). *Let  $X$  be an  $n$ -dimensional vector space,  $T \in L(X, X)$ ,  $U$  any basis of  $X$ , and  $A = Mtx_U(T)$ . Then the following are equivalent:*

1.  *$A$  can be diagonalized*
2. *there is a basis  $W$  for  $X$  consisting of eigenvectors of  $T$*
3. *there is a basis  $V$  for  $\mathbf{R}^n$  consisting of eigenvectors of  $A$*

*Proof.* Follows from Theorem 6.7 in de la Fuente and Theorem 4 from the Handout.  $\square$

# Diagonalization

**Theorem 8** (Thm. 6.8'). *Let  $X$  be a vector space and  $T \in L(X, X)$ .*

- 1. If  $\lambda_1, \dots, \lambda_m$  are distinct eigenvalues of  $T$  with corresponding eigenvectors  $v_1, \dots, v_m$ , then  $\{v_1, \dots, v_m\}$  is linearly independent.*
- 2. If  $\dim X = n$  and  $T$  has  $n$  distinct eigenvalues, then  $X$  has a basis consisting of eigenvectors of  $T$ ; consequently, if  $U$  is any basis of  $X$ , then  $Mtx_U(T)$  is diagonalizable.*

*Proof.* This is an adaptation of the proof of Theorem 6.8 in de la Fuente. □

$$|J(f) - J(g)| = \left| \int_0^1 e^{-x} \left( \frac{f(x)}{1+f'(x)^2} - \frac{g(x)}{1+g'(x)^2} \right) dx \right|$$

$$\leq \int_0^1 e^{-x} \left| \frac{f(x)(1+g'(x)^2) - g(x)(1+f'(x)^2)}{(1+f'(x)^2)(1+g'(x)^2)} \right| dx$$

$$\leq \int_0^1 e^{-x} |f(x)(1+g'(x)^2) - g(x)(1+f'(x)^2)| dx$$

$$\leq \int_0^1 e^{-x} | (f(x) - g(x)) + (f(x)g'(x)^2 - g(x)f'(x)^2) | dx$$

$$\leq \int_0^1 e^{-x} |f(x) - g(x)| dx + \int_0^1 e^{-x} |f(x)g'(x)^2 - g(x)f'(x)^2| dx$$

$$\leq \|f - g\|_\infty + \underbrace{\hspace{10em}}$$



$$\int_0^1 e^{-x} |f(x)g'(x)^2 - g(x)f'(x)^2| dx$$

$$= \int_0^1 e^{-x} |f(x)g'(x)^2 - f(x)f'(x)^2 + f(x)f'(x)^2 - g(x)f'(x)^2| dx$$

$$\leq \int_0^1 e^{-x} |f(x)g'(x)^2 - f(x)f'(x)^2| dx$$

$$+ \int_0^1 e^{-x} |f(x)f'(x)^2 - g(x)f'(x)^2| dx$$

$$\leq \int_0^1 e^{-x} |f(x)| |g'(x) - f'(x)| |g'(x) + f'(x)| dx$$

$$+ \int_0^1 e^{-x} |f'(x)|^2 |f(x) - g(x)| dx$$



$$\leq \|f'\|_\infty^2 \|f - g\|_\infty$$

$$\int_0^1 e^{-x} |f(x)| |g'(x) - f'(x)| |g'(x) + f'(x)| dx$$


$$\leq \|f\|_\infty \int_0^1 e^{-x} |g'(x) - f'(x)| |g'(x) + f'(x)| dx$$

$$\leq \|f\|_\infty \|g' - f'\|_\infty \|g' + f'\|_\infty \left( \int_0^1 e^{-x} dx \right)$$


$$\|f - g\| = \|f - g\|_\infty + \|f' - g'\|_\infty$$

$$\int_0^1 h(x) dx \leq \int_0^1 |h(x)| dx$$

$\{x: f(x) > \alpha\}$  open  $\rightarrow \alpha$



$$\liminf_{x_n \rightarrow x} f(x_n) \geq f(x)$$



$$\ker T = \{x \in X : T(x) = 0\}$$

$$K(y) = \{x \in X : T(x) = \alpha y \text{ for some } \alpha \in \mathbb{K}\}$$

$$T(x) = y \Rightarrow T(\alpha x) = \alpha y$$

$$T(x_1) = T(x_2) = y \quad T(x_1 + x_2) = 2y$$