

## Announcements

- look for bCourses email about exam

# Econ 204 2022

## Lecture 12

### Outline

0. Characterizing critical pts  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  w/ Taylor's Thm
1. An Example
  2. Regular and Critical Points and Values
  3. Inverse Function Theorem
  4. Implicit Function Theorem
  5. Sard's Theorem
  6. Transversality Theorem

$f: \mathbb{R}^n \rightarrow \mathbb{R}$  differentiable

## Characterizing Critical Points

**Definition 4.** We say  $f$  has a saddle at  $x$  if  $Df(x) = 0$  but  $f$  has neither a local maximum nor a local minimum at  $x$ .

$f$  has a critical point at  $x$  if  $Df(x) = 0$ .

$D^2f(x)$  symmetric  $\forall x$

of eigenvectors  
of  $Df(x)$

## Characterizing Critical Points

**Corollary 1.** Suppose  $X \subseteq \mathbb{R}^n$  is open and  $x \in X$ . If  $f : X \rightarrow \mathbb{R}$  is  $C^2$ , there is an orthonormal basis  $\{v_1, \dots, v_n\}$  and corresponding eigenvalues  $\lambda_1, \dots, \lambda_n \in \mathbb{R}$  of  $D^2f(x)$  such that

$$\begin{aligned} f(x+h) &= f(x + \gamma_1 v_1 + \dots + \gamma_n v_n) \\ &= f(x) + \sum_{i=1}^n (Df(x)v_i) \gamma_i + \frac{1}{2} \sum_{i=1}^n \lambda_i \gamma_i^2 + o(|\gamma|^2) \end{aligned}$$

$$h = \sum \gamma_i v_i$$

where  $\gamma_i = h \cdot v_i$ .

$$= f(x) + Df(x)h + \frac{1}{2} h^T D^2f(x) h + o(\|h\|^2)$$

1. If  $f \in C^3$ , we may strengthen  $o(|\gamma|^2)$  to  $O(|\gamma|^3)$ .

2. If  $f$  has a local maximum or local minimum at  $x$ , then

$$Df(x) = 0$$

3. If  $Df(x) = 0$ , then

- $\lambda_1, \dots, \lambda_n > 0 \Rightarrow f$  has a local minimum at  $x$
- $\lambda_1, \dots, \lambda_n < 0 \Rightarrow f$  has a local maximum at  $x$
- $\lambda_i < 0$  for some  $i$ ,  $\lambda_j > 0$  for some  $j \Rightarrow f$  has a saddle at  $x$
- $\lambda_1, \dots, \lambda_n \geq 0$ ,  $\lambda_i > 0$  for some  $i \Rightarrow f$  has a local minimum or a saddle at  $x$
- $\lambda_1, \dots, \lambda_n \leq 0$ ,  $\lambda_i < 0$  for some  $i \Rightarrow f$  has a local maximum or a saddle at  $x$
- $\lambda_1 = \dots = \lambda_n = 0$  gives no information.

*Proof.* (Sketch) From our study of quadratic forms, we know the behavior of the quadratic terms is determined by the signs of the eigenvalues. If  $\lambda_i = 0$  for some  $i$ , then we know that the quadratic form arising from the second partial derivatives is identically zero in the direction  $v_i$ , and the higher derivatives will determine the behavior of the function  $f$  in the direction  $v_i$ . For example, if  $f(x) = x^3$ , then  $f'(0) = 0$ ,  $f''(0) = 0$ , but we know that  $f$  has a saddle at  $x = 0$ ; however, if  $f(x) = x^4$ , then again  $f'(0) = 0$  and  $f''(0) = 0$  but  $f$  has a local (and global) minimum at  $x = 0$ . □

# Comparative Statics

In many problems we are interested in how endogenously determined variables are affected by exogenously given parameters. Here we study problems in which the variables of interest are characterized as solutions to a parameterized family of equations.

To formalize, let  $X \subseteq \mathbf{R}^n$  and  $A \subseteq \mathbf{R}^p$  be open, and let  $f : X \times A \rightarrow \mathbf{R}^m$ . For a given  $a \in A$ , consider solutions  $x \in X$  to the family of equations

$$f(x, a) = 0$$

*m equations*  
*n unknowns*  
*p parameters*

We want to characterize the set of solutions and study how this set depends on the parameter  $a$ .

## An Example

Consider the function  $f : (0, 2\pi) \times \mathbf{R} \rightarrow \mathbf{R}$  defined by

$$f(x, a) = \sin x + a$$

Let  $X = (0, 2\pi)$ . For fixed  $a$ , let

$$f_a(x) = f(x, a) = \sin x + a$$

$$f_a : (0, 2\pi) \rightarrow \mathbb{R}$$

We look for solutions  $x \in (0, 2\pi)$  to the equation

$$f_a(x) = f(x, a) = \sin x + a = 0$$

that is, the  $x \in (0, 2\pi)$  such that

$$\sin x = -a$$

Let  $\Psi : A \rightarrow 2^X$  denote the solution correspondence, so

$$\Psi(a) = \{x \in (0, 2\pi) : f_a(x) = \sin x + a = 0\}$$

## An Example

Start with  $a = 0$ . For  $x \in (0, 2\pi)$ ,

$$f_0(x) = \sin x = 0 \iff x = \pi$$

so  $\Psi(0) = \{\pi\}$ .

Notice that for  $x$  near  $\pi$ , for example in the neighborhood  $(\pi/2, 3\pi/2)$ , and for  $a$  near 0,  $\sin^{-1}(a)$  remains single-valued and depends smoothly on  $a$ .


In addition, we can predict the direction of change:  $x$  is increasing in  $a$ .



$$f_0(x) = \sin x + 0 = 0$$

$$f_a(x) = \sin x + a = 0$$

$$\Leftrightarrow \sin x = -a$$



## An Example

Now consider  $a = 1$ . For  $x \in (0, 2\pi)$ ,


$$\begin{aligned} f_1(x) &= \sin x + 1 = 0 \\ \iff \sin x &= -1 \\ \iff x &= \frac{3\pi}{2} \end{aligned}$$

So  $\Psi(1) = \{3\pi/2\}$ .

$$\sin x + a' = 0 \iff \sin x = -a'$$

But note that for  $a' > 1$ ,  $\Psi(a') = \emptyset$ , while for  $a < 1$  close to 1, there are two solutions near  $3\pi/2$ , one above and one below  $3\pi/2$ .

$\Psi$  is not lower hemicontinuous at  $a = 1$ .



# Regular and Critical Points and Values

Suppose  $X \subseteq \mathbf{R}^n$  is open. Suppose  $f : X \rightarrow \mathbf{R}^m$  is differentiable at  $x \in X$ , and let  $W = \{e_1, \dots, e_n\}$  denote the standard basis of  $\mathbf{R}^n$ . Then  $df_x \in L(\mathbf{R}^n, \mathbf{R}^m)$ , and

$$\begin{aligned} \text{Rank } df_x &= \dim \text{Im}(df_x) \\ &= \dim \text{span} \{df_x(e_1), \dots, df_x(e_n)\} \\ &= \dim \text{span} \{Df(x)e_1, \dots, Df(x)e_n\} \\ &= \dim \text{span} \{\text{column 1 of } Df(x), \dots, \text{column } n \text{ of } Df(x)\} \\ &= \text{Rank } Df(x) \end{aligned}$$

(linearity +  $\{e_1, \dots, e_n\}$   
basis for  $\mathbf{R}^n$ )

Thus,

$$\text{Rank } df_x \leq \min\{m, n\}$$

We say  $df_x$  has *full rank* if  $\text{Rank } df_x = \min\{m, n\}$ , that is, if  $df_x$  has maximum possible rank.

$$\begin{aligned} &\Leftrightarrow \text{Rank } Df(x) = \min\{m, n\} \\ &\Leftrightarrow Df(x) \text{ has full rank} \end{aligned}$$

# Regular and Critical Points and Values

**Definition 1.** Suppose  $X \subseteq \mathbb{R}^n$  is open. Suppose  $f : X \rightarrow \mathbb{R}^m$  is differentiable on  $X$ .

- $x$  is a regular point of  $f$  if  $\text{Rank } df_x = \min\{m, n\}$ .

$df_x$  has full rank ( $\Leftrightarrow Df(x)$  has full rank)

- $x$  is a critical point of  $f$  if  $\text{Rank } df_x < \min\{m, n\}$ .

$\Leftrightarrow \text{Rank } Df(x) < \min\{m, n\}$

- $y$  is a critical value of  $f$  if there exists  $x \in f^{-1}(y)$  such that  $x$  is a critical point of  $f$ .

- $y$  is a regular value of  $f$  if  $y$  is not a critical value of  $f$

$y$  regular value of  $f \Leftrightarrow \forall x \in f^{-1}(y), df_x$  has full rank  
 $\downarrow$   
 $Df(x)$

**Example:** Consider the function  $g : (0, 2\pi) \rightarrow \mathbf{R}$  defined by

$$g(x)(= f_0(x)) = \sin x$$

Note that  $g'(x) = \cos x$ , so  $g'(x) = 0 \iff x = \pi/2$  or  $x = 3\pi/2$ .

$Dg(x)$  is the  $1 \times 1$  matrix  $(g'(x))$ , so

$$\text{Rank } dg_x = \text{Rank } Dg(x) = 1 \iff \underline{g'(x) \neq 0}$$

critical points of  $g$ :  $\pi/2$  and  $3\pi/2$

regular points of  $g$ :  $(0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \frac{3\pi}{2}) \cup (\frac{3\pi}{2}, 2\pi)$

critical values of  $g$ :  $g(\pi/2) = \sin(\pi/2) = 1$  and  $g(3\pi/2) = \sin(3\pi/2) = -1$

regular values of  $g$ :  $(-\infty, -1) \cup (-1, 1) \cup (1, \infty)$

$$\begin{pmatrix} \frac{\partial f^1}{\partial x_1}(x) & \frac{\partial f^1}{\partial x_2}(x) \\ \frac{\partial f^2}{\partial x_1}(x) & \frac{\partial f^2}{\partial x_2}(x) \end{pmatrix}$$

In particular, notice that 0 is not a critical value of  $g$ .

Given  $a \in \mathbf{R}$ , as above consider the perturbed function

$$f_a(x) = g(x) + a$$

Notice that  $f'_a(x) = g'(x)$ , so the critical points of  $f_a$  are the same as those of  $g$ ,  $\pi/2$  and  $3\pi/2$ .

For  $a$  close to zero, the solution to the equation

$$f_a(x) = 0$$

near  $x = \pi$  moves smoothly with respect to changes in  $a$ . The direction the solution moves is determined by the sign of  $f'_a$ .

Now let  $a = 1$ . Since  $3\pi/2$  is a critical point of  $f_1$ , 0 is a critical value of  $f_1$ .

# Inverse Function Theorem

**Theorem 1** (Thm. 4.6, Inverse Function Theorem). *Suppose  $X \subseteq \mathbf{R}^n$  is open,  $f : X \rightarrow \mathbf{R}^n$  is  $C^1$  on  $X$ , and  $x_0 \in X$ . If  $\det Df(x_0) \neq 0$  (i.e.  $x_0$  is a regular point of  $f$ ) then there are open neighborhoods  $U$  of  $x_0$  and  $V$  of  $f(x_0)$  such that*

$$\begin{aligned} f : U &\rightarrow V \text{ is one-to-one and onto} \\ f^{-1} : V &\rightarrow U \text{ is } C^1 \\ Df^{-1}(f(x_0)) &= [Df(x_0)]^{-1} \end{aligned}$$

*If in addition  $f \in C^k$ , then  $f^{-1} \in C^k$ .*

**Remark:**  $f$  is one-to-one only on  $U$ ; it need not be one-to-one globally. Thus  $f^{-1}$  is only a local inverse.



*Proof.* Read the proof in de la Fuente. This is pretty hard. The idea is that since  $\det Df(x_0) \neq 0$ , then  $df_{x_0} : \mathbf{R}^n \rightarrow \mathbf{R}^n$  is one-to-one and onto. You need to find a neighborhood  $U$  of  $x_0$  sufficiently small such that the Contraction Mapping Theorem implies that  $f$  is one-to-one and onto. \*

To see the formula for  $Df^{-1}$ , let  $\text{id}_U$  denote the identity function from  $U$  to  $U$  and  $I$  denote the  $n \times n$  identity matrix. Then

$$\begin{aligned} Df^{-1}(f(x_0))Df(x_0) &= D(f^{-1} \circ f)(x_0) \\ &= D(\text{id}_U)(x_0) \\ &= I \\ \Rightarrow Df^{-1}(f(x_0)) &= [Df(x_0)]^{-1} \end{aligned}$$

□

local inverse? Given  $y$ , want unique  $x$  s.t.  $y = f(x)$ :

\* Let  $A = Df(x_0)$

Set

$$g(x) = x + A^{-1}(y - f(x))$$

Note  $y = f(x) \Leftrightarrow x$  is a fixed point of  $g$

Then show  $g$  is a contraction mapping on a suitably small neighborhood  $U$  around  $x_0$ .

# Inverse Function Theorem


**Example:** Let  $g : (0, 2\pi) \rightarrow \mathbf{R}$  be given by  $g(x) = \sin x$ . Let  $x_0 = \pi$ .

Then  $g'(x_0) = \cos \pi = \underline{-1} \neq 0$ , so by the Inverse Function Theorem there exists an open set  $U \subseteq (0, 2\pi)$  with  $\pi \in U$ , an open set  $V \subseteq \mathbf{R}$  with  $0 = g(\pi) \in V$  and a  $C^1$  function  $h : V \rightarrow U$  such that  $g(h(v)) = v$  for all  $v \in V$ .

At  $x = 3\pi/2$ ,  $g'(x) = \cos(3\pi/2) = 0$ , and  $g$  has no local inverse function there: for every open neighborhood  $U$  of  $3\pi/2$  and every open neighborhood  $V$  of  $-1 = g(3\pi/2)$ , there exists  $v \in V$  and  $x_1 \neq x_2 \in U$  such that

$$g(x_1) = \sin x_1 = v = \sin x_2 = g(x_2)$$

$$0 = f(x, a) = g(x) + a$$



When does equation  $f(x, a) = 0$   
determine  $x$  as a function of  $a$  (locally)?

## Implicit Function Theorem

**Theorem 2** (Thm. 2.2, Implicit Function Theorem). Suppose  $X \subseteq \mathbb{R}^n$  and  $A \subseteq \mathbb{R}^p$  are open and  $f : X \times A \rightarrow \mathbb{R}^n$  is  $C^1$ . Suppose  $\underline{f(x_0, a_0) = 0}$  and  $\det(D_x f(x_0, a_0)) \neq 0$ , i.e.  $\underline{x_0}$  is a regular point of  $f(\cdot, a_0)$ . Then there are open neighborhoods  $U$  of  $x_0$  ( $U \subseteq X$ ) and  $W$  of  $a_0$  such that

$$\forall a \in W \quad \exists! x \in U \text{ s.t. } f(x, a) = 0$$

For each  $a \in W$  let  $g(a)$  be that unique  $x$ . Then  $g : W \rightarrow X$  is  $C^1$  and

$$Dg(a_0) = - \underline{[D_x f(x_0, a_0)]^{-1}} [D_a f(x_0, a_0)]$$

If in addition  $f \in C^k$ , then  $g \in C^k$ . ↖ ↗

Notice:  $g(a_0) = x_0$

*Proof.* Use the Inverse Function Theorem in the right way.\* Why is the formula for  $Dg$  correct? Assuming the implicit function exists and is differentiable,  $g: W \rightarrow U$

$$\begin{aligned} h(a) = f(g(a), a) = 0 \quad \forall a \in W &\Rightarrow 0 = Df(g(a), a)(a_0) (= Dh(a_0)) \\ &= D_x f(x_0, a_0) Dg(a_0) + D_a f(x_0, a_0) \\ Dg(a_0) &= -[D_x f(x_0, a_0)]^{-1} D_a f(x_0, a_0) \end{aligned}$$

□

\* Look at

$$\begin{aligned} F(x, a) &= (f(x, a), a) \\ \text{So } F: \mathbb{R}^n \times \mathbb{R}^p &\rightarrow \mathbb{R}^n \times \mathbb{R}^p \end{aligned}$$

Show  $DF(x_0, a_0)$  invertible  $\Leftrightarrow D_x f(x_0, a_0)$  invertible

Then use Inverse F. Thm on  $F$ .



## Implicit Function Theorem

**Corollary 1.** Suppose  $X \subseteq \mathbf{R}^n$  and  $A \subseteq \mathbf{R}^p$  are open and  $f : X \times A \rightarrow \mathbf{R}^n$  is  $C^1$ . If 0 is a regular value of  $f(\cdot, a_0)$ , then the correspondence

$$a \mapsto \{x \in X : f(x, a) = 0\} = \gamma(a)$$

is lower hemicontinuous at  $a_0$ .

*Proof.* If 0 is a regular value of  $f(\cdot, a_0)$ , then given any  $x_0 \in \{x \in X : f(x, a_0) = 0\}$ , we can find a local implicit function  $g$ ; in other words, if  $a$  is sufficiently close to  $a_0$ , then  $g(a) \in \{x \in X : f(x, a) = 0\}$ ; the continuity of  $g$  then shows that the correspondence  $\{x \in X : f(x, a) = 0\}$  is lower hemicontinuous at  $a_0$ .  $\square$

# Implicit Function Theorem

**Example:** Back to our opening example:  $f : (0, 2\pi) \times \mathbf{R} \rightarrow \mathbf{R}$  given by  $f(x, a) = \sin x + a$ . Let  $x_0 = \pi$  and  $a_0 = 0$ .

Then  $f(x_0, a_0) = \sin \pi = 0$  and  $D_x f(x_0, a_0) = \cos \pi = -1 \neq 0$ .  
So  $x_0 = \pi$  is a regular point of  $f(\cdot, a_0)$ .  $= f_{a_0}(\cdot)$

By the Implicit Function Theorem,  $\exists$  open neighborhoods  $U \ni \pi$  and  $W \ni 0$  and a  $C^1$  function  $h : W \rightarrow U$  such that  $f(h(a), a) = 0$  for every  $a \in W$ , and

$$Dh(a_0) = -[\cos \pi]^{-1} \cdot 1 = 1$$

So the local solution is increasing in  $a$  near  $a_0$  (as we saw above).



Again at  $x = 3\pi/2$  and  $a = 1$ ,  $D_x f(x, a) = 0$  and no local implicit function exists:

for every open neighborhood  $U$  of  $3\pi/2$  and  $W$  of  $1$ , for any  $a' > 1$  there is no  $x' \in U$  such that  $f(x', a') = \sin x' + a' = 0$ .

- $g(x) = \sin x$

critical values =  $\{-1, 1\}$

regular values =  $\mathbb{R} \setminus \{-1, 1\}$

- $f(x, a) = \sin x + a$

0 regular value of  $f(\cdot, a)$   $\forall a \notin \{-1, 1\}$



## Lebesgue Measure Zero

**Definition 2.** Suppose  $A \subseteq \mathbf{R}^n$ .  $A$  has Lebesgue measure zero if for every  $\varepsilon > 0$  there is a countable collection of rectangles  $I_1, I_2, \dots$  such that

$$\sum_{k=1}^{\infty} \text{Vol}(I_k) < \varepsilon \text{ and } A \subseteq \bigcup_{k=1}^{\infty} I_k$$

Here by a rectangle we mean  $I_k = \times_{j=1}^n (a_j^k, b_j^k)$  for some

$a_j^k < b_j^k \in \mathbf{R}$ , and

$$= \{x \in \mathbf{R}^n : x_j \in (a_j^k, b_j^k) \forall j\}$$

define

$$\text{Vol}(I_k) = \prod_{j=1}^n |b_j^k - a_j^k|$$

Notice that this defines Lebesgue measure zero without defining Lebesgue measure.

# Lebesgue Measure Zero

## Examples:


1. “Lower-dimensional” sets have Lebesgue measure zero. For example,

$$A = \{x \in \mathbf{R}^2 : x_2 = 0\}$$

has measure zero.


$$\sum \frac{1}{2^k} = 2$$

Given  $\varepsilon > 0$ :



Choose  $h_k$  s.t.

$$\sum_{k=1}^{\infty} h_k < \varepsilon.$$



e.g. then:  $I^n = (n, n+2) \times \left(-\frac{\varepsilon}{2^{n+5}}, \frac{\varepsilon}{2^{n+5}}\right)$

$$\text{vol}(I^n) = \frac{\varepsilon}{2^{n+3}}$$


$$\sum_{n=0}^{\infty} \text{vol}(I^n) = \varepsilon \sum_{n=0}^{\infty} \frac{1}{2^{n+3}} = \varepsilon/4$$



•  $\{I^{-n} : n \in \mathbb{N} \cup \{0\}\} \cup \{I^n : n \in \mathbb{N} \cup \{0\}\}$  is countable

•  $A \subseteq \bigcup_n (I^{-n} \cup I^n) = (\bigcup_n I^{-n}) \cup (\bigcup_n I^n)$

•  $\sum_n \text{vol}(I^{-n}) + \sum_n \text{vol}(I^n) = \varepsilon/4 + \varepsilon/4 = \varepsilon/2 < \varepsilon$



e.g.  $|A| = k$ ,  $A \subseteq \mathbb{R}^2$

$x = (x_1, x_2) \in A$

$\forall n: I_x = (x_1 - \frac{\epsilon}{2}, x_1 + \frac{\epsilon}{2}) \times (x_2 - \frac{\epsilon}{k 2^{n+2}}, x_2 + \frac{\epsilon}{k 2^{n+2}})$

$\text{Vol}(I_x) = \epsilon / k 2^{n+1}$

$\sum_{n=1}^{\infty} \text{Vol}(I_x) = \epsilon / 2k$


2. Any finite set has Lebesgue measure zero in  $\mathbf{R}^n$ .
3. If  $A_n$  has Lebesgue measure zero  $\forall n$  then  $\cup_{n \in \mathbf{N}} A_n$  has Lebesgue measure zero.
4.  $\mathbf{Q}$  and every countable set has Lebesgue measure zero.



*with standard metric*

5. No open set in  $\mathbf{R}^n$  has Lebesgue measure zero.


If  $O \subset \mathbf{R}^n$  is open, then there exists a rectangle  $R$  such that  $\bar{R} \subseteq O$  and such that  $\text{Vol}(R) = r > 0$ . If  $\{I_j\}$  is any collection of rectangles such that  $O \subseteq \bigcup_{j=1}^{\infty} I_j$ , then  $\bar{R} \subseteq O \subseteq \bigcup_{j=1}^{\infty} I_j$ , so  $\sum_{j=1}^{\infty} \text{Vol}(I_j) \geq \text{Vol}(R) = r > 0$ .




# Genericity

Lebesgue measure zero is a natural formulation of the notion that  $A$  is a small set. Without specifying a probability measure explicitly, this expresses the idea that if  $x \in \mathbf{R}^n$  is chosen at random, then the probability that  $x \in A$  is zero.

A function may have many critical points; for example, if a function is constant on an interval, then every element of the interval is a critical point. But it can't have many critical *values*.

 a sufficiently well-behaved function



## Sard's Theorem

**Theorem 3** (Thm. 2.4, Sard's Theorem). *Let  $X \subseteq \mathbf{R}^n$  be open, and  $f : X \rightarrow \mathbf{R}^m$  be  $C^r$  with  $r \geq 1 + \max\{0, n - m\}$ . Then the set of all critical values of  $f$  has Lebesgue measure zero.*

*Proof.* First, we give a false proof that conveys the essential idea as to why the theorem is true; it can be turned into a correct proof. Suppose  $m = n$ . Let  $C$  be the set of critical points of  $f$ ,

$f(C) = V$  the set of critical values. Then

$$\begin{aligned}
 \text{Vol}(V) &= \text{Vol}(f(C)) = \int_{f(C)} 1 \, dy \\
 &\leq \int_C |\det Df(x)| \, dx \quad (\text{equality if } f \text{ is one-to-one}) \\
 &= \int_C 0 \, dx \\
 &= 0
 \end{aligned}$$

*linear*

Now, we outline how to turn this into a proof. First, show that we can write  $X = \cup_{j \in \mathbb{N}} X_j$ , where each  $X_j$  is a compact subset of  $[-j, j]^n$ . Let  $C_j = C \cap X_j$ . Fix  $j$  for now. Since  $f$  is  $C^1$ ,

$$\begin{aligned} x_k \rightarrow x &\Rightarrow \det Df(x_k) \rightarrow \det Df(x) \\ \text{det } Df(x_k) = 0 &\Rightarrow \{x_k\} \subseteq C_j, x_k \rightarrow x \Rightarrow \det Df(x) = 0 \Rightarrow x \in C_j \end{aligned}$$

so  $C_j$  is closed, hence compact. Since  $X$  is open and  $C_j$  is compact, there exists  $\delta_1 > 0$  such that

$$B_{\delta_1}[C_j] = \cup_{x \in C_j} B_{\delta_1}[x] \subseteq X$$

$B_{\delta_1}[C_j]$  is bounded, and, using the compactness of  $C_j$ , one can show it is closed, so it is compact. Since  $\det Df(x)$  is continuous on  $B_{\delta_1}[C_j]$ , it is uniformly continuous on  $B_{\delta_1}[C_j]$ . Then given  $\varepsilon > 0$ , we can find  $\delta \leq \delta_1$  such that  $B_\delta[C_j] \subseteq [-2j, 2j]^n$  and

$$x \in B_\delta[C_j] \Rightarrow |\det Df(x)| \leq \frac{\varepsilon}{2 \cdot 4^n j^n}$$

Then

$$\begin{aligned}
 f(C_j) &\subseteq f(B_\delta[C_j]) \\
 \int_{f(B_\delta[C_j])} 1 \, dx &= \text{Vol}(f(B_\delta[C_j])) \leq \int_{[-2j, 2j]^n} \frac{\varepsilon}{2 \cdot 4^n j^n} dx \\
 &= \frac{\varepsilon}{2}
 \end{aligned}$$

Since  $f$  is  $C^1$ , show that  $f(C_j)$  can be covered by a countable collection of rectangles of total volume less than  $\varepsilon$ . Since  $\varepsilon > 0$  is arbitrary,  $f(C_j)$  has Lebesgue measure zero. Then

$$f(C) = f\left(\bigcup_{j \in \mathbf{N}} C_j\right) = \bigcup_{j \in \mathbf{N}} f(C_j)$$

is a countable union of sets of Lebesgue measure zero, so  $f(C)$  has Lebesgue measure zero.  $\square$

# Sard's Theorem

**Remark:** Sard's Theorem has a number of powerful implications. Given a randomly chosen function  $f$ , it is very unlikely that zero will be a critical value of  $f$ . If by some fluke zero *is* a critical value of  $f$ , then a slight perturbation of  $f$  will make zero a regular value. We return to a more wide-ranging version of this statement below.

**Example:** Let  $g : (0, 2\pi) \rightarrow \mathbf{R}$  be given by  $g(x) = \sin x$ . We calculated by hand above that the set of critical values of  $g$  is  $\{-1, 1\}$ . Since this set is finite, it has Lebesgue measure zero.




# Transversality

Let  $g : \mathbf{R}^n \rightarrow \mathbf{R}^n$  be  $C^1$ . Consider the family of  $n$  equations in  $n$  variables:

$$g(x) = 0$$

Suppose for some  $x$  such that  $g(x) = 0$ ,  $\text{rank}(Dg(x)) < n$ . That is, some  $x \in g^{-1}(0)$  is a critical point of  $g$ , thus 0 is a critical value of  $g$ .

By Sard's Theorem, almost every  $a \neq 0$  is a regular value of  $g$ . So for  $a$  outside a set of Lebesgue measure 0,  $Dg(x)$  has full rank for every  $x$  solving  $g(x) = a$ . For any such  $a$  and any  $x \in g^{-1}(a)$ , we can use the Inverse Function Theorem to show that a local inverse  $x(a)$  exists, and give a formula for  $Dx(a)$ .



# Transversality

Suppose  $f : \mathbf{R}^n \times \mathbf{R}^p \rightarrow \mathbf{R}^m$ . We care about the parameterized family of equations

$$f(x, a) = 0$$

where, as above, we interpret  $a \in \mathbf{R}^p$  to be a vector of parameters that indexes the function  $f(\cdot, a)$ .

For a given  $a$ , we are interested in the set of solutions

$$\psi(a) = \{x \in X : f(x, a) = 0\}$$

and the way that this correspondence depends on  $a$ .

If  $f$  is separable in  $a$ , that is,  $f(x, a) = g(x) + a$ , then we can use Sard's Theorem (PS6 2010).

# Transversality Theorem

$$f(x, a) = g(x) + a$$

$\uparrow$

Separability is strong, and not required: If  $f$  depends on  $a$  in a nonseparable fashion, it is enough that from any solution  $f(x, a) = 0$ , any directional change in  $f$  can be achieved by arbitrarily small changes in  $x$  and  $a$ .

**Theorem 4** (Thm. 2.5', Transversality Theorem). *Let  $X \subseteq \mathbf{R}^n$  and  $A \subseteq \mathbf{R}^p$  be open, and  $f : X \times A \rightarrow \mathbf{R}^m$  be  $C^r$  with  $r \geq 1 + \max\{0, n - m\}$ . Suppose that 0 is a regular value of  $f$ . Then there is a set  $A_0 \subseteq A$  such that  $A \setminus A_0$  has Lebesgue measure zero and for all  $a \in A_0$ , 0 is a regular value of  $f_a = f(\cdot, a)$ .*

**Remark:** Notice the important difference between the statement that 0 is a regular value of  $f$  (one of the assumptions of the Transversality Theorem), and the statement that 0 is a regular value of  $f_a$  for a fixed  $a \in A_0$  (part of the conclusion of the Transversality Theorem). 0 is a regular value of  $f$  if and only if  $Df(x, a)$  has full rank for every  $(x, a)$  such that  $f(x, a) = 0$ . Instead, for fixed  $a_0 \in A_0$ , 0 is a regular value of  $f_{a_0} = f(\cdot, a_0)$  if and only if  $D_x f(x, a_0)$  has full rank for every  $x$  such that  $f_{a_0}(x) = f(x, a_0) = 0$ .

**Remark:** Consider the important special case in which  $n = m$ , so we have as many equations ( $m$ ) as endogenous variables ( $n$ ). In this case, suppose  $f$  is  $C^1$  (note that  $1 = 1 + \max\{0, n - n\}$ ). If 0 is a regular value of  $f$ , that is,  $Df(x, a)$  has rank  $n = m$  for every  $(x, a)$  such that  $f(x, a) = 0$ , then by the Transversality Theorem there is a set  $A_0 \subset A$  such that  $A \setminus A_0$  has Lebesgue measure zero and for every  $a_0 \in A_0$ ,  $D_x f(x, a_0)$  has rank  $n = m$  for all  $x$  such that  $f(x, a_0) = 0$ .

Fix  $a_0 \in A_0$  and  $x_0$  such that  $f(x_0, a_0) = 0$ . By the Implicit Function Theorem, there exist open sets  $A^*$  containing  $a_0$  and  $X^*$  containing  $x_0$ , and a  $C^1$  function  $x : A^* \rightarrow X^*$  such that

- $x(a_0) = x_0$

- $f(x(a), a) = 0$  for every  $a \in A^*$
- if  $(x, a) \in X^* \times A^*$  then

$$f(x, a) = 0 \iff x = x(a)$$

that is,  $x_0$  is locally unique, and  $x(a)$  is locally unique for each  $a \in A^*$

- $Dx(a_0) = -[D_x f(x_0, a_0)]^{-1} D_a f(x_0, a_0)$

# Transversality

**Example:** Back to the opening example:  $f : (0, 2\pi) \times \mathbf{R} \rightarrow \mathbf{R}$  given by  $f(x, a) = \sin x + a$ .

For any  $(x, a)$  such that  $f(x, a) = 0$ ,  $Df(x, a) = (\cos x, 1)$  which has rank  $1 = \min\{2, 1\}$ . Thus 0 is a regular value of  $f$ .

Set  $A_0 = \mathbf{R} \setminus \{-1, 1\}$ . Since  $\{-1, 1\}$  is a finite set, it has Lebesgue measure zero in  $\mathbf{R}$ .

Again we have already calculated by hand that for any  $a \in A_0$ , 0 is a regular value of  $f_a = f(\cdot, a)$ .