

Estimators for Persistent and Possibly Non-Stationary Data with Classical Properties

Yuriy Gorodnichenko*

Anna Mikusheva[†]

Serena Ng[‡]

First Draft: June 2007

This Draft: March 2011

Abstract

This paper considers a moments based estimator that is \sqrt{T} consistent and uniformly asymptotically normal irrespective of the degree of persistence of the forcing process. These properties hold for linear autoregressive models, linear predictive regressions, as well as certain non-linear dynamic models. Asymptotic normality is obtained because the moments are chosen so that the objective function is uniformly bounded in probability and that a central limit theorem can be applied. Critical values from the normal distribution can be used irrespective of the treatment of the deterministic terms. Simulations show that the estimates are precise, and the t -test has good size in the parameter region where the least squares estimates usually yield distorted inference.

JEL Classification: C1

Keywords: Persistence, quasi-differencing, unit root, predictive regression, local-to-unity

*Department of Economics, University of California, Berkeley, CA 94720 Email: ygorodni@econ.berkeley.edu.

[†]Department of Economics, M.I.T., E52-50 Memorial Drive, Cambridge, MA 02142. Email: amikushe@mit.edu.

[‡]Dept. of Economics, Columbia University, 420 W 118 St., New York, NY 10027 Email: Serena.Ng@columbia.edu. The authors thank the editor and three referees for many helpful comments and suggestions. Mikusheva acknowledges financial support from the Castle-Krob Career Development Chair. Ng acknowledges financial support from the National Science Foundation (SES-0962431).

1 Introduction

This paper considers a quasi-differencing (QD) framework that can yield \sqrt{T} consistent and uniformly asymptotically normal estimators of autoregressions and multiple regressions when the predictors are persistent and possibly non-stationary. The approach can also be used to estimate dynamic stochastic general equilibrium (DSGE) models. The critical values are invariant to the presence of deterministic trends.

Let θ be an unknown parameter vector and let θ^0 be its true value. The QD approach yields estimators that can generically be defined as

$$\hat{\theta}_J = \underset{\theta \in \Theta}{\operatorname{argmin}} \bar{g}(\theta)' W_T \bar{g}(\theta), \quad (1)$$

where $\bar{g}(\theta)$ is a $J \times 1$ vector of moments, W_T is a $J \times J$ positive-definite matrix, and Θ is a bounded set containing values of θ .¹ The basic premise of QD estimation is that for $\hat{\theta}_J$ to have classical properties, $\bar{g}(\theta)$ needs to be uniformly bounded in probability and a central limit theorem can be applied. To this end, $\bar{g}(\theta)$ is defined to be the difference between the normalized autocovariances of the variables in the model and the data, all quasi-differenced at a persistence parameter that is to be estimated jointly with other parameters of the model. This is a non-standard form of covariance structure estimation because the sample (not just the model-implied) moments depend on unknown parameters. This unique feature makes QD estimation robust to possible non-stationarity in the data.

Achieving asymptotic normality without knowing when the exogenous process has an autoregressive unit root can be very useful in applied work because the answers to many macroeconomic questions are sensitive to assumptions about the nature of the trend and to whether the corresponding regressions are estimated in levels or in first-differences. The price to pay for practical simplicity and robustness is that the proposed estimators are \sqrt{T} consistent rather than super-consistent when the regressors are truly non-stationary. While other asymptotically normal estimators robust to non-stationary regressors are available, they apply only to specific linear models. The QD estimation framework is general and can be used whenever the variables can be quasi-differenced in the way discussed below.

We establish uniform asymptotic normality of QD-based estimators in many different settings. Throughout, we use the notion of uniformity given in Mikusheva (2007a), who studied uniform coverage properties of various inference procedures for the AR(1) model.

¹The optimization is performed over an expanded neighborhood of the set of admissible values of θ so that the parameter of interest is not on the boundary of the support. For the AR(1) model, the admissible values are $(-1 + \delta, 1]$ where $\delta > 0$. We optimize over $\Theta = [C_1, C_2]$, where $C_1 < -1 + \delta < 1 < C_2$. When the context is clear, Θ will not be explicitly specified.

Definition 1 A family of distributions $F_{\theta,T}^{(1)}(x) = P_{\theta,T}\{\xi_1 < x\}$ is asymptotically approximated by (converges to) a family of distributions $F_{\theta}^{(2)}(x) = P_{\theta}\{\xi_2 < x\}$ **uniformly** over $\theta \in \Theta$ if

$$\lim_{T \rightarrow \infty} \sup_{\theta \in \Theta} \sup_x \left| F_{\theta,T}^{(1)}(x) - F_{\theta}^{(2)}(x) \right| = 0.$$

In our analysis, $F_{\theta}^{(2)}(x)$ is in the family of Gaussian distributions.

The paper is structured as follows. Section 2 provides a rigorous analysis of the AR(1) model. Section 3 extends the analysis to AR(p) models, while Section 4 studies predictive regressions. Section 5 considers non-linear estimation of structural parameters. Simulations are presented in Section 6. The relation of QD estimation to other \sqrt{T} -consistent estimators is discussed in Section 7. All proofs are in the Appendix.

As a matter of notation, we let ε_t be the deviation between the dependent variable and the conditional mean evaluated at the true parameter value, while e_t is the deviation evaluated at an arbitrary value of the parameter vector. The indicator function $\mathbb{I}(a)$ is one if a is true and zero otherwise. We let $W(\cdot)$ be the standard Brownian motion and use \Rightarrow to denote weak convergence.

2 The AR(1) Model

To systematically motivate the idea behind QD estimation, we begin with the simple AR(1) model with parameter α and whose true value is α^0 . For $t = 1, \dots, T$, the data are generated by

$$y_t = \alpha^0 y_{t-1} + \varepsilon_t, \quad y_0 = 0. \quad (2)$$

Assumption A $(\varepsilon_t, \mathcal{F}_t)$ is a stationary ergodic martingale-difference sequence with conditional variance $E(\varepsilon_t^2 | \mathcal{F}_{t-1}) = \sigma^2 = \gamma_0$ and $E((\varepsilon_t^2 - \sigma^2)^2 | \mathcal{F}_{t-1}) = \mu_4$.

The least squares estimator $\hat{\alpha}_{OLS}$ is defined as the solution to $\bar{g}(\alpha) = 0$, where $\bar{g}(\alpha) = \frac{1}{T} \sum_{t=1}^T (y_t - \alpha y_{t-1}) y_{t-1}$ is the sample analog of the moment condition

$$Eg_t(\alpha) = E[(y_t - \alpha y_{t-1}) y_{t-1}].$$

When $\alpha^0 < 1$, $\hat{\alpha}_{OLS}$ is \sqrt{T} consistent and asymptotically normal. While $\hat{\alpha}_{OLS}$ is super-consistent at $\alpha^0 = 1$, its distribution is non-standard which makes inference difficult. In particular, the t -statistic for testing $\alpha^0 = 1$ is non-normal in finite samples and converges to the so-called Dickey-Fuller distribution. The issue of non-standard inference arises because of two problems. First, when $\alpha^0 = 1$, the sample moment evaluated at a value $\alpha \neq \alpha^0 = 1$ explodes, and second, the normalized sample moment evaluated at the true value does not obey a central limit theorem. Specifically, $\sqrt{T} \bar{g}(\alpha) = \frac{1}{\sqrt{T}} \sum_{t=1}^T e_t(\alpha) y_{t-1}$ explodes, but $\frac{1}{T} \sum_{t=1}^T e_t(\alpha) y_{t-1} \Rightarrow \sigma^2 \int_0^1 W(r) dW(r)$.

Our starting point is to resolve the second issue by exploiting the autocovariance structure of the errors. Specifically, for $j \geq 1$, it holds that

$$E(\varepsilon_t \varepsilon_{t-j}) = 0. \quad (3)$$

This population moment condition has a sample analog that obeys a central limit theorem. That is, for all true values of α denoted α^0 , $|\alpha^0| \leq 1$, $\varepsilon_t = y_t - \alpha^0 y_{t-1}$,

$$\sqrt{T} \left(\frac{1}{T} \sum_{t=j+1}^T \varepsilon_t \varepsilon_{t-j} - E(\varepsilon_t \varepsilon_{t-j}) \right) \Rightarrow \xi_j \sim N(0, \tilde{\sigma}_j^2). \quad (4)$$

Although ε_t is not observed, we can exploit the autocovariance properties of y_t quasi-differenced at α . Let

$$e_t(\alpha) = y_t - \alpha y_{t-1}.$$

Let E_α denote the expectation taken under the assumption that α is the true value. Then

$$\gamma_j(\alpha) = E_\alpha e_t(\alpha) e_{t-j}(\alpha) = \mathbb{I}(j=0) \sigma^2.$$

In other words, $\gamma_j(\alpha) = 0$ for all $j \geq 1$ when evaluated at the assumed true value of α . Although $\gamma_j(\alpha)$ does not depend on α in this AR(1) model, we keep the explicit dependence of γ_j on α which is useful for subsequent analysis. The sample autocovariances are²

$$\hat{\gamma}_j(\alpha) = \frac{1}{T} \sum_{t=j+1}^T e_t(\alpha) e_{t-j}(\alpha).$$

Let $\bar{g}_{NQD}(\alpha) = (\bar{g}_{0,NQD}(\alpha), \dots, \bar{g}_{J,NQD}(\alpha))'$, where

$$\bar{g}_{j,NQD}(\alpha) = \hat{\gamma}_j(\alpha) - \gamma_j(\alpha).$$

Our NQD (non-classical QD) estimator is defined as

$$\hat{\alpha}_{J,NQD} = \underset{\alpha}{\operatorname{argmin}} \bar{g}_{NQD}(\alpha)' W_T \bar{g}_{NQD}(\alpha).$$

The NQD estimator exploits the difference between the model-implied and the sample autocovariance of $e_t(\alpha)$. In standard covariance structure where $\bar{g}_j(\alpha) = E_\alpha y_t y_{t-j} - \frac{1}{T} \sum_{t=j+1}^T y_t y_{t-j}$, the model implied autocovariance at lag j is a function of α . However, the data y_t depends on α^0 , not α . Our NQD is a non-standard covariance structure estimator because by considering the autocovariances of the quasi-differenced data, both the sample and model moments depend on α .

²To be more precise, we should write $\hat{\gamma}_j(\alpha, \alpha^0)$ because the data are generated under α^0 , and we quasi-difference the data at α . For notational simplicity, the dependence of $\hat{\gamma}_j$ on α^0 is suppressed.

The NQD estimator solves one of the two problems inherent in least squares estimation by making $\bar{g}_{j,NQD}(\alpha^0)$ asymptotically normal for all $|\alpha^0| \leq 1$. However, the estimator still has non-standard properties because the moment $\bar{g}_{j,NQD}(\alpha)$ explodes at $\alpha \neq \alpha^0$ when α^0 is unity or in the neighborhood of one. To resolve this problem, suppose γ_0 is known and define

$$\begin{aligned}\bar{g}_{j,FQD}(\alpha) &= \bar{g}_{j,NQD}(\alpha) - \bar{g}_{0,NQD}(\alpha) \\ &= \left(\hat{\gamma}_j(\alpha) - \hat{\gamma}_0(\alpha) \right) - \left(\gamma_j(\alpha) - \gamma_0 \right), \\ \hat{\alpha}_{J,FQD} &= \underset{\alpha}{\operatorname{argmin}} \bar{g}_{FQD}(\alpha)' W_T \bar{g}_{FQD}(\alpha).\end{aligned}\tag{5}$$

While it is fairly obvious that $\bar{g}_{j,FQD}(\alpha^0)$ obeys a central limit theorem, less obvious but important is that

$$\bar{g}_{j,FQD}(\alpha) = \frac{1}{T} \sum_{t=j+1}^T e_t(\alpha) (e_{t-j}(\alpha) - e_t(\alpha)) + \gamma_0$$

is uniformly bounded in probability for *all* values over which α is optimized, and $\alpha^0 \in (1 - \delta, 1]$. The proof of this fact is given in Lemma A-2 of the Appendix. Thus, while $\hat{\gamma}_j(\alpha)$ is unbounded when $\alpha^0 = 1$ and $\alpha \neq 1$, $\hat{\gamma}_j(\alpha) - \hat{\gamma}_0(\alpha)$ is bounded in the limit even when $\alpha^0 = 1$ and $\alpha \neq \alpha^0$. For this reason, the NQD has very different properties from quasi-differencing using a fixed γ_0 (FQD) in the local-to-unity framework.

To gain insight about the properties of the estimators, we let W_T be an identity matrix. This simplifying assumption makes it possible to obtain useful closed form expressions.

Proposition 1 *Let y_t be generated as in equation (2) with error terms satisfying Assumption A. Assume that W_T is an identity matrix of dimension J .*

- i. Let $n \in \{1, 2\}$ be the number of local minima in the optimization problem $\min_{\alpha} \sum_{j=1}^J \bar{g}_{j,NQD}^2(\alpha)$. In the local-to-unity framework in which $\alpha^0 = 1 + c/T$ with $c \leq 0$, $\hat{\alpha}_{J,NQD}$ is super-consistent and*

$$\begin{pmatrix} \mathbb{I}\{n = 2\} \cdot T^{3/2} (\hat{\alpha}_{J,NQD} - \alpha^0)^2 \\ \mathbb{I}\{n = 1\} \cdot T (\hat{\alpha}_{J,NQD} - \alpha^0) \end{pmatrix} \Rightarrow \begin{pmatrix} \frac{-\xi}{\int_0^1 J_c^2(s) ds} \mathbb{I}\{\xi < 0\} \\ \frac{1+2 \int_0^1 J_c(s) dW(s)}{2 \int_0^1 J_c^2(s) ds} \mathbb{I}\{\xi > 0\} \end{pmatrix}, \tag{6}$$

where J_c is an Ornstein-Uhlenbeck process generated by the Brownian motion W that is independent of $\xi \sim N(0, 1/J)$.

- ii. Let γ_0 be the true value of σ^2 . For any fixed $J > 1$, the estimator $\hat{\alpha}_{J,FQD}$ is consistent. Furthermore, uniformly over $-1 + \delta < \alpha^0 \leq 1$:*

$$\sqrt{T}(\hat{\alpha}_{J,FQD} - \alpha^0) \Rightarrow N(0, \sigma_{J,FQD}^2), \quad \text{where} \quad \sigma_{J,FQD}^2 = \frac{\left(\sum_{j=1}^J (\alpha^0)^{2(j-1)} \right)^2 \frac{\mu_4}{\sigma^4} + \sum_{j=1}^J (\alpha^0)^{2(j-1)}}{\left(\sum_{j=1}^J (\alpha^0)^{2(j-1)} \right)^2}.$$

The NQD estimator is the basis of the estimators we subsequently investigate. It is consistent and has a data dependent convergence rate. Since the objective function is a polynomial of the fourth order, there are multiple solutions. If the realization of data is such that there is a unique minimum to the optimization problem, the convergence rate is T . If there are two minima, a slower convergence rate of $T^{3/4}$ is obtained. In either case, the distribution of $\hat{\alpha}_{J,NQD}$ is not asymptotically normal because $\bar{g}_{NQD}(\alpha)$ is not well behaved for all values of α^0 . However, the problematic term that frustrates a quadratic expansion of $\hat{\gamma}_j(\alpha)$ around α^0 is asymptotically collinear with the corresponding term in $\hat{\gamma}_0(\alpha)$ in the local-to-unity framework. Normalizing each $\hat{\gamma}_j(\alpha)$ by $\hat{\gamma}_0(\alpha)$ and $\gamma_j(\alpha)$ by γ_0 results in an FQD estimator whose asymptotic distribution is normal uniformly over $\alpha^0 \in (-1 + \delta, 1]$. When $J > 1$, the FQD objective function has only one minimum asymptotically.³

The properties for $\hat{\alpha}_{J,FQD}$ are stated assuming the true value of σ^2 is known. The reason why $\gamma_0 = \sigma^2$ is not freely estimated along with α is that doing so would yield multiple solutions. The objective function is zero not only at the true solution $\alpha = \alpha^0, \sigma^2 = \gamma_0$, but also at $\alpha = 1/\alpha^0, \sigma^2 = \gamma_0/(\alpha^0)^2$. Without additional information, the FQD cannot uniquely identify α and σ^2 .

In practice, the true value of σ^2 is not known, and the FQD estimator is infeasible. This can be overcome by finding another moment that can identify σ^2 . Let $\theta = (\alpha, \sigma^2)$ and consider

$$\bar{g}_{QD}(\theta) = \begin{pmatrix} s^2 - \sigma^2 \\ \hat{\gamma}_1(\alpha) - \hat{\gamma}_0(\alpha) - (\gamma_1(\alpha) - s^2) \\ \vdots \\ \hat{\gamma}_J(\alpha) - \hat{\gamma}_0(\alpha) - (\gamma_J(\alpha) - s^2) \end{pmatrix}. \quad (7)$$

where $s^2 = \frac{1}{T}y'My$, and $M = I_T - z(z'z)^{-1}z'$ is the matrix that projects onto the space orthogonal to z with $z_t = y_{t-1}$. Let

$$\hat{\theta}_{J,QD} = (\hat{\alpha}_{J,QD}, \hat{\sigma}_{J,QD}^2) = \underset{\theta}{\operatorname{argmin}} \bar{g}_{QD}(\theta)' W_T \bar{g}_{QD}(\theta)$$

Proposition 2 *Let W_T be an identity matrix of dimension $J + 1$. For any fixed $J > 1$, the estimator $\hat{\alpha}_{J,QD}$ is consistent, and the following convergence holds uniformly over $-1 + \delta < \alpha^0 \leq 1$:*

$$\sqrt{T}(\hat{\alpha}_{J,QD} - \alpha^0) \Rightarrow N(0, \sigma_{J,QD}^2), \text{ where } \sigma_{J,QD}^2 = \frac{\sum_{j=1}^J (\alpha^0)^{2(j-1)}}{\left(\sum_{j=1}^J (\alpha^0)^{2(j-1)}\right)^2}.$$

Estimator $\hat{\alpha}_{J,QD}$ can also be implemented as a two-step estimator in which s^2 is first obtained, and its value is used as γ_0 in the moment function $\bar{g}_{FQD}(\alpha) = (\bar{g}_{1,FQD}(\alpha), \dots, \bar{g}_{J,FQD}(\alpha))'$. The

³When the NQD has two local minima, it does not matter which one is chosen as they are asymptotically symmetric around the true value. This was why we state our result as $T^{3/2}(\hat{\alpha}_{J,NQD} - \alpha^0)^2$ rather than $T^{3/4}(\hat{\alpha}_{J,NQD} - \alpha^0)$. If $J = 1$, the FQD objective function has 2 minima, only one of which is consistent for α^0 .

asymptotic variance of $\hat{\alpha}_{J,QD}$ takes into account the sampling uncertainty of s^2 . The surprising aspect of Proposition 2 is that $\hat{\alpha}_{J,QD}$ does not have an inflated variance as is typical of two-step estimators. Instead, the estimator is more efficient than $\hat{\alpha}_{J,FQD}$ that has a known σ^2 . Pierce (1982) showed in a framework for stationary data that using estimated values of nuisance parameters can yield statistics with smaller variance than if the nuisance parameters were known. This somewhat paradoxical result was also reported by Prokhorov and Schmidt (2009) and Han and Kim (2011) for GMM estimators. Our results suggest that this feature may also arise in the local-to-unity framework.

The closed form expression for the asymptotic variance of $\hat{\alpha}_{J,QD}$ in Proposition 2 was obtained under the assumption that W_T is an identity matrix. For an arbitrary positive-definite weighting matrix, the asymptotic variance of $\hat{\alpha}_{J,QD}$ is the (1,1)-th element of asymptotic variance matrix

$$Avar(\hat{\theta}_{J,QD}) = (G^{0'}WG^0)^{-1}G^{0'}WS^0WG^0(G^{0'}WG^0)^{-1} \quad (8)$$

where W, G^0 and S^0 are the probability limits of W_T , the derivative of $\bar{g}_{QD}(\theta)$ with respect to θ evaluated at θ^0 , and the asymptotic variance of $\bar{g}_{QD}(\theta^0)$, respectively. The asymptotic variance can thus be estimated as though the GMM estimator were developed in the stationary framework under regularity conditions such as those given in Newey and McFadden (1994). In theory, more efficient estimates can be obtained if W_T is an optimal weighting matrix. However, it has been documented in Abowd and Card (1989) and Altonji and Segal (1996) that an optimal weighting matrix may not be desirable for covariance structure estimation.

The key to the classical properties of $\hat{\alpha}_{J,QD}$ is the ability to exploit the autocovariance properties of the quasi-differenced data in an appropriate way. Quasi-differencing has a long tradition in econometrics and underlies GLS estimation. Canjels and Watson (1997) and Phillips and Lee (1996) found that quasi-differencing gives more precise estimates of trend parameters when the errors are highly persistent. Pesavento and Rossi (2006) suggest that for such data, quasi-differencing can improve the coverage of impulse response functions. In both studies, the data are quasi-differenced at $\alpha = \bar{\alpha}$ which is fixed at the value suggested by the local-to-unity framework. In contrast, the present paper estimates this parameter, and both the FQD and the QD estimates have classical properties. These properties also hold in the presence of deterministic terms. Consider the data generating process

$$y_t = d_t + x_t, \quad (9a)$$

$$x_t = \alpha^0 x_{t-1} + \varepsilon_t. \quad (9b)$$

The deterministic terms are captured by $d_t = \sum_{j=0}^r \psi_j t^j$ where r is the order of the deterministic trend function. In the intercept-only case, $d_t = \psi_0$, and in the linear trend case, $d_t = \psi_0 + \psi_1 t$.

Once the parameters of the trend function are consistently estimated, QD estimation proceeds by replacing y_t with demeaned or de-trended data, $\hat{x}_t = y_t - \hat{d}_t$. Let $\hat{e}_t(\alpha) = \hat{x}_t - \alpha\hat{x}_{t-k}$. The sample autocovariances can be constructed as

$$\hat{\gamma}_k(\alpha) = \frac{1}{T} \sum_{t=k+1}^T \hat{e}_t(\alpha)\hat{e}_{t-k}(\alpha).$$

Demeaning and de-trending do not affect the asymptotic distribution of the QD.⁴

The practical appeal of QD estimation is that asymptotic normality permits standard inference. The usual critical values of ± 1.96 and ± 1.64 can be used for two-tailed tests at the 5 and 10 percent significance levels, respectively. We will see in simulations that the size of tests and the coverage of the confidence sets based on the asymptotic normality of $\hat{\alpha}_{J,QD}$ are stable over the parameter set $\alpha^0 \in (-1 + \delta, 1]$.

To recapitulate, the proposed QD estimation of the AR(1) model is based on two simple premises: first, that for all $j \geq 1$, $E(\varepsilon_t \varepsilon_{t-j}) = 0$ and its sample analog obeys a central limit theorem, and, second, that the objective function is uniformly bounded in probability for all values of α and α^0 . The idea can be used whenever the variables can be quasi-differenced to form moment conditions that satisfy these two properties. The next two sections consider the AR(p) model and predictive regressions, respectively. We then show that the quasi-differenced variables can be serially correlated and that the QD framework can be used in non-linear estimations.

3 AR(p) Models

Consider the data generating process

$$y_t = \alpha^0 y_{t-1} + \sum_{j=1}^{p-1} b_j^0 \Delta y_{t-j} + \varepsilon_t. \quad (10)$$

Let $\beta = (\alpha, b_1, \dots, b_{p-1})$ be a $p \times 1$ parameter vector of interest. The true parameter vector is denoted β^0 , and the correct lag length is denoted p . Let $|\lambda_1| \leq |\lambda_2| \dots \leq |\lambda_p|$ be defined implicitly by the identity

$$1 - \alpha L - \sum_{j=1}^{p-1} b_j L^j (1 - L) = (1 - \lambda_1 L) \dots (1 - \lambda_p L).$$

We restrict the parameter set in such a way that the $p-1$ smallest roots do not exceed δ in absolute value for some fixed $0 < \delta < 1$. If the largest root exceeds δ in absolute value, then it is positive and not larger than 1.

⁴De-trending does not affect the asymptotic distribution of FQD, but the J_c in the distribution of the NQD estimator will depend on d_t . In the intercept only case, one should use the de-meaned Ornstein-Uhlenbeck process $\bar{J}_c(r) = J_c(r) - \int_0^1 J_c(s) ds$. In the linear trend case, the de-trended process is $\tilde{J}_c(r) = J_c(r) - \int_0^1 (4 - 6s) J_c(s) ds - r \int_0^1 (12 - 6s) J(s) ds$.

Definition 2 The parameter set \mathfrak{R}_δ consists of all β such that the corresponding roots satisfy the following two conditions: (i) $|\lambda_{p-1}| < \delta$, (ii) if $\lambda_p \in \mathbb{R}$, then $-\delta \leq \lambda_p \leq 1$.⁵

Define the quasi-differenced series, $e_t(\beta)$, by

$$e_t(\beta) = y_t - \alpha y_{t-1} - \sum_{j=1}^{p-1} b_j \Delta y_{t-j}.$$

Obviously, $e_t(\beta^0) = \varepsilon_t$ is white noise, but $e_t(\beta)$ is serially correlated when $\beta \neq \beta^0$. Thus, as in the AR(1) model, the model-implied autocovariances satisfy

$$\gamma_j(\beta) = E_\beta (e_t(\beta)e_{t-j}(\beta)) = 0, \quad j \geq 1 \quad \forall \beta \in \mathfrak{R}_\delta$$

with $\gamma_0(\beta) = \sigma^2$. The sample autocovariances of $e_t(\beta)$ are

$$\hat{\gamma}_j(\beta) = \frac{1}{T} \sum_{t=j+p+1}^T e_t(\beta)e_{t-j}(\beta).$$

Let $s^2 = \frac{1}{T} y' M y$ where M projects onto the space orthogonal to the one spanned by $X_t = (y_{t-1}, \Delta y_{t-1}, \dots, \Delta y_{t-p+1})'$, $t = 1, \dots, T$. Let γ_0 be the true value of σ^2 . Define

$$\hat{\beta}_{J,FQD} = \underset{\beta}{\operatorname{argmin}} \bar{g}_{FQD}(\beta)' W_T \bar{g}_{FQD}(\beta),$$

where

$$\bar{g}_{FQD}(\beta) = \begin{pmatrix} \bar{g}_{1,FQD}(\beta) \\ \vdots \\ \bar{g}_{J,FQD}(\beta) \end{pmatrix} = \begin{pmatrix} \hat{\gamma}_1(\beta) - \hat{\gamma}_0(\beta) - (\gamma_1(\beta) - \gamma_0) \\ \vdots \\ \hat{\gamma}_J(\beta) - \hat{\gamma}_0(\beta) - (\gamma_J(\beta) - \gamma_0) \end{pmatrix}.$$

Define

$$(\hat{\beta}_{J,QD}, \hat{\sigma}_{J,QD}^2) = \underset{\beta, \sigma^2}{\operatorname{argmin}} \bar{g}_{QD}(\beta, \sigma^2)' W_T \bar{g}_{QD}(\beta, \sigma^2),$$

where

$$\bar{g}_{QD}(\beta, \sigma^2) = \begin{pmatrix} \bar{g}_{0,QD}(\beta) \\ \bar{g}_{1,QD}(\beta) \\ \vdots \\ \bar{g}_{J,QD}(\beta) \end{pmatrix} = \begin{pmatrix} s^2 - \sigma^2 \\ \hat{\gamma}_1(\beta) - \hat{\gamma}_0(\beta) - (\gamma_1(\beta) - s^2) \\ \vdots \\ \hat{\gamma}_J(\beta) - \hat{\gamma}_0(\beta) - (\gamma_J(\beta) - s^2) \end{pmatrix}.$$

Proposition 3 Let y_t be generated as in equation (10) with error terms satisfying Assumption A. Let $a_j = E[(X_{t+j} + X_{t-j} - 2X_t)\varepsilon_t]$ and $G = \left(\sum_{j=1}^J a_j a_j'\right)^{-1}$. For any fixed $J > p > 1$, the estimators $\hat{\beta}_{J,QD}$ and $\hat{\beta}_{J,FQD}$ are consistent. Furthermore, when W_T is an identity weighting matrix, the following results hold uniformly over $\beta^0 \in \mathfrak{R}_\delta$:

⁵The optimization in equation (1) is done over a bounded set that includes a neighborhood of \mathfrak{R}_δ in order to avoid the boundary problem.

- (i) $\sqrt{T}(\widehat{\beta}_{J,FQD} - \beta^0) \Rightarrow N(0, \Sigma_{J,FQD})$, where $\Sigma_{J,FQD} = \sigma^4 G + \mu_4 G \left(\sum_{j=1}^J a_j \right) \left(\sum_{j=1}^J a_j \right)' G$;
(ii) $\sqrt{T}(\widehat{\beta}_{J,QD} - \beta^0) \Rightarrow N(0, \Sigma_{J,QD})$, where $\Sigma_{J,QD} = \sigma^4 G$.

The proof is a generalization of Propositions 1 and 2. A sketch of the arguments is as follows. From the definition that $e_t(\beta) = \varepsilon_t + (\beta^0 - \beta)' X_t$, we have

$$\widehat{\gamma}_j(\beta) = \frac{1}{T} \sum_{t=j+p+1}^T \varepsilon_t \varepsilon_{t-j} + (\beta^0 - \beta)' \frac{1}{T} \sum_{t=j+p+1}^T \left(X_t \varepsilon_{t-j} + X_{t-j} \varepsilon_t \right) + (\beta^0 - \beta)' \frac{1}{T} \sum_{t=j+p+1}^T X_t X_{t-j}' (\beta^0 - \beta).$$

The moment function can be rewritten as

$$\bar{g}_{j,FQD}(\beta) = A_{j,FQD} + (\beta^0 - \beta)' B_{j,FQD} + (\beta^0 - \beta)' C_{j,FQD} (\beta^0 - \beta).$$

The thrust of the proof is to show that for each $1 \leq j \leq J$ uniformly over \mathfrak{R}_δ ,

$$A_{j,FQD} = \left(\frac{1}{T} \sum_{t=j+p+1}^T \varepsilon_t \varepsilon_{t-j} - \gamma_j(\beta) \right) - \left(\frac{1}{T} \sum_{t=p+1}^T \varepsilon_t^2 - \gamma_0(\beta^0) \right) = O_p(T^{-1/2}), \quad (11)$$

$$B_{j,FQD} = \frac{1}{T} \sum_{t=j+p+1}^T \left(X_t \varepsilon_{t-j} + X_{t-j} \varepsilon_t - 2X_t \varepsilon_t \right) \rightarrow^p a_j, \quad (12)$$

$$C_{j,FQD} = \frac{1}{2T} \sum_{t=j+p+1}^T (X_t X_{t-j}' + X_{t-j} X_t' - 2X_t X_t') = O_p(1). \quad (13)$$

Equations (11)-(13) imply that the function $\bar{g}_{j,FQD}(\beta)$ is bounded in probability uniformly for all β in the optimization set and $\beta_0 \in \mathfrak{R}_\delta$. It also follows from equations (11)-(13) that

$$\frac{\partial \bar{g}_{j,FQD}}{\partial \beta}(\widehat{\beta}_{J,FQD}) \rightarrow^p a_j,$$

and

$$\sqrt{T} \bar{g}_{j,FQD}(\widehat{\beta}_{J,FQD}) = \sqrt{T} A_{j,FQD} + a_j' \sqrt{T} (\widehat{\beta}_{J,FQD} - \beta^0) + o_p(1).$$

The first order condition for the optimization problem implies:

$$\sqrt{T}(\widehat{\beta}_{J,FQD} - \beta^0) = G^{-1} \sum_{j=1}^J \sqrt{T} A_{j,FQD} a_j + o_p(1)$$

In view of equation (4), $\sqrt{T} A_{j,FQD} = \sqrt{T} \bar{g}_{j,FQD}(\beta^0) \Rightarrow \xi_j - \xi_0$ uniformly over \mathfrak{R}_δ , part (i) of the proposition follows. Part (ii) uses a similar argument with one exception: $A_{j,QD} = A_{j,FQD} + s^2 - \sigma^2 = \frac{1}{T} \sum_{t=j+1}^T \varepsilon_t \varepsilon_{t-j}$. As in the AR(1) case, $\widehat{\beta}_{J,QD}$ has a smaller variance than $\widehat{\beta}_{J,FQD}$. Furthermore, one can use other weighting matrices in the estimation. The asymptotic variance of $\widehat{\beta}_{J,QD}$ can be computed from the expression given in equation (8).

4 Predictive Regressions

Consider the predictive regression with scalar predictor x_{t-1} :

$$y_t = \beta^0 x_{t-1} + \varepsilon_{yt} \quad (14a)$$

$$x_t = \alpha^0 x_{t-1} + \varepsilon_{xt}. \quad (14b)$$

If $\alpha^0 = 1$, then $(1; -\beta^0)$ is a cointegrating vector, and ordinary least squares provide super-consistent estimates but inference is non-standard. Unfortunately, the finite sample distribution of $\widehat{\beta}_{OLS}$ is not well approximated by the normal distribution if x_t is highly persistent. The challenge is how to conduct inference robust to the dynamic properties of x_t . Let $\varepsilon_t = (\varepsilon_{yt}, \varepsilon_{xt})'$ be a martingale-difference sequence with $E(\varepsilon_t \varepsilon_t' | \mathcal{F}_{t-1}) = \Omega^0 = \begin{pmatrix} \sigma_{yy} & \sigma_{yx} \\ \sigma_{yx} & \sigma_{xx} \end{pmatrix}$. Consider quasi-differencing the data at $\theta = (\beta, \alpha)$ to obtain

$$e_{yt}(\theta) = y_t - \beta x_{t-1},$$

$$e_{xt}(\theta) = x_t - \alpha x_{t-1}.$$

Now $Y_t = \theta^0 x_{t-1} + \varepsilon_t$ where $Y_t = \begin{pmatrix} y_t \\ x_t \end{pmatrix}$ and $\theta^0 = (\beta^0, \alpha^0)'$. Let $e_t(\theta) = \begin{pmatrix} e_{yt}(\theta) \\ e_{xt}(\theta) \end{pmatrix}$. Then

$$e_t(\theta) = (\theta^0 - \theta)x_{t-1} + \varepsilon_t.$$

Let $\Gamma_j(\theta) = E_\theta(e_t(\theta)e_{t-j}(\theta)')$ where E_θ denotes the expectation under the assumption that θ is the true value. The model implies

$$\Gamma_j(\theta) = 0, \quad j \neq 0,$$

$$\Gamma_0(\theta) = \Omega.$$

The sample autocovariance at lag j is

$$\widehat{\Gamma}_j(\theta) = \frac{1}{T} \sum_{t=j+1}^T e_t(\theta)e_{t-j}(\theta)'$$

A central limit theorem applies to $\sqrt{T} \widehat{\Gamma}_j(\theta^0)$. Evaluating Γ_0 at the true value of Ω and letting $S = \frac{1}{T} Y'(I_2 \otimes M)Y$, $M = I_T - z(z'z)^{-1}z'$, $z_t = x_{t-1}$, we can define, for $j = 1, \dots, J$:

$$\bar{g}_{j,FQD}(\theta) = \text{vec}(\widehat{\Gamma}_j(\theta) - \widehat{\Gamma}_0(\theta) - (\Gamma_j(\theta) - \Gamma_0)).$$

Let $\bar{g}_{FQD}(\theta) = (\bar{g}_{1,FQD}(\theta)', \dots, \bar{g}_{J,FQD}(\theta)')$. The FQD estimator is

$$\widehat{\theta}_{J,FQD} = \arg \min_{\theta} \bar{g}_{FQD}(\theta)' W_T \bar{g}_{FQD}(\theta).$$

Analogously, let $\bar{g}_{QD}(\theta, \Omega) = (\bar{g}_{0,QD}(\theta, \Omega)', \dots, \bar{g}_{J,QD}(\theta, \Omega)')$ where

$$\begin{aligned}\bar{g}_{j,QD}(\theta, \Omega) &= \text{vec}(\hat{\Gamma}_j(\theta) - \hat{\Gamma}_0(\theta) - (\Gamma_j(\theta) - S)), \quad j \geq 1 \\ \bar{g}_{0,QD}(\theta, \Omega) &= \text{vech}(S - \Omega).\end{aligned}$$

Define the QD estimator as

$$(\hat{\theta}_{J,QD}, \hat{\Omega}_{J,QD}) = \arg \min_{\theta, \Omega} \bar{g}_{QD}(\theta, \Omega)' W_T \bar{g}_{QD}(\theta, \Omega).$$

Proposition 4 *Let the data be generated according to formulas (14a) and (14b) and error terms be a martingale-difference sequence with $E(\varepsilon_t \varepsilon_t' | \mathcal{F}_{t-1}) = \Omega^0$ and finite four moments. Define $a_j = E[x_{t-1}(\varepsilon_{t-j} - \varepsilon_t)]$. Then for any fixed $J > 1$, the estimators $\hat{\theta}_{J,FQD}$ and $\hat{\theta}_{J,QD}$ are consistent. Furthermore, when W_T is an identity matrix, the following asymptotic results hold uniformly over all possible values of β , and uniformly over all possible values of $\alpha \in (-1 + \delta, 1]$:*

(i) *Let $\Gamma_0 = \Omega^0$. Then $\sqrt{T}(\hat{\theta}_{J,FQD} - \theta^0) \Rightarrow N(0, \Sigma_{J,FQD})$, where*

$$\Sigma_{J,FQD} = \left(\frac{1}{\sum_{j=1}^J a_j' a_j} \right)^2 \sum_{j=1}^J (a_j' \Omega^0 a_j) \Omega^0 + \left(\frac{1}{\sum_{j=1}^J a_j' a_j} \right)^2 E \left[\left(\varepsilon_t' \sum_{j=1}^J a_j \right)^2 \varepsilon_t \varepsilon_t' \right];$$

(ii) *$\sqrt{T}(\hat{\theta}_{J,QD} - \theta^0) \Rightarrow N(0, \Sigma_{J,QD})$ where*

$$\Sigma_{J,QD} = \left(\frac{1}{\sum_{j=1}^J a_j' a_j} \right)^2 \sum_{j=1}^J (a_j' \Omega^0 a_j) \Omega^0.$$

As in the case of autoregressions, the FQD moments alone cannot globally identify both θ and Ω . Thus, the properties of $\hat{\theta}_{J,FQD}$ are stated by evaluating Ω at the true value of Ω^0 . Proposition 4 shows that $\hat{\theta}_{J,QD}$ has classical properties both in the stationary and the local-to-unity framework and is more efficient than the estimator $\hat{\theta}_{J,FQD}$ that uses the known Ω . The QD can be implemented as a sequential estimator in which the covariance matrix of the shocks is obtained from two least squares regressions: one by regressing y_t on x_t to get \hat{e}_{yt} , and another autoregression in x_t to obtain \hat{e}_{xt} .

Proposition 4 has useful implications for applied work because there does not exist an estimator that is robust to the persistent properties of the predictors. The approach of Jansson and Moreira (2006) relies on model-specific conditional critical values and, in any event, their inference procedure does not yield an estimator per se. In contrast, the QD estimator is simple and robust.

The predictive regression can be generalized to accommodate stationary and pre-determined regressors, z_t . Suppose the data generating process is

$$\begin{aligned} y_t &= x_{t-1}\beta^0 + z_t\gamma^0 + \varepsilon_{yt} \\ x_t &= \alpha^0 x_{t-1} + \varepsilon_{xt}, \quad (\varepsilon_{yt}, \varepsilon_{xt})' \sim (0, \Omega), \quad \Omega = \begin{pmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{xy} & \sigma_x^2 \end{pmatrix}. \end{aligned}$$

Let $\theta = (\beta, \alpha)$ and as before, $\Gamma_j = E_\theta(\varepsilon_t \varepsilon_{t-j}') = 0$ for all $j \neq 0$ with $\Gamma_0 = \Omega^0$. Let $\hat{\gamma}_{OLS}$ be obtained from least squares regression of y_t on x_{t-1} and z_t , and let $\hat{\Omega}_{OLS}$ be the estimated covariance matrix of the errors. Since z_t is stationary, the estimator $\hat{\gamma}_{OLS}$ is \sqrt{T} consistent and asymptotically normal uniformly over $\alpha \in (-1 + \delta, 1]$. Define the quasi-differenced sequence

$$e_{yt}(\theta) = y_t - x_{t-1}\beta - z_t\hat{\gamma}_{OLS}, \quad e_{xt}(\theta) = x_t - \alpha x_{t-1}, \quad e_t(\theta) = \begin{pmatrix} e_{yt}(\theta) \\ e_{xt}(\theta) \end{pmatrix}.$$

Let $\hat{\Gamma}_j(\theta) = \frac{1}{T} \sum_{t=j+1}^T e_t(\theta) e_{t-j}(\theta)'$ and define $\bar{g}_{j,QD}(\theta, \Omega)$ as in the absence of z_t . Using arguments analogous to Proposition 4, it can be shown that $\hat{\theta}_{J,QD}$ is still \sqrt{T} consistent and asymptotically normal. Proposition 4 assumes that the regression error ε_{yt} is white noise. This is not restrictive as lags of Δy_t and Δx_t can be added to z_t to control for residual serial correlation.

5 Non-Linear Models and Minimum Distance Estimation

So far, the QD framework has been used to estimate linear models, where the model autocovariances are such that $\gamma_j(\theta) = 0$ for all θ if $j \geq 1$. The analysis also holds for $\gamma_j(\theta) \neq 0$ provided that $\gamma_j(\theta)$ is known or can be computed numerically. For example, if x_t is an ARMA(1,1) instead of an AR(1), $\gamma_j(\theta)$ will depend on the parameters of the model. A more explicit example is now illustrated in the context of a DSGE model.

To fix ideas, consider the simple one sector stochastic growth model presented in Uhlig (1999). Let Q_t, C_t, K_t, I_t be output, consumption, capital stock, and investment, respectively. The problem facing the central planner is to maximize expected utility $E_{t-1} \sum_{t=0}^{\infty} (1 + \rho)^{-t} \log C_t$ subject to the constraints $Q_t = K_{t-1}^\psi Z_t^{1-\psi} = C_t + I_t$ and $K_t = (1 - \delta)K_{t-1} + I_t$ where $u_t = \log Z_t$ evolves as

$$u_t = \alpha u_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim (0, \sigma^2).$$

Denote the deviation of a variable from its mean by lower case letters. Let $Y_t = (y_{1t}, \dots, y_{Nt})'$ be the collection of endogenous variables in the model (such as consumption, output, etc.). As shown in Uhlig (1999), this simple model has an analytic solution:

$$k_t = v_{kk} k_{t-1} + v_{kz} u_t$$

where $v_{kk} < 1$ does not depend on α , but v_{kz} depends on α . For each $y_{nt} \in Y_t$,

$$(1 - v_{kk}L)(1 - \alpha L)y_{nt} = u_t + \vartheta_n u_{t-1}$$

is an ARMA(2,1) with a moving-average parameter ϑ_n that is a function of the structural parameters. Note that all series in Y_t have the same autoregressive dynamics as k_t . The parameters of the linearized solution are $\beta = (v_{kk}, v_{kz}, \vartheta_1, \dots, \vartheta_N)$. The parameters of the model are $\theta = (\psi, \alpha, \sigma^2, \rho, \delta)$. Let Θ be a compact set containing possible values of θ .

Estimation of DSGE models depends on whether the observables are stationary in two ways. First, the model needs to be linearized around a stationary equilibrium whose precise definition depends on the value of α . Second, the likelihood function is not well defined at $\alpha = 1$. Furthermore, likelihood estimation is infeasible when there are more variables than shocks, a problem known as stochastic singularity

We propose to estimate the parameters of the model without making *a priori* assumptions about the degree of persistence of the shocks. Two variations of the QD framework are considered. The *first* method proceeds as follows. For given θ , let $e_t(\theta) = Y_t - \alpha Y_{t-1}$ with $\Gamma_j(\theta) = E(e_t(\theta)e_t(\theta)')$. Define the moment $\omega_j(\theta) = (\Gamma_j(\theta) - \Gamma_0(\theta))$ whose sample analog is $\hat{\omega}_j(\theta) = (\hat{\Gamma}_j(\theta) - \hat{\Gamma}_0(\theta))$. Note that since $e_t(\theta)$ can be serially correlated, $\Gamma_j(\theta)$ need not be a null matrix as in the applications considered thus far. Let $\bar{g}_{QD}(\theta) = (\bar{g}_{1,QD}(\theta)', \dots, \bar{g}_{J,QD}(\theta)')$ where $\bar{g}_{j,QD}(\theta) = \text{vec}(\hat{\omega}_j(\theta) - \omega_j(\theta))$. The QD estimator considered in Gorodnichenko and Ng (2010) is defined as⁶

$$\hat{\theta}_{J,QD} = \underset{\theta \in \Theta}{\text{argmin}} \bar{g}_{QD}(\theta)' \bar{g}_{QD}(\theta). \quad (15)$$

As written, $\hat{\theta}_{J,QD}$ is an equally weighted estimator. An optimal weighting matrix can be used subject to constraints imposed by stochastic singularity. In the one-shock stochastic growth model considered, the autocovariance at lag one of both output and consumption can both be used to construct an efficient $\hat{\theta}_{J,QD}$, but additional autocovariances will not add independent information. In contrast, the use of data for both output and consumption in likelihood estimation would not even be possible.

Another QD-based estimator can be obtained if we entertain the possibility of a reduced form model. Consider a finite order AR(p) model:

$$y_t = a^0 y_{t-1} + \sum_{j=1}^{p-1} b_j^0 \Delta y_{t-j} + \varepsilon_{tp} \quad (16)$$

⁶Identification requires $\text{rank} \frac{\partial \bar{g}_{j,QD}(\theta)}{\partial \theta} = \dim \theta$. Now $\bar{g}_j(\theta)$ depends on θ through the parameters in the solution to expectational equations. Formal identification conditions are given in Komunjer and Ng (2009).

where $\beta^0 = (a^0, b_1^0, \dots, b_{p-1}^0) = \beta(\theta^0)$ are the true ‘reduced-form’ parameters that can be computed analytically or numerically. We also need the following:

Assumptions B. (Identification) (i) there is a unique θ^0 such that $\beta(\theta^0) = \beta^0$; (ii) the function $\beta(\theta)$ is twice continuously differentiable; (iii) $B(\theta^0) = \nabla\beta(\theta^0)$ has full rank $k = \dim(\theta) \leq p$.

The method proceeds as follows. For $y_{nt} \in Y_t$, define $e_{nt}(\beta) = y_{nt} - ay_{nt-1} - \sum_{j=1}^{p-1} b_j y_{nt-j}$ where $\beta = (a, b_1, \dots, b_{p-1})$. Note that Y_t is now quasi-differenced using the ‘reduced-form’ parameter β instead of the structural parameter α as in method 1. Once the data are quasi-differenced, estimation proceeds by defining $\omega_j(\theta; p) = (\Gamma_j(\theta; p) - \Gamma_0(\theta; p))$ with $\bar{g}_{j,QD}(\theta; p) = \hat{\omega}_j(\theta; p) - \omega_j(\theta; p)$ and sample analog as in (15). Let $\bar{g}_{QD}(\theta; p) = (\bar{g}_{1,QD}(\theta; p)', \dots, \bar{g}_{J,QD}(\theta; p)')$. The estimator is

$$\hat{\theta}_{J,QD} = \underset{\theta \in \Theta}{\operatorname{argmin}} \bar{g}_{QD}(\theta; p)' \bar{g}_{QD}(\theta; p). \quad (17)$$

Because β is p -dimensional, this second estimator also depends on the choice of p . Since e_{nt} is not necessarily exactly white noise, $\Gamma_j(\theta)$ will not be zero. However, its autocovariances can be computed for any given θ .

We have presented two uses of the QD framework that can yield estimators that are robust to non-stationary exogenous variables in DSGE models. The data-dependent transformations allow us to construct moments that are uniformly bounded. Applying the central limit theorem to the sample moments yields estimators with classical properties.

6 Simulations

We consider the finite sample properties of the estimators in linear AR(p) models, predictive regressions, and DSGE models. Results for three estimators are presented: OLS, FQD with γ_0 fixed at the true σ_0^2 , and QD. Even though the FQD estimator is infeasible in practice, it is a useful benchmark. The simulations are based on 2,000 replications. We use the standard Newey-West plug-in estimator for the variance of the moments. As starting values, we use 0.9 times the true values of the parameters. The QD estimator requires evaluation of the model implied autocovariances $\Gamma_j(\theta)$. This is straightforward once a model is cast in a state space. For example, the system

$$\begin{bmatrix} y_t \\ x_t \end{bmatrix} = \begin{bmatrix} 0 & \beta \\ 0 & \alpha \end{bmatrix} \begin{bmatrix} y_{t-1} \\ x_{t-1} \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \varepsilon_{yt} \\ \varepsilon_{xt} \end{bmatrix}$$

in quasi-differenced form is

$$\begin{bmatrix} e_{yt}(\theta) \\ e_{xt}(\theta) \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} e_{yt-1}(\theta) \\ e_{xt-1}(\theta) \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \varepsilon_{yt} \\ \varepsilon_{xt} \end{bmatrix}.$$

More generally, every ARMA model has a state-space representation from which a state-space model for the quasi-differenced data can be expressed as

$$w_t = D_0 w_{t-1} + D_1 \varepsilon_t$$

where w_t includes $e_t(\theta)$ (and possibly its lags), and ε_t is the set of exogenous white noise shocks with variance Ω_ε . The variance matrix $\Omega_w(0) = E(w_t w_t')$ can be computed by iterating the equation

$$\Omega_w^{(i)}(0) = D_0 \Omega_w^{(i-1)}(0) D_0' + D_1 \Omega_\varepsilon D_1' \quad (18)$$

until convergence. The autocovariance matrices can then be computed as $\Omega_w(j) = D_0^j \Omega_w(0)$. Now $\Gamma_j(\theta)$ are submatrices of $\Omega_w(j)$. If we are only interested in computing the moments $w_t^d \subset w_t$, we iterate equation (18) until the block that corresponds to w_t^d converges, i.e. $\|\Omega_{w^d}^{(i)}(0) - \Omega_{w^d}^{(i-1)}(0)\| < c$.

AR(p) Models: Data are generated from the AR(2) model:

$$(1 - \lambda_1^0 L)(1 - \lambda_2^0 L)y_t = \varepsilon_t, \quad \varepsilon_t \sim N(0, 1).$$

The process can be written as

$$y_t = \alpha^0 y_{t-1} + b^0 \Delta y_{t-1} + \varepsilon_t.$$

The parameter of interest is $\alpha^0 = \lambda_1^0 + \lambda_2^0 - \lambda_1^0 \lambda_2^0$ with $b^0 = \lambda_1^0 \lambda_2^0$. The OLS estimate of α has a non-standard distribution when the roots are unity, in which case $\alpha^0 = 1$.

We consider estimation of an AR(2) model when $\lambda_2^0 = 0$ (Table 1), and an AR(3) model when $\lambda_2^0 \neq 0$ (Table 2). Demeaned data are used to compute the sample autocovariances in the intercept case, and linearly detrended data are used in the linear trend case. We report the mean of the QD, FQD, and OLS estimates when $T = 200$ and 500 , the J test for overidentifying restrictions, along with the finite sample power for one-sided t tests evaluated at $\alpha = \alpha^0 - 0.05$.

Table 1 shows that all three estimators are precise when $\alpha^0 \leq 0.8$. The t -statistic for the null hypothesis that $\alpha = \alpha^0$ for all three estimators has rejection rates close to the nominal size of 0.05 when $\alpha^0 \leq 0.8$. The picture is, however, very different at larger values of α^0 . The FQD has slightly smaller bias but is much less efficient. While OLS has the largest bias, its root-mean-squared error (RMSE) is much smaller than the FQD. The QD is neither the most accurate nor the most efficient, but has RMSE closer to OLS and much smaller than the FQD, in support of Proposition 2.

Efficiency of OLS comes at the cost of size distortion, however. At $T = 200$, the OLS-based t -statistic has a rejection rate of 0.473 when $\alpha^0 = 1$ and 0.15 when $\alpha^0 = 0.95$, much larger than the nominal size of 0.05. Even at $T = 500$, the rejection rates are 0.462 and 0.127, well above the nominal rate of 5%. The rejection rates for the FQD and QD are 0.107 and 0.08 when $T = 200$,

and are 0.066 and 0.076 when $T = 500$, much closer to the nominal size of 0.05. The QD has accurate rejection rates that are always around 0.05 for all values of α^0 , but it has less power than OLS. Figure 1 plots the distribution of t -statistics for QD at $T=200$ and $T=500$. The normal approximation to the finite sample distribution is good.

The results in Table 2 are based on overparameterized autoregressions, as we estimate an AR(3) when the true model is an AR(2). We fix λ_2^0 to 0.2 and set λ_1^0 such that the sum of the AR(2) parameters is the same as in Table 1. Statistics are reported for estimates of α . The results are similar to those in Table 1: the FQD is inefficient; the t -statistic associated with it has good size but less power. The OLS has large size distortion when α^0 is close to one, but the estimates are tightly estimated. The QD strikes a balance between the two. In the intercept case, the t statistic has very similar size property for all values of $|\alpha^0| \leq 1$ even when $T = 200$. In the linear time trend case, the uniform size property can be achieved with $T = 500$. Importantly, the QD achieves a more accurate size when α^0 is close to unity without sacrificing too much power when α^0 is outside the persistent range. For example, when α^0 is 0.5, the power of OLS and QD are quite similar. The reason is that when α^0 is far from the unit circle, OLS is \sqrt{T} consistent, but so are FQD and QD. Power is also fairly similar for both the intercept-only and the linear trend model.

Predictive Regressions Data for the predictive regressions are generated as in equations (14a) and (14b). We let $\beta^0 = 1$, $\sigma_y^2 = 1$, $\sigma_x^2 = 1$, $\sigma_{yx} = 0.5$. The results, reported in Table 3, are generally similar to the AR(p) case. The QD is slightly less efficient than OLS when α^0 is close to one but no less efficient when α^0 is far from the unit circle. This loss of efficiency arises from the fact that at $\alpha^0 = 1$, OLS is T -consistent but the QD is only \sqrt{T} -consistent. The benefit to this small efficiency loss is robustness. As one can see, the size of the t test for $H_0 : \beta = \beta^0$ is fairly constant around the nominal size of 0.05, but the size of the test based on OLS varies with α^0 . Furthermore, as σ_{yx}^0 rises, the size of the t -test based on OLS becomes increasingly distorted while the QD continues to have sizes close to nominal.

DSGE Models Data are generated from the stochastic growth model presented in Section 5. We fix the true value of capital intensity ψ to 0.25 and σ^2 to 1 and consider five values of α : 1, 0.98, 0.95, 0.9, 0.8. We use data on consumption to estimate ψ , α and σ^2 using the second method discussed in Section 5. This consists of solving for ψ , α and σ from the six autocovariances of $e_t(\beta)$, where β are the parameters of an AR(3) model. For the sake of comparison, we also use the Kalman filter to obtain the maximum likelihood estimates. The results are reported in Table 4. Evidently, the QD estimates of ψ are close to the true value of 0.25, and the size of the t -test is

close to the nominal value for all values of α . In contrast, the MLE estimates are biased when α is close to one, and the t test for the null hypothesis that $\psi = 0.25$ is severely distorted.

7 Relation to other \sqrt{T} -Consistent Linear Estimators

This section shows how the ideas underlying the QD estimation can be used to construct a two-step *linear* estimator for the AR(1) model. We then relate it to other known linear estimators with classical properties in the local-to-unity framework.

As we emphasized in Section 2, one of the problems with the OLS estimator when a unit root is present is that the moment condition at the true value $\sqrt{T}\bar{g}(\alpha^0) = \frac{1}{\sqrt{T}} \sum_{t=1}^T y_{t-1}\varepsilon_t$ does not satisfy a central limit theorem. Although y_{t-1} is orthogonal to ε_t , the persistence of y_{t-1} requires a stronger normalization and standard distribution theory cannot be used. One of the ideas behind the QD estimator is to use moment conditions that satisfy a central limit theorem uniformly over values of α^0 .

For the AR(1) model, the QD moment condition (3) replaces y_{t-1} with ε_{t-j} . As seen from equation (4), the central limit theorem holds whether or not there is a unit root present. But the moment condition can be understood in an instrumental variable setup because ε_{t-j} is uncorrelated with ε_t and is hence a valid instrument. The only problem is that ε_{t-j} is not observed. But $\hat{\alpha}_{OLS}$ is consistent for all $|\alpha^0| \leq 1$. Thus, let $\tilde{e}_{t-1} = y_t - \hat{\alpha}_{OLS}y_{t-1}$, noting that generated instruments do not require a correction for the standard errors like generated regressors do. We can now define a (hybrid differencing) HD estimator using the following moment condition:⁷

$$\bar{g}_{HD}(\hat{\alpha}_{HD}) = \frac{1}{T} \sum_{t=k+1}^T \tilde{e}_{t-k}(y_t - \hat{\alpha}_{HD}y_{t-1}) = 0.$$

This leads to the estimator

$$\hat{\alpha}_{HD} = \frac{\sum_{t=k}^T y_t \tilde{e}_{t-k}}{\sum_{t=k}^T y_{t-1} \tilde{e}_{t-k}} = \alpha^0 + \frac{\sum_{t=k}^T e_t \tilde{e}_{t-k}}{\sum_{t=k}^T y_{t-1} \tilde{e}_{t-k}}.$$

We refer to $\hat{\alpha}^{HD}$ as a hybrid estimator because it is based on the covariance between the quasi-difference of y_t and a stationary random variable. Notice that the HD and QD use the same moment condition. What distinguishes the HD from the QD is that the objective function of the HD is now linear in α . Consistency of $\hat{\alpha}_{HD}$ follows from the fact that $\frac{1}{T} \sum_{t=1}^T e_t(\alpha^0) \tilde{e}_{t-k} \xrightarrow{p} E\varepsilon_t \varepsilon_{t-k} = 0$. It is straightforward to show that in the local-to-unity framework,

$$\sqrt{T}(\hat{\alpha}_{HD} - \alpha^0) \Rightarrow 2(1 + J_c(1)^2)^{-1}N(0, 1).$$

⁷Laroque and Salanie (1997) used two OLS regressions in stationary variables to obtain a \sqrt{T} -consistent estimate of the cointegrating vector.

Once the HD is understood as an instrumental estimator, other possibilities arise. Instead of \tilde{e}_{t-1} , we can use any stationary series uncorrelated with the error term.⁸ For example, using Δy_{t-1} would by-pass the need for a preliminary least squares estimation. The first differencing (FD) estimator is:

$$\hat{\alpha}_{FD} = \frac{\sum_{t=2}^T \Delta y_{t-1} y_t}{\sum_{t=2}^T \Delta y_{t-1} y_{t-1}}.$$

The FD is a special case of estimators analyzed in So and Shin (1999). These authors used the sign of y_{t-1} as instrument x_t to construct

$$\hat{\alpha}_{SS} = \frac{\sum_{t=2}^T x_t y_t}{\sum_{t=1}^T x_t y_{t-1}}.$$

Another estimator with classical properties in the local-to-unity framework is that of Phillips and Han (2008). The PH estimator, defined as

$$\hat{\alpha}_{PH} = \frac{\sum_{t=2}^T \Delta y_{t-1} (2\Delta y_t + \Delta y_{t-1})}{\sum_{t=2}^T (\Delta y_{t-1})^2},$$

has the property that $\sqrt{T}(\hat{\alpha}_{PH} - \alpha^0) \Rightarrow N(0, 2(1 + \alpha^0))$ for all $\alpha^0 \in (-1, 1]$. As shown in the Appendix, the FD estimator is asymptotically equivalent to the PH estimator in the stationary case when α^0 is far from unit circle. That is, for the AR(1) model, $\hat{\alpha}_{PH} = \hat{\alpha}_{FD} + O_p(1/T)$ under stationary classical asymptotics. However, these two estimators differ in the local-to-unity setting. While $\sqrt{T}(\hat{\alpha}_{FD} - \alpha^0) \Rightarrow 2(1 + W(1)^2)^{-1}N(0, 1)$ when $\alpha^0 = 1$, $\sqrt{T}(\hat{\alpha}_{PH} - \alpha^0) \Rightarrow N(0, 4)$. The FD is thus more efficient at $\alpha^0 = 1$. Table 5 shows the properties of the FD, HD and PH estimators using data generated as in Table 1 to facilitate comparison with the QD. Overall, in this simple AR(1) setting, the QD appears to dominate the FD and PH, but is comparable to the HD. These simulation results are in line with our theoretical predictions that the FD, HD, and QD are all asymptotically normal and \sqrt{T} consistent.

No estimator is perfect and QD estimation has its drawbacks. As mentioned in the introduction, the price we pay for asymptotic normality is that $\hat{\alpha}_{QD}$ converges at a rate of \sqrt{T} instead of T when there is a unit root. Han et al. (2009) aggregate L stationary moment conditions and showed that by suitable choice of L , uniform asymptotic normality can be achieved at a rate faster than \sqrt{T} . Extension of their result outside of the AR(1) model is, however, not straightforward. In contrast, the QD framework is broadly applicable.

⁸As suggested by a referee, $E(e_t(e_{t-j} - e_{t-k})) = 0$, $1 \leq j < k$ is also a valid moment condition.

8 Concluding Comments

In this paper, we use a quasi-differencing framework to obtain estimators with classical properties even when the underlying data are highly persistent. Quasi-differencing can render non-stationary processes stationary so that classical limit theorems can be applied. However, the QD estimation approach yields \sqrt{T} consistent rather than super-consistent in the local-to-unity framework. In exchange for this slower convergence is generality, as QD estimation can be used in a broad range of linear and non-linear models.

Appendix A Proofs

The proofs proceed with the assumption that the weighting matrix W_T is an identity matrix.

Proof of Proposition 1 (i): First, consider the problem of matching the j -th autocovariance. That is, $Q_j(\alpha) = (\hat{\gamma}_j(\alpha) - \gamma_j(\alpha))^2$, and $\hat{\alpha}_j = \arg \min_{\alpha} Q_j(\alpha)$. Under the assumption that α is the true value, $\gamma_0(\alpha) = \sigma^2$, and $\gamma_j(\alpha) = 0$ for all $j > 0$. Note that

$$\hat{\gamma}_j(\alpha) - \gamma_j(\alpha) = \frac{1}{T} \sum_{t=j+1}^T \varepsilon_t \varepsilon_{t-j} - \gamma_j(\alpha) - \frac{\alpha - \alpha^0}{T} \sum_{t=j+1}^T [\varepsilon_t y_{t-j-1} + \varepsilon_{t-j} y_{t-1}] + \frac{(\alpha - \alpha^0)^2}{T} \sum_{t=j+1}^T y_{t-1} y_{t-j-1}.$$

As a result, the NQD objective function is the fourth-order polynomial:

$$Q_j(\alpha) = Q_j^{(0)} + (\alpha - \alpha^0) Q_j^{(1)} + (\alpha - \alpha^0)^2 Q_j^{(2)} + (\alpha - \alpha^0)^3 Q_j^{(3)} + (\alpha - \alpha^0)^4 Q_j^{(4)}. \quad (\text{A.1})$$

In the local-to-unity framework with $\alpha_0 = 1 + c/T$, the following results hold as $T \rightarrow \infty$:

$$\frac{1}{T} \sum \varepsilon_{t-j} y_{t-1} \Rightarrow \sigma^2 + \sigma^2 \int_0^1 J_c(s) dW(s), \quad (\text{A.2})$$

$$\frac{1}{T} \sum \varepsilon_t y_{t-1-j} \Rightarrow \sigma^2 \int_0^1 J_c(s) dW(s), \quad (\text{A.3})$$

$$\frac{1}{T^2} \sum y_{t-1} y_{t-j-1} \Rightarrow \sigma^2 \int_0^1 J_c^2(s) ds. \quad (\text{A.4})$$

It follows from equations (4) and (A.2) - (A.4) that

$$\begin{aligned} T^{1/2} Q_j^{(1)} &= -2\sqrt{T} \left(\frac{1}{T} \sum_{t=j+1}^T \varepsilon_t \varepsilon_{t-j} - \gamma_j(\alpha) \right) \frac{1}{T} \sum_{t=j+1}^T [\varepsilon_t y_{t-j-1} + \varepsilon_{t-j} y_{t-1}] \\ &\Rightarrow -2\xi_j \left(\sigma^2 + 2\sigma^2 \int_0^1 J_c(s) dW(s) \right); \end{aligned}$$

$$\begin{aligned} T^{-1/2} Q_j^{(2)} &= 2\sqrt{T} \left(\frac{1}{T} \sum_{t=j+1}^T \varepsilon_t \varepsilon_{t-j} - \gamma_j(\alpha) \right) \left(\frac{1}{T^2} \sum_{t=j+1}^T y_{t-1} y_{t-j-1} \right) \\ &\quad + \frac{1}{\sqrt{T}} \left(\frac{1}{T} \sum_{t=j+1}^T [\varepsilon_t y_{t-j-1} + \varepsilon_{t-j} y_{t-1}] \right)^2 \Rightarrow 2\xi_j \sigma^2 \int_0^1 J_c^2(s) ds; \end{aligned}$$

$$\begin{aligned} T^{-1} Q_j^{(3)} &= -2 \left(\frac{1}{T^2} \sum_{t=j+1}^T y_{t-1} y_{t-j-1} \right) \left(\frac{1}{T} \sum_{t=j+1}^T [\varepsilon_t y_{t-j-1} + \varepsilon_{t-j} y_{t-1}] \right) \\ &\Rightarrow -2\sigma^2 \int_0^1 J_c^2(s) ds \left(\sigma^2 + 2\sigma^2 \int_0^1 J_c(s) dW(s) \right); \end{aligned}$$

$$T^{-2}Q_j^{(4)} = \left(\frac{1}{T^2} \sum_1^T y_{t-1}y_{t-2} \right)^2 \Rightarrow \left(\sigma^2 \int_0^1 J_c^2(s)ds \right)^2 > 0.$$

To summarize:

$$Q_j^{(1)} = O_p(T^{-1/2}), \quad Q_j^{(2)} = O_p(T^{1/2}), \quad Q_j^{(3)} = O_p(T^1), \quad Q_j^{(4)} = O_p(T^2). \quad (\text{A.5})$$

It follows that

$$\frac{Q_j(\alpha) - Q_j^{(0)}}{T^2} \Rightarrow \left(\sigma^2 \int_0^1 J_c^2(s)ds \right)^2 (\alpha - \alpha^0)^4$$

uniformly over a bounded parameter space for α . As a result, $\hat{\alpha}_j$ is a consistent estimate of α^0 .

To study the large sample properties of $\hat{\alpha}_j$, consider the first order condition:

$$Q_j^{(1)} + 2(\hat{\alpha}_j - \alpha^0)Q_j^{(2)} + 3(\hat{\alpha}_j - \alpha^0)^2Q_j^{(3)} + 4(\hat{\alpha}_j - \alpha^0)^3Q_j^{(4)} = 0. \quad (\text{A.6})$$

This is a cubic equation of the form $ax^3 + bx^2 + cx + d = 0$, where x stands for $(\hat{\alpha}_j - \alpha^0)$ with the obvious correspondence between the coefficients. The cubic equation may have one or three real roots depending on the sign of the determinant:

$$\Delta = 18abcd - 4b^3d + b^2c^2 - 4ac^3 - 27a^2d^2.$$

Given the orders established in A.5, it can be shown that

$$T^{-7/2}\Delta \Rightarrow -32\sigma^{10} \left(\int_0^1 J_c^2(s)ds \right)^5 \xi_j^3.$$

The sign of the determinant Δ is asymptotically defined by the sign of ξ_j . When the sign of Δ is negative, there is a unique real root to equation (A.6); otherwise, there are three real roots. However, in the case of three real roots, the middle one corresponds to the local maxima, while the other two roots are the local minima of (A.1).

The next step is to work out the formulas for the roots and to check their rates of convergence toward zero. For example, when there is only one real root, the formula is:

$$x_1 = -\frac{1}{3a} \left(b + \sqrt[3]{b^3 - \frac{9}{2}abc + \frac{27}{2}a^2d + \frac{1}{2}\sqrt{-27a^2\Delta}} + \sqrt[3]{b^3 - \frac{9}{2}abc + \frac{27}{2}a^2d - \frac{1}{2}\sqrt{-27a^2\Delta}} \right).$$

Using the asymptotic orders of the terms in (A.5) and after tedious algebra, we can deduce that $Tx_1 = O_p(1)$ (that is $T(\hat{\alpha}_j - \alpha^0) = O_p(1)$). Similarly, using the explicit formula for cubic roots and denoting the two non-central roots by x_2 and x_3 , we can deduce that when $\Delta > 0$, $T^{3/4}x_2 = O_p(1)$ and $T^{3/4}x_3 = O_p(1)$ (that is, $T^{3/4}(\hat{\alpha}_j - \alpha^0) = O_p(1)$ in this case).

To find the asymptotic distribution of these roots, we start with the case of one root. Because $T(\hat{\alpha}_j - \alpha^0) = O_p(1)$, some terms in equation (A.6) are asymptotically negligible. Thus, asymptotically we have $Q_j^{(1)} + 2(\hat{\alpha}_j - \alpha^0)Q_j^{(2)} = 0$. Equivalently,

$$T(\hat{\alpha}_j - \alpha^0) = -T \frac{Q_j^{(1)}}{2Q_j^{(2)}} \Rightarrow \frac{\sigma^2 + 2\sigma^2 \int_0^1 J_c(s)dw(s)}{2\sigma^2 \int_0^1 J_c^2(s)ds}.$$

Similarly, for the case of two local maxima $T^{3/4}(\hat{\alpha}_j - \alpha^0) = O_p(1)$ and asymptotically $2(\hat{\alpha}_j - \alpha^0)Q_j^{(2)} + 4(\hat{\alpha}_j - \alpha^0)^3Q_j^{(4)} = 0$. Equivalently,

$$T^{3/2}(\hat{\alpha}_j - \alpha^0)^2 = -T^{3/2} \frac{Q_j^{(2)}}{2Q_j^{(4)}} \Rightarrow \frac{-\xi_j}{\sigma^2 \int_0^1 J_c^2(s)ds}.$$

The equation has a solution only when $\xi_j < 0$. As shown above, this is the condition for the cubic equation to have three roots. Thus, we proved that equation (6) holds for $\hat{\alpha}_j$ with $J = 1$. Analogous arguments hold in the general case $\hat{\alpha}_{J,NQD} = \arg \min_{\alpha} \sum_{j=1}^J Q_j(\alpha)$ with ξ_j replaced by $\sum_{j=1}^J \xi_j/J \sim N(0, \sigma^4/J)$. This also shows that in AR(1) case, matching more than one autocovariance leads to increase in efficiency. \square

Proposition 1 (ii) and Proposition 2 are special cases of Proposition 3. Observe that

$$\bar{g}_{j,FQD}(\beta) = A_{j,FQD} + (\beta^0 - \beta)' B_{j,FQD} + (\beta^0 - \beta)' C_{j,FQD}(\beta^0 - \beta),$$

where $A_{j,FQD}$, $B_{j,FQD}$ and $C_{j,FQD}$ are defined in equations (11)-(13). The following three lemmas will be used to prove Proposition 3.

Lemma A-1 (Uniform Law of Large Numbers) *Let $\varepsilon_t = (\varepsilon_{t,1}, \varepsilon_{t,2})'$ be martingale-difference sequence with $\Omega = E(\varepsilon_t \varepsilon_t' | \mathcal{F}_{t-1})$ and finite fourth moments, $\Omega = (\sigma_{i,j})$ and $\eta_{t,i} = \sum_{j=0}^{\infty} c_j^i \varepsilon_{t-j,i}$ for $i = 1, 2$. Uniformly over the set of all sequences c_j^i satisfying $\sum_{j=0}^{\infty} |c_j^i| < C$ for some constant C ,*

$$\frac{1}{T} \sum_{t=1}^T \eta_{t,1} \eta_{t,2} \xrightarrow{p} E[\eta_{t,1} \eta_{t,2}].$$

Proof of Lemma A-1 Notice first that

$$\gamma_j^{i_1, i_2} = \text{cov}(\eta_{t, i_1}, \eta_{t+j, i_2}) = \sigma_{i_1, i_2} \sum_{n=0}^{\infty} c_{n+j}^{i_1} c_n^{i_2}$$

and for any i_1 and i_2 , $\sum_{j=0}^{\infty} |\gamma_j^{i_1, i_2}| < \|\Omega\|C^2$. Furthermore,

$$E[\eta_{t,1} \eta_{t,2} \eta_{t+j,1} \eta_{t+j,2}] = (\gamma_0^{1,2})^2 + \gamma_j^{1,2} \gamma_j^{2,1} + \gamma_j^{1,1} \gamma_j^{2,2} + E(\varepsilon_{1,t}^2 \varepsilon_{2,t}^2) \sum_{n=0}^{\infty} c_n^1 c_{n+j}^1 c_n^2 c_{n+j}^2.$$

As a result,

$$\text{cov}(\eta_{t,1}\eta_{t,2}, \eta_{t+j,1}\eta_{t+j,2}) = \gamma_j^{1,2}\gamma_j^{2,1} + \gamma_j^{1,1}\gamma_j^{2,2} + E(\varepsilon_{1,t}^2\varepsilon_{2,t}^2) \sum_{n=0}^{\infty} c_n^1 c_{n+j}^1 c_n^2 c_{n+j}^2$$

and

$$\sum_{j=1}^{\infty} \text{cov}((\eta_{t,1}\eta_{t,2}), (\eta_{t+j,1}\eta_{t+j,2})) \leq 2\|\Omega\|^2 C^4 + E(\varepsilon_{t,1}^2\varepsilon_{t,2}^2) \left(\sum_{n=0}^{\infty} |c_n^1 c_n^2| \right)^2 \leq (2\|\Omega\|^2 + E(\varepsilon_{t,1}^2\varepsilon_{t,2}^2)) C^4$$

Chebyshev's inequality implies the statement of the Lemma. \square

Lemma A-2 *The following three statements hold asymptotically uniformly over \mathfrak{R}_δ and uniformly over $1 \leq j \leq J$*

(a) $\sqrt{T}(A_{1,FQD}, \dots, A_{J,FQD})' \Rightarrow (\xi_1 - \xi_0, \dots, \xi_J - \xi_0)$, where $(\xi_0, \xi_1, \dots, \xi_J)'$ is a normally distributed random vector with mean zero and diagonal covariance matrix, $E\xi_0^2 = \mu_4$, $E\xi_j^2 = \sigma^4$ for all $1 \leq j \leq J$;

(b) $B_{j,FQD} \xrightarrow{p} a_j = E[(X_{t+j} + X_{t-j} - 2X_t)\varepsilon_t]$;

(c) $C_{j,FQD} = O_p(1)$.

Proof of Lemma A-2: Part (a) follows from applying the Central Limit Theorem (4) to the sums of $\varepsilon_t^2 - \sigma^2$ and $\varepsilon_t\varepsilon_{t-j}$ for $1 \leq j \leq J$. For (b) we need to show that the Uniform Law of Large Numbers holds for

$$B_{j,FQD} = \frac{1}{T} \sum_{t=p+j+1}^T \left(X_t\varepsilon_{t-j} + X_{t-j}\varepsilon_t - 2X_t\varepsilon_t \right) = \frac{1}{T} \sum_{t=p+j+1}^{T-j} \left(X_{t+j} + X_{t-j} - 2X_t \right) \varepsilon_t + O_p(1/\sqrt{T}),$$

where the $O_p(1/\sqrt{T})$ term appears due to the change of limits of summation by a finite number of summands. To apply Lemma A-1, we need to show that the process $X_{t+j} + X_{t-j} - 2X_t$ has absolutely summable MA coefficients. From Lemma S8 in the web Appendix of Mikusheva (2007b), the process $Z_t = X_t - X_{t-1}$ has absolutely summable MA coefficients uniformly over \mathfrak{R}_δ . Now

$$X_{t+j} + X_{t-j} - 2X_t = \sum_{k=1}^j Z_{t+k} - \sum_{k=0}^{j-1} Z_{t-k}.$$

Our process of interest is the sum of a finite number of processes each with summable MA coefficients and thus its MA coefficients are absolutely summable. Lemma A-1 implies that uniformly over \mathfrak{R}_δ

$$B_{j,FQD} \xrightarrow{p} E \left[\left(X_{t+j} + X_{t-j} - 2X_t \right) \varepsilon_t \right] = a_j.$$

Turning to (c), the object of interest is the $p \times p$ matrix:

$$C_{j,FQD} = \frac{1}{2T} \sum_{t=j+p+1}^T (X_t X'_{t-j} + X_{t-j} X'_t - 2X_t X'_t), \quad (\text{A.7})$$

where all elements except possibly the top-left element satisfy the uniform Law of Large Numbers, and thus are of order $O_p(1)$. Now the last $p-1$ elements of X_t are $\Delta y_{t-1}, \dots, \Delta y_{t-p+1}$. From Lemma S8 in Mikusheva (2007b), they have absolutely summable MA coefficients uniformly over \mathfrak{R}_δ . Thus, the elements of the right-bottom $(p-1) \times (p-1)$ sub-matrix satisfy conditions of Lemma A-1. The elements in the first row and the first column (except the top-left element) are of the form

$$\frac{1}{2T} \sum_{t=j+p+1}^T (y_{t-1} z_{t-j} + y_{t-1-j} z_t - 2y_{t-1} z_t) = \frac{1}{2T} \sum_{t=j+1}^T (y_{t-1+j} + y_{t-1-j} - 2y_{t-1}) z_t + O_p(1/\sqrt{T}),$$

where z_t is one of $\Delta y_{t-1}, \dots, \Delta y_{t-p+1}$. From the proof of (b), the series $y_{t-1+j} + y_{t-1-j} - 2y_{t-1}$ also has absolutely summable MA coefficients. Thus, the conditions of Lemma A-1 are satisfied.

It remains to consider the top-left element of the matrix $C_{j,FQD}$ which is given by

$$(C_{j,FQD})_{11} = \frac{1}{T} \sum_{t=j+p+1}^T [y_{t-1} y_{t-j-1} - y_{t-1}^2].$$

If the largest (in absolute value) root λ_p is not real, then by definition of \mathfrak{R}_δ , it is less than $\delta < 1$ in absolute value, and the process y_t is uniformly stationary. Thus $(C_{j,FQD})_{11}$ satisfies the conditions of Lemma A-1. Assume now that the largest root λ_p is a real number. We have $1 - \alpha L - \sum_{j=1}^{p-1} b_j L^j (1 - L) = (1 - \lambda_p) B(L)$, where all inverse roots of $B(L)$ are strictly inside the circle of radius δ .

Let $u_t = y_t - \lambda_p y_{t-1}$ and thus $B(L)u_t = \varepsilon_t$. Now u_t has absolutely summable MA coefficients uniformly over \mathfrak{R}_δ .

$$\begin{aligned} \frac{1}{T} \sum_{t=j+1}^T y_{t-1} y_{t-j-1} &= \frac{1}{T} \sum_{t=j+1}^T y_{t-j-1} (\lambda_p^j y_{t-j-1} + \sum_{k=0}^{j-1} \lambda_p^k u_{t-k-1}) = \\ &= \lambda_p^j \frac{1}{T} \sum_{t=1}^{T-j} y_{t-1}^2 + \sum_{k=0}^{j-1} \lambda_p^k \frac{1}{T} \sum_{t=j+1}^T y_{t-j-1} u_{t-k-1}. \end{aligned}$$

As a result,

$$(C_{j,FQD})_{11} = -(1 - \lambda_p^j) \frac{1}{T} \sum_{t=1}^T y_{t-1}^2 + \sum_{k=1}^{j-1} \lambda_p^{k-1} \frac{1}{T} \sum_{t=j+1}^T y_{t-j-1} u_{t-k-1} + O_p(T^{-1/2}),$$

again the O_p term appears due to change of summation bounds. First, observe that

$$\begin{aligned} \text{Var} \left(\frac{1}{T} \sum_{t=1}^{T-j} y_t u_{t+k} \right) &= \text{Var} \left(\frac{1}{T} \sum_{t=1}^{T-j} \sum_{s=0}^t \lambda_p^s u_{t-s} u_{t+k} \right) = \\ &= \frac{1}{T^2} \sum_{t=1}^{T-j} \sum_{s=0}^t \lambda_p^s \text{cov}(u_{t+k}, u_{t-s}) < \text{Var}(u_t) < \text{const}(\delta). \end{aligned}$$

The variance of u_t is uniformly bounded because all roots of this process are uniformly separated from the unit circle. That is, $\frac{1}{T} \sum_{t=1}^{T-j} y_t u_{t+k} = O_p(1)$ uniformly over $\beta^0 \in \mathfrak{R}_\delta$ and for all $1 \leq k \leq j \leq J$.

Next, consider the term $(1 - \lambda_p^j) \frac{1}{T} \sum_{t=1}^T y_{t-1}^2$. From Theorem 1 in Mikusheva (2010), $(1 - \lambda_p) \frac{1}{T} \sum_{t=1}^T y_{t-1}^2$ is uniformly approximated by $\frac{\sigma^2}{g(c)} \int_0^1 J_c^2(t) dt$, where $g(c) = E \int_0^1 J_c^2(t) dt$, where $J_c(t)$ is the Ornstein-Uhlenbeck process, and $c = T \log(|\lambda_p|)$. It follows from Lemma 4(h) and Lemma 10 in Mikusheva (2007a) that $\frac{1}{g(c)} \int_0^1 J_c^2(t) dt$ is uniformly bounded in probability over c . Summing up, $C_{j,FQD}$ is asymptotically uniformly $O_p(1)$ over \mathfrak{R}_δ and the proof of Lemma A-2 is complete. \square

Lemma A-3 *Under assumptions of Proposition 3 the estimator $\widehat{\beta}_{J,FQD}$ is consistent for any $J > p$.*

Proof of Lemma A-3 Let $f(x) = (f_1(x), \dots, f_{p+1}(x))$, where $f_j(x) = x' B_{j,FQD} + x' C_{j,FQD} x$ and $Q(x) = \sum_{j=1}^J f_j^2(x)$. Any $J \geq p + 1$ suffices for consistency of $\widehat{\beta}_{J,FQD}$, though additional moments may improve efficiency. For any bounded set C in the parameter space, and by Lemma A-2, it holds that:

$$\sup_{\beta \in C} \left| Q(\beta^0 - \beta) - \sum_{j=1}^J (\bar{g}_{j,FQD}(\beta))^2 \right| = o_p(1).$$

Since $Q(0) = 0$, for consistency of $\widehat{\beta}_{J,FQD}$, it is enough to show that for any $\varsigma > 0$, there is $\varepsilon > 0$ such that

$$\lim_{T \rightarrow \infty} P(\inf_{|x| > \varsigma} Q(x) > \varepsilon) = 1, \tag{A.8}$$

where $x = \beta^0 - \beta$. Since $\beta^0 \in \mathfrak{R}_\delta$ and β belongs to bounded neighborhood of \mathfrak{R}_δ , x is bounded. There are two cases to consider: $|\lambda_p| < \delta_1 < 1$, and $\lambda_p \geq \delta_1$.

Case (i) $|\lambda_p| < \delta_1$: We will show that for *any* fixed $0 < \delta_1 < 1$ statement (A.8) holds uniformly over $\beta^0 \in \mathfrak{R}_\delta \cap \{|\lambda_p| < \delta_1\}$.

Since $J > p$, $Q(x) \geq f(x)'f(x)$. For any orthonormal transformation A , $Q(x) \geq (Af(x))'(Af(x)) \geq (Af(x))_1^2$, where $(Af)_1$ is the first component of vector $Af(x)$. Consider a linear transformation, the first component of which is

$$(Af(x))_1 = (f_{p+1} - \alpha^0 f_p - \sum_{j=1}^{p-1} b_j^0 (f_{p+1-j} - f_{p-j})) \frac{1}{a}$$

where $a = \sqrt{1 + (\alpha^0 + b_1^0)^2 + (b_2^0 - b_1^0)^2 + \dots}$ is a (non-zero) multiplier that normalizes the linear transformation. Let $\mathcal{A}(L) = 1 - \alpha^0 L - \sum_{j=1}^{p-1} b_j^0 L^j (1 - L)$ be a lag operator. Given the definition of f_j and linearity of the transformation,

$$(Af(x))_1 = \frac{1}{a} \mathcal{A}(L) f_{p+1} = \frac{1}{a} (x'(\mathcal{A}(L)B_{p+1,FQD}) + x'(\mathcal{A}(L)C_{p+1,FQD})x). \quad (\text{A.9})$$

From (b) of Lemma A-2, $B_{j,FQD} \xrightarrow{p} a_j = E[X_{t+j}\varepsilon_t]$. From (10) and the definition of X_t ,

$$\mathcal{A}(L)X_{t+p+1} = [\varepsilon_{t+p+1}, \Delta\varepsilon_{t+p+1}, \dots, \Delta\varepsilon_{t+2}]' = \tilde{e}_{t+p+1}. \quad (\text{A.10})$$

Since $E\tilde{e}_{t+p+1}\varepsilon_s = 0$ for any $s \leq t$,

$$\mathcal{A}(L)B_{p+1,FQD} \xrightarrow{p} \mathcal{A}(L)E[X_{t+j}\varepsilon_t] = E[\mathcal{A}(L)X_{t+j}\varepsilon_t] = E[\tilde{e}_{t+p+1}\varepsilon_t] = 0.$$

Thus, uniformly over all x in a bounded set,

$$(Af(x))_1 = \frac{1}{a} (x'(\mathcal{A}(L)C_{p+1,FQD})x) + o_p(1). \quad (\text{A.11})$$

It follows from (13) and (A.10) that

$$\mathcal{A}(L)C_{p+1,FQD} = \frac{1}{2T} \sum_{t=p+1}^T (\tilde{e}_{t+p+1}X_t' + X_t\tilde{e}_{t+p+1}') - \mathcal{A}(1)\frac{1}{T} \sum_{t=p+1}^T X_tX_t' + o_p(1), \quad (\text{A.12})$$

where $o_p(1)$ appears from change in the bounds of summation. Since $|\lambda_p| < \delta_1$ by assumption, the process is stationary. Thus $\frac{1}{2T} \sum_{t=p+1}^T \tilde{e}_{t+p+1}X_t' \xrightarrow{p} 0$ uniformly over $|\lambda_p| < \delta_1$ and $\mathcal{A}(1)\frac{1}{T} \sum_{t=p+1}^T X_tX_t'$ is uniformly positive definite. This gives us the needed bound in (A.8) for processes with $|\lambda_p| < \delta_1$.

Case (ii) $|\lambda_p| \geq \delta_1$: To show that for $\delta_1 < 1$ close enough to the unity, (A.8) holds uniformly over $\beta^0 \in \mathfrak{R}_\delta \cap \{|\lambda_p| \geq \delta_1\}$, we divide the area $|x| > \varsigma$ from (A.8) into two regions: $I_1 = \{x : |x| > \varsigma, |x_1| > \varsigma_1\}$ and $I_2 = \{x : |x| > \varsigma, |x_1| \leq \varsigma_1\}$, where $0 < \varsigma_1 < \varsigma$.

Consider $x \in I_1$. We will prove that for *any* fixed $\varsigma_1 > 0$, one can choose δ_1 close enough to the unity such that uniformly over $\beta^0 \in \mathfrak{R}_\delta \cap \{|\lambda_p| > \delta_1\}$, an analog of (A.8) holds where the infimum is taken over $x \in I_1$.

Applying the arguments and transformation as in (A.9), it can be shown that equations (A.11) and (A.12) hold. Since $\mathcal{A}(1)$ converges to zero as δ_1 converges to 1, one can choose δ_1 close enough to 1 to make all terms except the (1,1)-th element of $\mathcal{A}(1)\frac{1}{T}\sum_{t=p+1}^T X_t X_t'$ sufficiently small, and all but the (1,1)-th element of $\frac{1}{T}\sum_{t=p+1}^T \tilde{e}_{t+p+1} X_t'$ converge in probability to its expected value of zero. In consequence, the following holds uniformly over $\beta^0 \in \mathfrak{R}_\delta \cap \{|\lambda_p| > \delta_1\}$ and $x \in I_1$:

$$x' \mathcal{A}(L) C_{p+1, FQD} x = x_1^2 \left(\frac{1}{T} \sum_{t=p+1}^T \varepsilon_{t+p+1} y_{t-1} - \mathcal{A}(1) \frac{1}{T} \sum_{t=p+1}^T y_{t-1}^2 \right) + o_p(1) + o_p(1 - \delta_1).$$

It remains to show that $\frac{1}{T} \sum_{t=p+1}^T (\varepsilon_{t+p+1} - \mathcal{A}(1)y_{t-1}) y_{t-1}$ satisfies the uniform Law of Large Numbers, and thus converges uniformly to a non-zero constant. To do so, we use the Beveridge-Nelson decomposition (Hamilton (1995), chapter 17) that $\varepsilon_{t+p+1} - \mathcal{A}(1)y_{t-1} = u_t - u_{t-1}$, where u_t is a series with absolutely summable MA coefficients. Since $\frac{1}{T}(u_t - u_{t-1})y_{t-1} = -\frac{1}{T}(y_t - y_{t-1})u_t + O_p(1/\sqrt{T})$, Lemma A-1 applies, and $\frac{1}{T} \sum_{t=p+1}^T (\varepsilon_{t+p+1} - \mathcal{A}(1)y_{t-1}) y_{t-1}$ converges in probability to its expectation. Since $\frac{\mathcal{A}(1)}{T} E \sum_{t=p+1}^T y_{t-1}^2$ is uniformly different from zero, this implies that for any fixed $\varsigma_1 > 0$ there exists $\delta_1 < 1$ such that uniformly over $|\lambda_p| > \delta_1$ an analog of (A.8) holds where the infimum is taken over $x \in I_1$.

Consider now $x \in I_2$. One can choose ς_1 small enough and δ_1 close enough to the unity such that uniformly over $\beta^0 \in \mathfrak{R}_\delta \cap \{|\lambda_p| > \delta_1\}$, an analog of (A.8) holds, where the infimum is taken over $x \in I_2$. Given that $B_{j, FQD}$ and $C_{j, FQD}$ are uniformly bounded,

$$f_j(x) = x'_{-1} B_{j, -1} + x'_{-1} C_{j, -1} x_{-1} + o_p(\varsigma_1).$$

where $x_{-1} = (x_2, \dots, x_p)$ is $(p-1) \times 1$ sub-vector of x , and $B_{j, -1}$ and $C_{j, -1}$ are the $(p-1) \times 1$ and $(p-1) \times (p-1)$ sub-matrices of $B_{j, FQD}$ and $C_{j, FQD}$ corresponding to the last $p-1$ components of β .

Let $Z_t = (\Delta y_{t-1}, \dots, \Delta y_{t-p+1})$ and $\tilde{Z}_t = (y_{t-1} - \lambda_p y_{t-2}, \dots, y_{t-p+1} - \lambda_p y_{t-p})'$ be two $(p-1) \times 1$ uniformly stationary vector-processes. Note that the matrices $B_{j, -1}$ and $C_{j, -1}$ satisfy equations analogous to (12) and (13) with Z_t in place of X_t . Similarly, \tilde{B}_j and \tilde{C}_j are defined as in (12) and (13) with \tilde{Z}_t in place of X_t . Observe that $Z_t' = \tilde{Z}_t' - (1 - \lambda_p)(y_{t-2}, \dots, y_{t-p})$. It is easy to see that

$$f_j(x) = x'_{-1} \tilde{B}_j + x'_{-1} \tilde{C}_j x_{-1} + o_p(1 - \delta_1) + o_p(\varsigma_1).$$

The function $\tilde{f}_j = x'_{-1} \tilde{B}_j + x'_{-1} \tilde{C}_j x_{-1}$ corresponds to that of the uniformly stationary process $y_t - \lambda_p y_{t-1}$ with all roots smaller than δ in absolute value. The rest of the proof follows arguments as in Case i. \square

Proof of Proposition 3 (i): To establish the asymptotic distribution of $\widehat{\beta}_{J,FQD}$, consider first order condition:

$$\sum_{j=1}^J \bar{g}_{j,FQD}(\widehat{\beta}_{J,FQD}) \frac{\partial \bar{g}_{j,FQD}}{\partial \beta}(\widehat{\beta}_{J,FQD}) = 0.$$

From Lemma A-2 and consistency of $\widehat{\beta}_{J,FQD}$,

$$\frac{\partial \bar{g}_{j,FQD}}{\partial \beta}(\widehat{\beta}_{J,FQD}) = -B_{j,FQD} + o_p(1) \rightarrow^p a_j,$$

and uniformly over \mathfrak{R}_δ :

$$\sqrt{T} \bar{g}_{j,FQD}(\widehat{\beta}_{J,FQD}) = \sqrt{T} A_{j,FQD} + a'_j \sqrt{T} (\widehat{\beta}_{J,FQD} - \beta^0) + o_p(1).$$

As a result, the following holds uniformly:

$$\sqrt{T} (\widehat{\beta}_{J,FQD} - \beta^0) \Rightarrow \left(\sum_{j=1}^J a_j a'_j \right)^{-1} \left(\sum_{j=1}^J a_j (\xi_j - \xi_0) \right) = N(0, \Sigma_{J,FQD}),$$

where $G = \left(\sum_{j=1}^J a_j a'_j \right)^{-1}$, and $\Sigma_{J,FQD} = \sigma^4 G + \mu_4 G \left(\sum_{j=1}^J a_j \right) \left(\sum_{j=1}^J a_j \right)' G$.

Proof of Proposition 3 (ii): The proof proceeds by treating QD as a two-step estimator. First, note that

$$s^2 - \frac{1}{T} \sum_{t=p+1}^T \varepsilon_t^2 = -\frac{1}{T} \left(\sum_t X_t \varepsilon_t \right)' \left(\sum_t X_t X_t' \right)^{-1} \left(\sum_t X_t \varepsilon_t \right)$$

Theorem 1 in Mikusheva (2010) shows that the statistic $\left(\sum_t X_t \varepsilon_t \right)' \left(\sum_t X_t X_t' \right)^{-1} \left(\sum_t X_t \varepsilon_t \right)$ is uniformly approximated by the distribution $(t^c + N(0, p-1))^2$, where $t^c = \frac{\int J_c(t) dw(t)}{\sqrt{\int J_c^2(t) dt}}$ is a local-to-unity limit of a t-statistic, and $c = T \log(|\lambda_p|)$. Given that t^c is uniformly bounded in probability over all possible values of $c \leq 0$, the following holds uniformly over \mathfrak{R}_δ :

$$s^2 = \frac{1}{T} \sum_t \varepsilon_t^2 + O_p(1/T). \quad (\text{A.13})$$

Since $\bar{g}_{j,QD}(\beta) = \bar{g}_{j,FQD}(\beta) - \gamma_0 + s^2$,

$$\bar{g}_{j,QD}(\beta) = A_{j,QD} + (\beta^0 - \beta)' B_{j,QD} + (\beta^0 - \beta)' C_{j,QD} (\beta^0 - \beta),$$

where $A_{j,QD} = A_{j,FQD} + s^2 - \sigma^2$, $B_{j,QD} = B_{j,FQD}$ and $C_{j,QD} = C_{j,FQD}$. A result analogous of Lemma A-2 holds for $A_{j,QD}$, $B_{j,QD}$ and $C_{j,QD}$ with one correction: $\sqrt{T}(A_{1,QD}, \dots, A_{J,QD}) \Rightarrow (\xi_1, \dots, \xi_J)$. This gives us consistency and asymptotic normality of $\widehat{\beta}_{J,QD}$ with asymptotic covariance matrix $\Sigma_{J,QD} = \sigma^4 G$. \square

The following lemma will be used to prove Proposition 4.

Lemma A-4 *Uniformly over all possible values of θ ,*

- (a) $\sqrt{T}(A_j + \Omega^0) \Rightarrow \xi_j - \xi_0$;
- (b) $\sqrt{T}(A_j + S) \Rightarrow \xi_j$;
- (c) $B_{j,1} \xrightarrow{p} E[\varepsilon_t(x_{t-j-1} - x_{t-1})] = 0$;
- (d) $B_{j,2} \xrightarrow{p} E[x_{t-1}(\varepsilon_{t-j} - \varepsilon_t)'] = a_j$;
- (e) $C_j = O_p(1)$.

where $\frac{1}{\sqrt{T}} \sum_{t=j+1}^T \varepsilon_t \varepsilon_t' \Rightarrow \xi_j$ and ξ_j is a 2×2 matrix with normally distributed components such that for any non-random vector \mathbf{a} the vector $\xi_j \mathbf{a}$ is normally distributed with variance-covariance matrix $\Omega^0 \mathbf{a}' \Omega^0 \mathbf{a}$. We also have $\frac{1}{\sqrt{T}} \sum_{t=j+1}^T \varepsilon_t \varepsilon_t' \Rightarrow \xi_0$ where ξ_0 is a 2×2 matrix with normally distributed components such that for any non-random vector \mathbf{a} , the vector $\xi_0 \mathbf{a}$ is normally distributed with variance-covariance matrix $E[(\varepsilon_t \mathbf{a})^2 \varepsilon_t \varepsilon_t']$. The variables ξ_j are independent for any $j \geq 0$.

Proof of Lemma A-4 Result (a) follows from Central Limit Theorem. To prove (b), note that

$$S = \frac{1}{T} \sum_{t=1}^T \varepsilon_t \varepsilon_t' - \frac{1}{T} \frac{1}{\sum_{s=1}^T x_{s-1}^2} \begin{pmatrix} (\sum_s \varepsilon_{xs} x_{s-1})^2 & (\sum_s \varepsilon_{xs} x_{s-1})(\sum_s \varepsilon_{ys} x_{s-1}) \\ (\sum_s \varepsilon_{ys} x_{s-1})(\sum_s \varepsilon_{xs} x_{s-1}) & (\sum_s \varepsilon_{ys} x_{s-1})^2 \end{pmatrix}$$

Now $\frac{\sum_{s=1}^T \varepsilon_{xs} x_{s-1}}{\sqrt{\sum_{s=1}^T x_{s-1}^2}} \Rightarrow t^c$ uniformly over $\alpha^0 \in (-1 + \delta, 1]$, and the family t^c is uniformly bounded. The sum $\frac{\sum_{s=1}^T \varepsilon_{ys} x_{s-1}}{\sqrt{\sum_{s=1}^T E x_{s-1}^2}}$ has a bounded second moment since $\varepsilon_{ys} x_{s-1}$ is martingale-difference sequence,

and thus it is uniformly bounded by Chebyshev's inequality. Lastly, $\frac{\sum_{s=1}^T x_{s-1}^2}{\sum_{s=1}^T E x_{s-1}^2}$ is uniformly separated from zero, a result that follows from Lemma 4(h) and Lemma 10 in Mikusheva (2007a). Summing up, we have $S = \frac{1}{T} \sum_{t=1}^T \varepsilon_t \varepsilon_t' + O_p(\frac{1}{T})$. As a result, $\sqrt{T}(A_j + S) = \frac{1}{\sqrt{T}} \sum_{t=j+1}^T \varepsilon_t \varepsilon_t' \Rightarrow \xi_j$.

The proof of part (c) follows from Lemma A-1, since we show in the proof of Proposition 3 that $x_{t-j-1} - x_{t-1}$ has absolutely summable MA coefficients uniformly over α .

To prove (d), re-write

$$B_{j,2} = \frac{1}{T} \sum_{t=j+1}^T x_{t-1}(\varepsilon_{t-j} - \varepsilon_t)' = \frac{1}{T} \sum_{t=j+1}^T (x_{t+j-1} - x_{t-1})\varepsilon_t' + \frac{1}{T} \sum_{t=j+1}^{2j} x_{t-1}\varepsilon_{t-j} - \frac{1}{T} \sum_{t=T+1}^{T+j} x_{t-1}\varepsilon_{t-j}.$$

The terms $\frac{1}{T} \sum_{t=j+1}^{2j} x_{t-1}\varepsilon_{t-j}$ and $\frac{1}{T} \sum_{t=T+1}^{T+j} x_{t-1}\varepsilon_{t-j}$ both have j summands each of which are of order $O_p(\sqrt{T})$. This means that for any $j \leq J$ where J is fixed, the following holds uniformly:

$$B_{j,2} = \frac{1}{T} \sum_{t=j+1}^T (x_{t+j-1} - x_{t-1})\varepsilon_t' + O_p\left(\frac{1}{\sqrt{T}}\right).$$

The rest of the proof is the same as for part (c). Part (e) follows from Proposition 3.(c). \square

Proof of Proposition 4: Note first that

$$\begin{aligned}
\widehat{\Gamma}_j(\theta) - \widehat{\Gamma}_0(\theta) &= \frac{1}{T} \sum_{t=j+1}^T e_t(\theta)(e_{t-j}(\theta) - e_t(\theta))' \\
&= \frac{1}{T} \sum_{t=j+1}^T ((\theta^0 - \theta)x_{t-1} + \varepsilon_t)((\theta^0 - \theta)(x_{t-j-1} - x_{t-1}) + \varepsilon_{t-j} - \varepsilon_t)' \\
&= A_j + B_{j,1}(\theta^0 - \theta)' + (\theta^0 - \theta)B_{j,2} + (\theta^0 - \theta)C_j(\theta^0 - \theta)',
\end{aligned}$$

where

$$\begin{aligned}
A_j &= \frac{1}{T} \sum_{t=j+1}^T \varepsilon_t(\varepsilon_{t-j} - \varepsilon_t)'; B_{j,1} = \frac{1}{T} \sum_{t=j+1}^T \varepsilon_t(x_{t-j-1} - x_{t-1}); \\
B_{j,2} &= \frac{1}{T} \sum_{t=j+1}^T x_{t-1}(\varepsilon_{t-j} - \varepsilon_t)'; C_j = \frac{1}{T} \sum_{t=j+1}^T x_{t-1}(x_{t-j-1} - x_{t-1}).
\end{aligned}$$

Lemma A-4 showed that uniformly over α :

$$\|\widehat{\Gamma}_j(\theta) - \widehat{\Gamma}_0(\theta) + \Omega^0\|_2^2 = \|(\theta^0 - \theta)B_{j,2} + C_j(\theta^0 - \theta)(\theta^0 - \theta)'\|_2^2 + o_p(1),$$

and

$$\|\widehat{\Gamma}_j(\theta) - \widehat{\Gamma}_0(\theta) + S\|_2^2 = \|(\theta^0 - \theta)B_{j,2} + C_j(\theta^0 - \theta)(\theta^0 - \theta)'\|_2^2 + o_p(1).$$

We minimize the sum of such functions for $j = 1, \dots, J$. Obviously, the minimized function is non-negative and one of its minimal value of zero is achieved at $\theta = \theta^0$. The question is whether there are any other minima. For this, there should exist θ such that $\|(\theta^0 - \theta)B_{j,2} + C_j(\theta^0 - \theta)(\theta^0 - \theta)'\|_2$ is zero for all j . For a given j , the only non-trivial null of function $\|(\theta^0 - \theta)B_{j,2} + C_j(\theta^0 - \theta)(\theta^0 - \theta)'\|_2$ implies $\theta_j = \theta^0 + \frac{1}{C_j}B_{j,2}$ which is asymptotically different for different j . This implies that for $J \geq 2$ no other asymptotic null of the objective function other than $\theta = \theta^0$ exists, and thus $\widehat{\theta}_{J,FQD}$ and $\widehat{\theta}_{J,QD}$ are consistent.

To derive the limit distribution of $\widehat{\theta}_{J,FQD}$, we use the fact that the first order condition must be satisfied at $\theta = \widehat{\theta}_{J,FQD}$. Now the first order condition is

$$\begin{aligned}
&\nabla_{\theta} \|\widehat{\Gamma}_j(\theta) - \widehat{\Gamma}_0(\theta) + \Omega^0\|_2^2 \\
&= -2(A_j + \Sigma^0 + B_{j,1}(\theta^0 - \theta)' + (\theta^0 - \theta)B_{j,2} + C_j(\theta^0 - \theta)(\theta^0 - \theta)') (B'_{j,1} + B_{j,2} + 2C_j(\theta^0 - \theta)')'.
\end{aligned}$$

Since we proved that $\widehat{\theta}_{J,FQD}$ is uniformly consistent, and given statements (d) and (e),

$$B_{j,2} + 2C_j(\theta^0 - \widehat{\theta}_{J,FQD})' \rightarrow^p a_j.$$

Furthermore,

$$\begin{aligned} & \sqrt{T} \left(A_j + \Omega^0 + B_{j,1}(\theta^0 - \hat{\theta}_{J,FQD}) + (\theta^0 - \hat{\theta}_{J,FQD})B_{j,2} + C_j(\theta^0 - \hat{\theta}_{J,FQD})(\theta^0 - \hat{\theta}_{J,FQD})' \right) \\ &= \sqrt{T}(A_j + \Omega^0) + \sqrt{T}(\theta^0 - \hat{\theta}_{J,FQD})a_j + o_p(1). \end{aligned}$$

As a result,

$$\sqrt{T}(\hat{\theta}_{J,FQD} - \theta^0) \Rightarrow \frac{1}{\sum_{j=1}^J a_j a_j'} \sum_{j=1}^J (\xi_j - \xi_0) a_j$$

uniformly over α . Similarly,

$$\sqrt{T}(\hat{\theta}_{J,QD} - \theta^0) \Rightarrow \frac{1}{\sum_{j=1}^J a_j a_j'} \sum_{j=1}^J \xi_j a_j.$$

The last two formulas lead to the conclusion of Lemma 4.

Relation between PH and FD: Observe that

$$\begin{aligned} \sum_{t=2}^T (\Delta y_{t-1})^2 &= \sum_{t=2}^T \Delta y_{t-1} y_{t-1} - \sum_{t=2}^T (y_{t-1} - y_{t-2}) y_{t-2} \\ &= \sum_{t=2}^T \Delta y_{t-1} y_{t-1} + \sum_{t=2}^T y_{t-1} (y_{t-1} - y_{t-2}) - y_{T-1}^2 + y_0^2 = 2 \sum_{t=2}^T \Delta y_{t-1} y_{t-1} - y_{T-1}^2 + y_0^2. \end{aligned}$$

Thus if $|\alpha^0| < 1$ is fixed and $T \rightarrow \infty$,

$$\frac{1}{T} \sum_{t=2}^T (\Delta y_{t-1})^2 = 2 \frac{1}{T} \sum_{t=2}^T \Delta y_{t-1} y_{t-1} + O_p(1/T).$$

Similarly,

$$\begin{aligned} \sum_{t=2}^T \Delta y_{t-1} (2\Delta y_t + \Delta y_{t-1}) &= 2 \sum_{t=2}^T \Delta y_{t-1} y_t - 2 \sum_{t=2}^T \Delta y_{t-1} y_{t-1} + \sum_{t=2}^T (\Delta y_{t-1})^2 = \\ &= 2 \sum_{t=2}^T \Delta y_{t-1} y_t - y_{T-1}^2 + y_0^2 \end{aligned}$$

and

$$\frac{1}{T} \sum_{t=2}^T \Delta y_{t-1} (2\Delta y_t + \Delta y_{t-1}) = 2 \frac{1}{T} \sum_{t=2}^T \Delta y_{t-1} y_t + O_p(1/T).$$

This leads us to the result that $\hat{\alpha}_{PH} = \hat{\alpha}_{FD} + O_p(T^{-1})$ under stationary asymptotics.

References

- Abowd, J. M. and Card, D. 1989, On the Covariance Structure of Earnings and Hours Changes, *Econometrica* **57(2)**, 411–445.
- Altonji, J. and Segal, L. 1996, Small-Sample Bias in GMM estimation of Covariance Structures, *Journal of Business and Economic Statistics* **14(3)**, 353–366.
- Canjels, E. and Watson, M. W. 1997, Estimating Deterministic Trends in the Presence of Serially Correlated Errors, *Review of Economics and Statistics* **May**, 184–200.
- Gorodnichenko, Y. and Ng, S. 2010, Estimation of DSGE Models When the Data are Persistent, *Journal of Monetary Economics* **57**, 325–340.
- Han, C. and Kim, B. 2011, A GMM Interpretation of the Paradox in the Inverse Probability Weighting Estimation of the Average Treatment Effect on the Treated, *Economics Letters* **110**, 163–165.
- Han, C., Phillips, P. and Sul, D. 2009, Uniform Asymptotic Normality in Stationary and Unit Root Autoregression, unpublished manuscript.
- Jansson, M. and Moreira, M. 2006, Optimal Inference In Regression with Nearly Integrated Regressors, *Econometrica* **74**, 681–714.
- Komunjer, I. and Ng, S. 2009, Dynamic Identification of DSGE models, unpublished manuscript, Columbia University.
- Laroque, B. and Salanie, B. 1997, Normal Estimators for Cointegrating Relationships, *Economics Letters* **55**, 185–189.
- Mikusheva, A. 2007a, Uniform Inference in Autoregressive Models, *Econometrica* **75:5**, 1411–1452.
- Mikusheva, A. 2007b, Uniform Inference in Autoregressive Models, *Econometrica* **75:5**, Supplementary Material.
- Mikusheva, A. 2010, One Dimensional Inferences in Autoregressive Models with the Potential Presence of a Unit Root, mimeo, M.I.T.
- Newey, W. and McFadden, D. 1994, Large Sample Estimators and Hypothesis Testing, *Handbook of Econometrics*, Vol. 4, Chapter 36, North Holland.
- Pesavento, E. and Rossi, B. 2006, Small Sample Confidence Interevals for Multivariate Impulse Response Functions at Long Lags, *Journal of Applied Econometrics* **21:8**, 1135–1155.
- Phillips, P. and Lee, C. 1996, Efficiency Gains from Quasi-differencing Under Non-stationarity, in P. Robinson and M. Rosenblatt (eds), *Athens Conference on Applied Probability and Time Series: Volume II Time Series Analysis in Honor of E.J. Hannan*.
- Phillips, P. C. B. and Han, C. 2008, Gaussian Inference in AR(1) Time Series With or Without a Unit Root, *Econometric Theory* **63**, 1023–1078.
- Pierce, D. 1982, The Asymptotic Effect of Substituting Estimators for Parameters in Certain Types of Statistics, *Annals of Statistics* **10:2**, 475–478.
- Prokhorov, A. and Schmidt, P. 2009, GMM Redundancy Results for the Missing Data Problem, *Journal of Econometrics* **151**, 47–55.

- So, B. and Shin, D. 1999, Cauchy Estimators for Autoregressive Processes with Applications to Unit Root Tests and Confidence Intervals, *Econometric Theory* **15**, 166–176.
- Uhlig, H. 1999, A Toolkit for Analyzing Nonlinear Dynamic Stochastic Models Easily, in R. Marimon and A. Scott (eds), *Computational Methods for the Study of Dynamic Economies*, Oxford University Press, pp. 30–61.

Table 1. AR(1) model.

T	α	QD						FQD						OLS					
		mean		t-test		J-test		mean		t-test		J-test		mean		rmse		t-test	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)				
Panel A: Intercept model																			
200	1.00	0.975	0.059	0.107	0.533	0.085	0.991	0.122	0.033	0.098	0.078	0.973	0.035	0.473	1.000				
200	0.98	0.960	0.052	0.102	0.484	0.083	0.971	0.113	0.032	0.085	0.069	0.956	0.035	0.219	1.000				
200	0.95	0.934	0.050	0.080	0.428	0.101	0.942	0.114	0.028	0.066	0.095	0.929	0.036	0.150	0.969				
200	0.90	0.883	0.054	0.098	0.417	0.088	0.888	0.117	0.030	0.084	0.083	0.880	0.040	0.111	0.687				
200	0.80	0.782	0.058	0.091	0.359	0.075	0.794	0.126	0.035	0.081	0.077	0.782	0.050	0.097	0.413				
500	1.00	0.990	0.030	0.066	0.748	0.069	0.997	0.072	0.023	0.136	0.061	0.989	0.014	0.462	1.000				
500	0.98	0.974	0.028	0.064	0.681	0.056	0.977	0.073	0.035	0.150	0.054	0.971	0.015	0.138	1.000				
500	0.95	0.943	0.030	0.076	0.676	0.077	0.946	0.072	0.037	0.154	0.071	0.941	0.019	0.127	1.000				
500	0.90	0.894	0.031	0.073	0.598	0.068	0.896	0.074	0.029	0.144	0.067	0.893	0.022	0.082	0.930				
500	0.80	0.793	0.035	0.085	0.545	0.047	0.792	0.080	0.043	0.146	0.052	0.792	0.030	0.082	0.687				
Panel B: Linear trend model																			
200	1.00	0.964	0.063	0.224	0.641	0.108	0.991	0.112	0.025	0.091	0.101	0.950	0.057	0.775	1.000				
200	0.98	0.954	0.058	0.164	0.547	0.099	0.967	0.114	0.028	0.081	0.098	0.940	0.050	0.454	1.000				
200	0.95	0.926	0.056	0.146	0.511	0.090	0.945	0.111	0.022	0.078	0.077	0.916	0.047	0.278	0.986				
200	0.90	0.876	0.059	0.143	0.481	0.108	0.892	0.119	0.032	0.071	0.105	0.869	0.049	0.180	0.775				
200	0.80	0.775	0.060	0.121	0.432	0.075	0.788	0.127	0.043	0.081	0.078	0.770	0.054	0.133	0.512				
500	1.00	0.985	0.033	0.137	0.775	0.072	0.999	0.069	0.020	0.160	0.063	0.980	0.023	0.777	1.000				
500	0.98	0.972	0.029	0.080	0.717	0.082	0.976	0.071	0.041	0.153	0.081	0.967	0.018	0.245	1.000				
500	0.95	0.942	0.031	0.086	0.678	0.073	0.950	0.074	0.034	0.161	0.068	0.938	0.021	0.185	1.000				
500	0.90	0.891	0.032	0.090	0.654	0.061	0.898	0.076	0.041	0.151	0.068	0.889	0.025	0.123	0.957				
500	0.80	0.792	0.034	0.089	0.567	0.043	0.797	0.078	0.025	0.122	0.054	0.789	0.030	0.090	0.729				

Note: DGP is AR(1) model while AR(2) model is fitted. Three autocovariances are used in QD estimation. T-test and J-test sizes are for 5 percent level. Power of the t-test is computed for the null of $H_0 : \alpha = \alpha_0 - 0.05$. Additional results are available in the online appendix.

Table 2. AR(2) model.

T	α	QD						FQD						OLS					
		mean		t-test		J-test		mean		t-test		J-test		mean		rmse		t-test	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)				
Panel A: Intercept model																			
200	1.00	0.962	0.101	0.073	0.322	0.050	0.983	0.155	0.057	0.097	0.044	0.968	0.042	0.468	1.000				
200	0.98	0.965	0.070	0.059	0.269	0.045	0.975	0.149	0.046	0.074	0.039	0.952	0.041	0.247	0.999				
200	0.95	0.939	0.070	0.062	0.266	0.051	0.947	0.158	0.040	0.073	0.039	0.923	0.044	0.179	0.928				
200	0.90	0.890	0.072	0.064	0.246	0.038	0.900	0.163	0.051	0.081	0.033	0.877	0.048	0.121	0.590				
200	0.80	0.787	0.077	0.074	0.215	0.048	0.795	0.184	0.058	0.094	0.047	0.776	0.059	0.089	0.364				
500	1.00	0.982	0.067	0.045	0.503	0.036	0.987	0.099	0.050	0.121	0.030	0.987	0.017	0.464	1.000				
500	0.98	0.975	0.037	0.046	0.447	0.031	0.977	0.094	0.041	0.119	0.029	0.969	0.017	0.176	1.000				
500	0.95	0.946	0.040	0.056	0.431	0.034	0.951	0.093	0.047	0.117	0.030	0.940	0.022	0.122	0.999				
500	0.90	0.897	0.041	0.051	0.405	0.032	0.895	0.101	0.060	0.138	0.030	0.891	0.025	0.079	0.874				
500	0.80	0.798	0.045	0.059	0.344	0.025	0.798	0.122	0.077	0.155	0.031	0.793	0.032	0.074	0.546				
Panel B: Linear trend model																			
200	1.00	0.962	0.088	0.158	0.423	0.045	0.989	0.152	0.055	0.094	0.045	0.940	0.070	0.762	1.000				
200	0.98	0.949	0.080	0.142	0.395	0.040	0.975	0.153	0.045	0.094	0.039	0.928	0.064	0.493	0.999				
200	0.95	0.928	0.075	0.110	0.343	0.045	0.940	0.155	0.043	0.085	0.041	0.907	0.058	0.301	0.969				
200	0.90	0.881	0.077	0.104	0.306	0.053	0.899	0.162	0.041	0.080	0.045	0.862	0.059	0.210	0.699				
200	0.80	0.775	0.087	0.117	0.300	0.037	0.796	0.186	0.066	0.099	0.037	0.757	0.072	0.158	0.520				
500	1.00	0.984	0.046	0.107	0.566	0.023	0.997	0.093	0.045	0.145	0.024	0.975	0.029	0.760	1.000				
500	0.98	0.972	0.039	0.062	0.482	0.028	0.980	0.092	0.049	0.120	0.031	0.962	0.024	0.317	1.000				
500	0.95	0.942	0.041	0.078	0.484	0.029	0.950	0.087	0.028	0.105	0.030	0.934	0.026	0.184	1.000				
500	0.90	0.891	0.043	0.084	0.447	0.034	0.897	0.104	0.061	0.137	0.033	0.884	0.030	0.145	0.921				
500	0.80	0.793	0.046	0.072	0.395	0.027	0.795	0.116	0.063	0.143	0.026	0.785	0.035	0.097	0.649				

Note: DGP is AR(2) model while AR(3) model is fitted. All results are for the sum of autoregressive coefficients. Four autocovariances are used in QD estimation. T-test and J-test sizes are for 5 percent level. Power of the t-test is computed for the null of $H_0 : \alpha = \alpha_0 - 0.05$. Additional results are available in the online appendix.

Table 3. Predictive regression. (True parameter β is equal to one)

T	α	QD						FQD						OLS					
		mean		t-test		J-test		mean		t-test		J-test		mean		rmse		t-test	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)				
Panel A: Intercept model																			
200	1.00	0.986	0.051	0.054	0.302	0.166	0.999	0.106	0.026	0.087	0.243	0.987	0.022	0.169	0.985				
200	0.98	0.988	0.049	0.042	0.274	0.181	0.996	0.105	0.027	0.097	0.255	0.989	0.025	0.104	0.937				
200	0.95	0.992	0.052	0.044	0.258	0.182	1.001	0.108	0.035	0.099	0.304	0.990	0.030	0.097	0.751				
200	0.90	0.991	0.055	0.057	0.265	0.191	1.004	0.114	0.031	0.092	0.296	0.990	0.036	0.081	0.554				
200	0.80	0.992	0.057	0.047	0.224	0.195	1.001	0.117	0.033	0.101	0.309	0.994	0.045	0.064	0.343				
500	1.00	0.995	0.030	0.034	0.494	0.068	0.999	0.063	0.022	0.147	0.109	0.995	0.009	0.172	1.000				
500	0.98	0.995	0.030	0.032	0.502	0.089	0.999	0.066	0.031	0.181	0.150	0.995	0.012	0.093	1.000				
500	0.95	0.995	0.031	0.042	0.466	0.073	0.999	0.071	0.041	0.172	0.157	0.996	0.016	0.082	0.987				
500	0.90	0.995	0.032	0.045	0.466	0.076	0.996	0.069	0.035	0.173	0.147	0.996	0.021	0.072	0.871				
500	0.80	0.997	0.034	0.051	0.381	0.083	1.002	0.077	0.039	0.147	0.153	0.997	0.028	0.061	0.631				
Panel B: Linear trend model																			
200	1.00	0.978	0.056	0.089	0.374	0.175	0.994	0.108	0.032	0.102	0.270	0.973	0.036	0.310	0.983				
200	0.98	0.985	0.053	0.064	0.303	0.180	1.004	0.109	0.027	0.097	0.288	0.980	0.033	0.180	0.927				
200	0.95	0.983	0.055	0.073	0.316	0.191	0.999	0.110	0.037	0.119	0.302	0.981	0.037	0.151	0.793				
200	0.90	0.984	0.055	0.062	0.292	0.177	1.002	0.115	0.038	0.104	0.299	0.984	0.039	0.109	0.591				
200	0.80	0.984	0.060	0.075	0.279	0.180	0.998	0.119	0.036	0.099	0.302	0.987	0.047	0.084	0.388				
500	1.00	0.992	0.031	0.054	0.543	0.086	0.998	0.068	0.030	0.188	0.139	0.990	0.014	0.278	1.000				
500	0.98	0.993	0.030	0.035	0.522	0.080	1.001	0.067	0.039	0.178	0.149	0.993	0.014	0.148	1.000				
500	0.95	0.993	0.031	0.042	0.514	0.074	0.999	0.066	0.033	0.161	0.146	0.993	0.018	0.100	0.983				
500	0.90	0.995	0.031	0.033	0.479	0.095	1.002	0.071	0.027	0.150	0.154	0.994	0.022	0.080	0.891				
500	0.80	0.994	0.035	0.045	0.448	0.085	1.009	0.077	0.029	0.121	0.158	0.995	0.027	0.062	0.647				

Note: DGP is given by equations (14a) and (14b). All results are for the parameter β . In all simulations $\rho_{ue} = 0.5$. Two autocovariances are used in QD estimation. T-test and J-test sizes are for 5 percent level. Power of the t-test is computed for the null of $H_0 : \beta = \beta_0 - 0.05$. Additional results are available in the online appendix.

Table 4. DSGE model, capital intensity ψ .

T	α	QD					MLE			
		mean	rmse	t-test		J-test	mean	rmse	t-test	
				size	power	size			size	power
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)		
Panel A: Intercept model										
200	1.00	0.255	0.122	0.077	0.245	0.088	0.181	0.117	0.309	0.577
200	0.98	0.255	0.120	0.098	0.259	0.085	0.220	0.094	0.142	0.409
200	0.95	0.268	0.133	0.076	0.238	0.095	0.246	0.094	0.076	0.294
200	0.90	0.284	0.147	0.066	0.201	0.094	0.266	0.114	0.065	0.235
200	0.80	0.289	0.144	0.046	0.148	0.079	0.285	0.131	0.065	0.184
500	1.00	0.260	0.072	0.050	0.304	0.049	0.173	0.110	0.517	0.830
500	0.98	0.261	0.071	0.059	0.322	0.053	0.231	0.058	0.151	0.626
500	0.95	0.262	0.073	0.066	0.331	0.049	0.245	0.054	0.076	0.522
500	0.90	0.266	0.082	0.052	0.306	0.043	0.253	0.062	0.053	0.428
500	0.80	0.281	0.110	0.035	0.197	0.035	0.270	0.096	0.062	0.315
Panel B: Linear trend model										
200	1.00	0.255	0.123	0.088	0.264	0.093	0.223	0.094	0.153	0.399
200	0.98	0.267	0.128	0.075	0.222	0.090	0.243	0.091	0.102	0.315
200	0.95	0.272	0.138	0.085	0.233	0.090	0.257	0.097	0.065	0.249
200	0.90	0.282	0.145	0.066	0.204	0.083	0.273	0.113	0.057	0.205
200	0.80	0.298	0.146	0.053	0.130	0.081	0.296	0.135	0.057	0.150
500	1.00	0.259	0.071	0.050	0.308	0.041	0.209	0.079	0.324	0.730
500	0.98	0.262	0.072	0.057	0.309	0.040	0.237	0.059	0.139	0.593
500	0.95	0.263	0.073	0.058	0.315	0.039	0.250	0.056	0.069	0.477
500	0.90	0.268	0.082	0.042	0.280	0.034	0.259	0.064	0.042	0.387
500	0.80	0.284	0.108	0.024	0.162	0.028	0.277	0.097	0.053	0.295

Note: The true value of capital intensity is $\psi = 0.25$. The observed series is consumption. *QD* uses OLS estimate of the standard deviation of innovations consumption for QD estimation. *MLE* corresponds to the maximum likelihood estimation (Kalman filter) of the structural parameters. Three autocorrelation coefficients (i.e, the fitted model is AR(3)) and six autocovariances are used in QD estimation. T-test and J-test sizes are for 5 percent level. Power of the t-test is computed for the null of $H_0 : \psi = \psi_0 - 0.1$.

Table 5. Linear QD Estimators: AR(1) model.

T	α	HD					FD					PH				
		mean	rmse	t-test	J-test		mean	rmse	t-test	J-test		mean	rmse	t-test	J-test	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)
Panel A: Intercept model																
200	1.00	0.975	0.048	0.156	0.662	0.073	0.954	0.154	0.025	0.189	0.072	1.003	0.171	0.092	0.157	
200	0.98	0.961	0.046	0.132	0.561	0.077	0.958	0.093	0.030	0.145	0.084	0.977	0.180	0.111	0.173	
200	0.95	0.933	0.045	0.114	0.532	0.068	0.931	0.081	0.035	0.170	0.073	0.952	0.179	0.116	0.173	
200	0.90	0.883	0.049	0.122	0.481	0.066	0.881	0.081	0.052	0.179	0.075	0.896	0.174	0.108	0.173	
200	0.80	0.781	0.053	0.099	0.424	0.072	0.779	0.083	0.058	0.190	0.080	0.792	0.170	0.115	0.188	
500	1.00	0.991	0.026	0.081	0.805	0.058	0.977	0.122	0.016	0.313	0.058	1.001	0.114	0.106	0.210	
500	0.98	0.974	0.025	0.086	0.772	0.062	0.973	0.048	0.028	0.301	0.064	0.982	0.116	0.114	0.208	
500	0.95	0.943	0.026	0.088	0.754	0.061	0.943	0.047	0.038	0.314	0.063	0.952	0.115	0.112	0.204	
500	0.90	0.894	0.028	0.078	0.694	0.065	0.894	0.047	0.041	0.290	0.069	0.902	0.112	0.101	0.203	
500	0.80	0.794	0.032	0.081	0.613	0.065	0.794	0.050	0.048	0.264	0.069	0.805	0.110	0.101	0.200	
Panel B: Linear trend model																
200	1.00	0.963	0.058	0.263	0.710	0.075	0.969	0.107	0.062	0.218	0.078	0.989	0.169	0.109	0.171	
200	0.98	0.952	0.054	0.208	0.624	0.081	0.959	0.086	0.053	0.204	0.085	0.972	0.174	0.119	0.170	
200	0.95	0.924	0.052	0.173	0.591	0.081	0.931	0.079	0.046	0.192	0.088	0.954	0.177	0.107	0.168	
200	0.90	0.877	0.052	0.152	0.529	0.065	0.884	0.080	0.057	0.189	0.068	0.906	0.173	0.103	0.162	
200	0.80	0.775	0.057	0.132	0.452	0.070	0.781	0.082	0.057	0.180	0.077	0.801	0.172	0.106	0.170	
500	1.00	0.986	0.028	0.152	0.850	0.057	0.987	0.081	0.026	0.323	0.056	1.003	0.115	0.103	0.208	
500	0.98	0.971	0.026	0.104	0.804	0.059	0.974	0.047	0.030	0.309	0.062	0.981	0.116	0.106	0.205	
500	0.95	0.941	0.028	0.109	0.777	0.058	0.943	0.048	0.048	0.295	0.061	0.951	0.113	0.104	0.210	
500	0.90	0.891	0.029	0.094	0.724	0.062	0.893	0.049	0.053	0.306	0.068	0.903	0.111	0.100	0.195	
500	0.80	0.791	0.032	0.090	0.638	0.066	0.792	0.051	0.052	0.273	0.074	0.805	0.109	0.102	0.207	

Note: HD is the hybrid estimator. FD is the first differencing estimator. PH is the Phillips-Han (2008) estimator. T-test and J-test sizes are for 5 percent level. Power of the t-test is computed for the null of $H_0 : \alpha = \alpha_0 - 0.05$. Additional results are available in the online appendix.






Figure 1: Distribution of the t-statistic for the largest autoregressive root in the intercept-only model with $\alpha^0 = 1$. See Table 1 and the text for more details.